



9/25/2003

1/9

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

SEP 25 2003

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

Mr. Henry E. Rohland  
EPA Registration Manager  
Lebanon Seaboard Corporation  
1600 East Cumberland Street  
Lebanon, PA 17042

Dear Mr. Rohland:

Subject: Lebanon Country Club® With Ronstar®  
EPA Registration No. 961-340  
Lebanon Country Club® With Ronstar®  
EPA Registration No. 961-371  
Par Ex® Fertilizer Plus 1.00 Ronstar®  
EPA Registration No. 961-379  
Fertilizer Plus 1.5% Ronstar® Par Ex® Professional  
Products  
EPA Registration No. 961-380  
Slow Release Fertilizer Plus Ronstar® Par ex®  
Professional Products  
EPA Registration No. 961-382  
Applications and Letters Dated July 29, 2003, Requests  
To Amend Registrations To Comply with PR Notices  
2001-1 and 2001-6, Minor Editorial Changes per PR  
Notices and Marketing Statements; Your Re-submissions  
of Labeling and Revised Confidential Statements of  
Formula (CSF) Submitted with Your Letter Dated  
September 15, 2003

The resubmissions of labeling associated with the subject applications have been reviewed and found to be acceptable amendments under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, provided that you:

1. Either delete the descriptors "excellent" and "selective" where they occur in the resubmitted labeling for these pesticide products, or for "selective" clearly describe the meaning of this word in context with the associated labeling. As it occurs in the "General Information" sections, the claim could be written as: ".....and a selective preemergence herbicide for the control of broadleaf weeds and annual grasses, listed on and applied as described on this label without harm to the ornamental turf grasses at the sites listed on this label". The claim that it is an "excellent" preemergence herbicide is unacceptable as it implies that it is better than others and could be misleading (See PR Notices 91-7 and 93-6).

2/9


2. Submit one copy of each of the final printed labels prior to shipment under the enclosed stamped copy of the revised labels.

If these conditions are not complied with, these registrations will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment of this product under this labeling constitutes acceptance of this condition.

A stamped copy of each of the labels is enclosed for your records.

The proposed basic Confidential Statement of Formula (CSFs submitted in EPA Form 8570-4 and dated September 9, 2003) for each of the subject pesticide products have been reviewed; and each was found acceptable under FIFRA as an amendment to the respective registration. Each CSF has been placed in its respective record. The CSF dated September 9, 2003 replaces any existing CSF for the respective product.

Sincerely yours,



Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505C)

Enclosures (5 Stamped Labels)

3/9

# LEBANON COUNTRY CLUB® WITH RONSTAR®

**FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY, TURF AND LANDSCAPE PERSONNEL ONLY.  
NOT FOR USE ON HOME LAWNS**

**ACTIVE INGREDIENT:**

Oxadiazon ..... 1.73%

**OTHER INGREDIENTS:** ..... 98.27%

**TOTAL** ..... 100.00%

CAS No. 19666-30-9

EPA Reg. No. 961-340

Ronstar® is a registered trademark of Bayer AG.

(L) (D)

EPA Est. 961-PA-01:961-IL-01

Superscript used is first letter of lot number.

### KEEP OUT OF REACH OF CHILDREN WARNING - ADVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

See back panel for additional Precautionary Statements.

**READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.**

**Before buying this product, read the Warranty Section elsewhere on this label.**

**If terms are not acceptable, return the unopened container to the place of purchase at once.**

### GUARANTEED ANALYSIS

Total Nitrogen (N) ..... X.X%

X.X% Ammoniacal Nitrogen

X.X% Water Insoluble Nitrogen

X.X% Urea Nitrogen

X.X% Other Water Soluble Nitrogen

Available Phosphate (P<sub>2</sub>O<sub>5</sub>) ..... X.X%

Soluble Potash (K<sub>2</sub>O) ..... X.X%

Nutrient Sources: (Guaranteed Analysis and Nutrient Sources to be supplied based on fertilizer analysis used by manufacturer.)

**MANUFACTURED BY:**

**LEBANON SEABOARD CORPORATION**

1600 EAST CUMBERLAND STREET

LEBANON, PA 17042

www.lebsea.com

**NET WEIGHT: XX LBS. (XX.X kg)**

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

SEP 25 2003

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

961-340

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
WARNING**

Causes skin and eye irritation. Harmful if inhaled. Do not get on skin or on clothing. Avoid contact with eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Applicators and other handlers must wear coveralls over short-sleeved shirt and short pants, waterproof gloves, a chemical-resistant apron when cleaning equipment, protective eyewear, chemical resistant headgear for overhead exposure and chemical resistant footwear plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash body thoroughly and put on clean clothing.

**FIRST AID**

- SWALLOWED:** • Call a poison control center or doctor immediately for treatment advice.
  - Have person sip a glass of water if able to swallow.
  - Do not induce vomiting unless told to do so by a poison control center or doctor.
  - Do not give anything by mouth to an unconscious person.

**IF ON SKIN:** • Take off contaminated clothing.

**OR CLOTHING:** • Rinse skin immediately with plenty of water for 15-20 minutes.  
• Call a poison control center or doctor for treatment advice.

**IF IN EYES:** • Hold eye open and rinse slowly and gently with water for 15-20 minutes.  
• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.  
• Call a poison control center or doctor for treatment advice.

**IF INHALED:** • Move person to fresh air.  
• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.  
• Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Emergency Medical Information call (888) 208-1368.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. RONSTAR® should not be used in areas where irrigation or rainfall results in the direct or indirect (via ditches and canals) contamination of surface waters through dissolved runoff or erosion of soil particles to which oxadiazon is absorbed.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry to treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls over short-sleeved shirt and short pants, waterproof gloves, chemical-resistant footwear plus socks and chemical-resistant headgear for overhead exposure.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store this product only in its original container in a cool, dry, secured storage area out of reach of children. In case of spill, sweep up, and use according to directions or dispose of as follows. Do not store in or around the home.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary land fill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### GENERAL INFORMATION

This product is a combination product consisting of a fine quality fertilizer and an excellent selective preemergence herbicide for the control of many broadleaf weeds and annual grasses in ornamental turf, and on a wide variety of woody ornamentals. Weeds are controlled by killing the young weed seedlings as they come in contact with the herbicide during germination.

### USE PRECAUTIONS

DO NOT apply through any type of irrigation system. Avoid contamination of water intended for irrigation and domestic use.

DO NOT cut treated vegetation for forage or hay nor graze domestic animals on treated areas for one year following application.

DO NOT apply to turf with heavy thatch accumulation or to turf which has been affected by disease or nematode activity.

DO NOT use on putting greens or tees because of the varying cultural practices used.

DO NOT apply to Dichondra or Centipedegrass.

DO NOT apply to Bentgrass turfs mowed at less than 3/8 inch.

For St. Augustine and Bentgrass turf, do not exceed the rate of 4 lbs. per 1,000 sq. ft. (3 lbs. active ingredient per acre).

May cause discoloration of Bentgrass, Bermudagrass and St. Augustinegrass. This discoloration is temporary and normally outgrown within 2 to 3 weeks.

DO NOT apply to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment.

Avoid contact with flowers and shrubs.

### TURF

For use in established perennial Bluegrass, Bentgrass, Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum, Tall Fescue, and Zoysia turf; such as fairways, parks, golf courses and lawns. This product controls Crabgrass (*Digitaria* spp.), Annual Bluegrass (*Poa annua*) and Goosegrass (*Elyusine indica*). Also controls Florida Pusley (*Richardia scabra*), Oxalis (*Oxalis* spp.), Stinging Nettle (*Urtica dioica*), Carpetgrass (*Axonopus officinis*) and Pigweed (*Amaranthus* spp.).

Rake turf area free of leaves, mowing if necessary, so granules reach soil surface. Measure area to be treated and uniformly apply required amount, using a fertilizer spreader. Cross-treating with one-half required amount in each direction will give best results. **DO NOT apply to wet turf.** If rain is not expected shortly after application, sprinkling turf thoroughly will increase effectiveness.

### AMOUNT TO USE

(NOTE: Application rates listed can be shown on actual label as lbs. per acre and/or lbs. per 1,000 sq. ft.)

**CRABGRASS AND GOOSEGRASS CONTROL:** For preemergent control, apply [175 lbs. per acre (3 lbs. active ingredient) or 4 lbs. per 1,000 sq. ft.] in early Spring prior to seed germination. Application made after March or April may not be fully effective.

**ANNUAL BLUEGRASS CONTROL:** For preemergent control, apply [230 lbs. per acre (4 lbs. active ingredient) or 5 1/2 lbs. per 1,000 sq. ft.] in late Summer to early Fall or in early Spring prior to seed germination.

**PIGWEED, FLORIDA PUSLEY, OXALIS, STINGING NETTLE AND CARPETGRASS CONTROL:** For preemergent control of these weeds, apply [230 lbs. per acre (4 lbs. active ingredient) or 5 1/2 lbs. per 1,000 sq. ft.] in early Spring prior to weed seed germination.

For **ST. AUGUSTINEGRASS** turf, **DO NOT exceed the rate of [175 lbs. per acre (3 lbs. active ingredient) or 4 lbs. per 1,000 sq. ft.]**.

### WOODY ORNAMENTAL SHRUBS, VINES AND TREES APPLICATION

Apply this product anytime during the year, prior to weed seed germination, for weed control. Can be applied to both newly transplanted and established ornamentals. Application can be made to actively growing or dormant ornamentals. Rainfall or overhead irrigation after application will improve weed control activity of Ronstar®. **DO NOT apply to wet foliage or under conditions in which granules will collect on leaves.**

### AMOUNT TO USE

Ground Application: Apply this product broadcast to ornamentals at [175 lbs. of product per acre (3 lbs. active ingredient) or 4 lbs. per 1,000 sq. ft.] depending on the weeds to be controlled, as listed below. Remove existing weed growth before application.

Application of [175 lbs. per acre (3 lbs. active ingredient) or 4 lbs. per 1,000 sq. ft.], is effective in controlling annual grasses and broadleaf weeds such as:

**ANNUAL BROADLEAF WEEDS**

Bittercress	Common Purslane	Golden Ragwort	Petty Spurge	Spotted Catsear
Carpetweed	Evening Primrose	Lambsquarters	Redroot Pigweed	Swinecress
Common Groundsel	Galinsoga	Pennsylvania Smartweed	Speedwell	Yellow Woodsorrel

**ANNUAL GRASSES**

Annual Bluegrass	Barnyardgrass	Crabgrass	Fall Panicum	Green Foxtail
------------------	---------------	-----------	--------------	---------------

**NOTE:** For the control of Liverwort in California, Goosegrass and Prostrate Spurge, use [230 lbs. per acre (4.0 lbs. active ingredient) or 5 1/4 lbs. per 1,000 sq. ft.].

**CROP TOLERANCE:** Recommended for use on a wide variety of container and field grown woody ornamental shrubs, vines and trees. May be used at the recommended rates on the following ornamental species:

**CONTAINER GROWN**

<b>SPECIES</b>	<b>VARIETIES</b>	<b>SPECIES</b>	<b>VARIETIES</b>
<b>Agapanthus</b>	Africanus	<b>Holly</b>	Dwarf Okinowan, Convexa, Rotundifolia, Hetzi, Helleri, Burfordi, English, Hellus, Dwarf Yaupon, Nellie R. Stevens, Boxleafed, Green Island, Japanese
<b>Andromeda</b>	Japanese		
<b>Arborvitae</b>	American, Eastern, Oriental, Aureanana, Globe, Pyramidal, Woodward	<b>Jasmine</b>	Yellow Star, Star
<b>Arctostaphylos</b>	Fred Oelher	<b>Juniper</b>	Barharbor, Wiltoni, Torulosa, Pfitzeriana Compacti, Pfitzeriana Aurea, Torulosa Variegata, Pfitzeriana Glauca, Gulf Stream, Tam, Armstrongii, Shore, Andora, San Jose, Blue Haven, Stricta, Jade River, Broadmoor, Meyeri, Rocket, Plumosa, Royal Beauty, Praecox, Greek, Apiculata, Blackburn, Plumosa Compacta Procumbens, Blue Pfitzer
<b>Azalea</b>	Hino Red, Red, Boudoir, Snowball, Karen, Kaempp, Stuartonian, Mollishybrid, Sweetheart Supreme, Salmon Beauty, Caroline Gable, Louise, Stewartstonian, Rosebud		
<b>Barberry</b>	Redleaf, Marshall Red, Darwinii, Irwinii, Purpleleaf	<b>Justica</b>	Brandegeana
<b>Boxwood</b>	Common, Japanese	<b>Leucothoe</b>	Catesbaei, Axillaris
<b>Callistemon</b>	Bottlebrush	<b>Myrtle</b>	Compacta Noellii
<b>Camellia</b>	Glen #40, Sasanqua, Colonel Fiery	<b>Magnolia</b>	Dr. Merrilli, Samuel Somers, Grandiflora
<b>Ceanothus</b>		<b>Mahonia</b>	Oregon Grape-Holly
<b>Cedar</b>	Atlas, Portoford (Allumii)	<b>Osmanthus</b>	Fortunes
<b>Cotoneaster</b>	Paineyi, Bearberry, Coral Beauty, Greyleaf, Cranberry, Silverleaf, Peking, Bright Bead, Dammeri Skolgholm, Microphylla, Lowfast, Glaucophylla	<b>Periwinkle</b>	Vinca Minor
		<b>Photinia</b>	Frazerii
<b>Cypress</b>	Monterey	<b>Pine</b>	Slash, Japanese Black, Montana, Scotch, Digger, Shore, Thumbergii
<b>Dogwood</b>	Red Osier		
<b>Douglas Fir</b>	Pseudotsugamenziesii	<b>Pittosporum</b>	Variegated, Japanese
<b>Eucalyptus</b>	Sideroxylon, Nicholli	<b>Podocarpus</b>	Macrophylla
<b>Euonymus</b>	Greenlane, Microphylla Variegata, Emeraldgaety, Dwarf, Gold, Spot, Golden Green, Silver King, Wintercreeper Vegetus, Emerald Cushion, Spreading	<b>Privet</b>	Japanese, Waxleaf, Glossy, Vicary, Golden Vicary, Curly Leaf
		<b>Pyracantha</b>	Victory, Scarlet Firethorn, Wyattii, Graberri
<b>Forsythia</b>	Lynwood Gold	<b>Quercus</b>	
<b>Gardenia</b>	Amy Yashioka, August Beauty, Mystery	<b>Raphiolepis</b>	India Hawthorn
<b>Ginko</b>	Biloba	<b>Rhododendron</b>	Cunningham, Blue Diamond, Snow, Christmas, Chionoides, 'PJM', Dr. Rutgers, Coral Bells, President Clay, Summer Snow, Bigleaf Winter Creeper, Charlotte Armstrong
<b>Heath</b>			
<b>Heather</b>	Mrs. Maxwell	<b>Star Jasmine</b>	Yellow Star Jasmine
<b>Hedera</b>	English Ivy	<b>Syzygium</b>	Paniculatum
<b>Honeysuckle</b>	Purpleleaf, Purpurea	<b>Viburnum</b>	Sandankwa, Laurenstinus, David
		<b>Xylosma</b>	Congestum

**NEWLY-TRANSPLANTED**

SPECIES	VARIETIES
Andromeda	Japanese
Arborvitae	Globe, Woodward
Azalea	Hino Red, Louise, Rosebud, Stewartsonian
Barberry	Marshall Red, Darwinii, Irwinii, Purpleleaf
Cedar	Portoford (Alkumii)
Cotoneaster	Cranberry, Peking, Coral Beauty, Bright Bead, Paineyi, Dammeri, Skolgholm, Microphylla, Glaucophylla
Cypress	Monterey
Dogwood	Red Osier
Euonymus	Greenlane, Emerald Cushion, Spreading
Forsythia	Lynwood Gold

SPECIES	VARIETIES
Holly	English, Japanese, Rotundifolia, Helleri
Juniper	Plumosa, Plumosa Compacta, Old Gold, Pfizeriana Aurea, Torulosa, Armstrongii, Tam Wiltoni, Procumbens
Leucothoe	Axillaris
Magnolia	Dr. Merrilli
Mahonia	Oregon Grape-Holly
Pine	Slash
Privet	Japanese, Waxleaf
Pyracantha	Graberii
Rhododendron	Snow, Blue Diamond, Chionoides, 'PJM', Dr. Rutgers
Rose	Summer Snow, Bigleaf Winter Creeper, Charlotte Armstrong

**ESTABLISHED, FIELD-GROWN**

SPECIES	VARIETIES
Arborvitae	Pyramidal
Ash	Green
Azalea	Hinodigeri
Barberry	Red, Mentor
Birch	Clump, Paper
Camellia	Tiffany, Marie Bracey, Elizabeth Dowd, Tom Knudsen, Crimson Robe, R.D. Wheeler
Cotoneaster	Lowfast, Skolgholm
Crabapple	Flowering, Torringo
Dogwood	Redstemmed, Red, Flowering
Euonymus	Burning Bush, Dwarf Burning Bush, Jewell
Forsythia	Showy Border, Lynngold, Lynwood Gold, Goldenbell
Holly	Boxleafed, Green Island, Japanese
Honeysuckle	Zabeli
Juniper	Plumosa, Mint Julip, Arcardia, Variegata, Plumosa Compacta, Andora, Wiltoni, Hetzi
Lilac	Persian, Dwarf Korean
Maple	Amur
Oak	Red
Periwinkle	Vinca Minor

SPECIES	VARIETIES
Pine	Austrian, Scotch, Norway, Stone, Japanese Black, Loblolly, Slash, Eastern White
Pittosporum	Undulatum
Privet	California, Japanese
Rhododendron	Rosebud
Rose	Texas Centennial, Sutters Gold, Show Girl
Russian Olive Spruce	Black Hills, Norway, Serbian
Taxus	Taunton, Spreading Yew, Hicks
Tsuga	Canadian Hemlock
Viburnum	European Cranberry Bush
<u>Xylosma Ground Covers</u>	
Carpobrotus	Edule
Osteospermum	
Sedum	Album
Algerian Ivy	
Vinca Major	
English Ivy	
Pachysandra	Terminalis
Ajuga	Reptans
Euonymus	Fortunei Radicans

**NEWLY-TRANSPLANTED, FIELD-GROWN**

SPECIES	VARIETIES
Camellia	Tiffany, Marie Bracey, Elizabeth Dowd, R.D. Wheeler, Tom Knudsen, Crimson Robe
Dogwood	Redstemmed, Red, Flowering
Forsythia	Showy Border, Lynwood Gold, Goldenbell
Hedera	English Ivy
Holly	Green Island, Japanese
Juniper	Plumosa, Andora, Wiltoni, Hetzi
Oak	Red

SPECIES	VARIETIES
Periwinkle	Vinca Minor
Pine	White Austrian, Eastern White
Privet	California, Japanese
Rhododendron	Rosebud
Russian Olive Spruce	Black Hills, Norway, Serbian
Taxus	Hicks
Tsuga	Canadian Hemlock

**ADDITIONAL APPLICATIONS**

For continued weed control, additional applications may be made 60 to 120 days after the previous application in the following situations:

**2 REPEAT TREATMENTS, CONTAINER GROWN, ESTABLISHED**

SPECIES	VARIETIES
Abelia	Grandiflora
Common Boxwood	
English Ivy	
Holly	Helleri
Juniper	Tam
Pine	Slash
Rhododendron	Obtusa, Kurume, Formosa, Kurume Snow

**CONTAINER GROWN, NEWLY-TRANSPLANTED**

SPECIES	VARIETIES
Common Boxwood	
English Ivy	
Juniper	Tam
Pine	Slash
Rhododendron	Obtusa, Kurume Formosa

**FIELD GROWN, ESTABLISHED AND NEWLY-TRANSPLANTED**

SPECIES	VARIETIES
Holly	Nellie R. Stevens, Dwarf Burford, Burford

**3 REPEAT TREATMENTS, CONTAINER GROWN, ESTABLISHED**

SPECIES	VARIETIES
Abelia	Grandiflora
Holly	Helleri, Rotundifolia
Rhododendron	Kurume Formosa, Kurume Snow

**GREENHOUSE GROWN, ESTABLISHED**

SPECIES	VARIETIES
Rose	Sonia, Forever Yours, Coed, Golden Wave

**FOREST NURSERIES, SEEDLINGS AT LEAST 4 WEEKS OLD**

SPECIES	VARIETIES
Pine	Loblolly, Slash, Eastern White

Apply this product with a properly calibrated granular applicator that will apply the granules uniformly (uniform application is essential for satisfactory weed control). Calibrate the applicator prior to use, according to the manufacturer's directions, and check frequently during the application to be certain that the equipment is working properly.

**SUGGESTED SPREADER SETTINGS**

SPREADER	SPEED		Width of Coverage	Rate Table and Spreader Settings	
	GROUND MPH	PTO RPM		4#/1000	5'/./1000
				175#/Acre	230#/Acre

[Note: Registrant/Subregistrant may insert appropriate spreader setting information based on desired spreaders and recommended use rates.]

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

**WARRANTY**

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Manufacturer, and Buyer assumes the risk of any such use.

Country Club® is a registered trademark of Lebanon Seaboard Corporation.



**Advertising claims that may be presented on the label**

A selective preemergent herbicide for the control of Crabgrass, Goosegrass and other Annual Weeds in Turf and Woody Ornamentals.

**Additional Labeling Requirements:**

If the fertilizer guarantee contains iron, the following statement must appear:

*This product contains the secondary nutrient iron. Iron may stain concrete and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash off with water.*

If the recommended use rate applies less than 1 pound nitrogen per 1,000 sq. ft. per application, the following statement must appear:

*Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.*

For products distributed in the State of Washington, the following statement must appear:

*Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-lebsea.com>*