

JUL 7 2003

Mr. Henry E. Rohland
Lebanon Seaboard Corporation
1600 East Cumberland Street
Lebanon, PA 17042

Dear Mr. Rohland:

Subject: Product Reregistration, First-Aid and Disposal Instructions
Lebanon Lawn Crabgrass Preventer with Fertilizer
EPA Registration No. 961-346
Your Submission Dated January 4, 2002

The label amendment referred to above, submitted in accordance with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable provided that you:

1. Within the First Aid section, place the skin exposure ("If on skin") first aid statements first, followed by the eye exposure and oral ingestion first aid statements.
2. Submit one (1) copy of your final printed labeling before you release the product for shipment.

This amended label supersedes all previously accepted labels. A stamped copy of the label is enclosed for your records. If you have any questions, about this letter, contact me at (703) 305-6224.

Please note, EPA has determined that you have fulfilled the requirements of the Trifluralin RED for the subject product. However, we are unable to reregister the product under FIFRA 4(g)(2)(c) at this time, since it also contains the active ingredient, benefin, for which a reregistration eligibility decision has not yet been made. You will be notified of the Agency's reregistration decision for products containing benefin when the Benefin RED is completed.

Sincerely yours,

Joanne I. Miller
Product Manager 23
Herbicide Branch
Registration Division (7505C)

Enclosure

CURRENCES

SIMBOL	7505C							
SURNAME	JIMiller							
DATE	Jul 7, 2003							

2 2 5

LEBANON LAWN CRABGRASS PREVENTER WITH FERTILIZER

ACTIVE INGREDIENTS:

Benefin	0.770%
Trifluralin	0.385%
OTHER INGREDIENTS	98.845%
TOTAL	100.000%

EPA Reg. No. 961-346
 Benefin CAS. No. 1861-40-1
 Trifluralin CAS. No. 1582-09-8

(L) (D)
 EPA Est. 961-PA-01:961-IL-01
 Superscript used is first letter of lot number.

GUARANTEED ANALYSIS

Total Nitrogen (N)	XX%
X.X% Ammoniacal Nitrogen	
X.X% Water Insoluble Nitrogen	
X.X% Urea Nitrogen	
X.X% Other Water Soluble Nitrogen	
Available Phosphate (P ₂ O ₅)	XX%
Soluble Potash (K ₂ O)	XX%

Nutrient Sources: (*Guaranteed Analysis and Nutrient Sources to be supplied based on fertilizer analysis used by manufacturer.*)

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for additional Precautionary Statements.

NOTICE: Read the entire label. Use only according to label directions.
Before buying this product, read the Warranty Section elsewhere on this label.
If terms are not acceptable, return the unopened container at once.

MANUFACTURED BY:
LEBANON SEABOARD CORPORATION
1600 EAST CUMBERLAND STREET
LEBANON, PA 17042
www.lebsea.com

NET WEIGHT: 40 LBS. (18.14 kg)
COVERS: 10,000 Sq. Ft.

**ACCEPTED
 with COMMENTS
 in EPA Letter Dated**

JUL 7 2003

**Under the Federal Insecticide,
 Fungicide, and Rodenticide Act
 as amended, for the pesticide
 registered under EPA Reg. No.
961-346**

375

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if absorbed through skin. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

Personal Protective Equipment:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Protective eyewear

User Safety Requirements: Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

- Users should wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF IN EYES: • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

- Call a poison control center or doctor for treatment advice.

IF SWALLOWED: • Call a poison control center or doctor immediately for treatment advice.

- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN • Take off contaminated clothing.

OR CLOTHING: • Rinse skin immediately with plenty of water for 15-20 minutes.

- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contradict the use of gastric lavage.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater, marine and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all directions carefully before applying this product.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food, feed, seeds or other pesticides.

DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

In case of emergency endangering health or the environment involving this product, call Chemtrec day or night at 1-800-424-9300.

GENERAL INFORMATION

This turf fertilizer product contains TEAM, a preemergence herbicide which is applied to established turfgrasses for the control of the winter annual grass: Annual Bluegrass (*Poa annua*); and summer annual grasses: Smooth and Hairy Crabgrass, Goosegrass (silver crabgrass or crowfootgrass), Barnyardgrass (watergrass) and Green and Yellow Foxtail. This product also provides partial control of prostrate spurge (*Euphorbia humistrata*), spotted spurge (*Euphorbia maculata*), creeping spurge (*Euphorbia serpens*), creeping woodsorrel (*Oxalis corniculata*) and yellow woodsorrel (*Oxalis stricta*).

475

This Product Is Recommended For Use On The Following Established Warm and Cool Season Perennial Turfgrass Species:

- Established Cool Season Turfgrasses** - Bentgrass Fairways, Kentucky Bluegrass, Perennial Ryegrass, Fescues
- Established Warm Season Turfgrasses** - Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass, Zoysiagrass.

This product should be applied one to two weeks prior to the anticipated germination of the target annual weeds. Apply using a properly calibrated granular fertilizer spreader which will uniformly distribute the product over the target area. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur. Optimum weed control performance will be obtained if treated areas are irrigated soon after application. This product will not harm most nearby ornamentals.

In areas with extended growing seasons or where desired, a second application at the recommended rate can be made 8 to 10 weeks following the initial application on cool season turfgrasses and 10 to 12 weeks following the initial application on warm season turfgrasses.

GENERAL PRECAUTIONS AND RESTRICTIONS

Do Not Ship or Store with Food, Feeds, Drugs or Clothing.

NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION OR FOR RESEARCH PURPOSES.

Application Restrictions: Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Entry Restrictions: Do not allow people or pets to enter the treated area until dusts have settled. If watering-in is required after the application, do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

Reseeding: Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Do not reseed treated areas for a minimum of 8 weeks after a single application of 4 lbs. per 1,000 sq. ft. (174 lbs. per acre) or 12 to 16 weeks after a single application of 6 lbs. per 1,000 sq. ft. (260 lbs. per acre). For split applications, do not reseed treated areas for 8 to 10 weeks after the last 3 lbs. per 1,000 sq. ft. rate (130 lbs. per acre) or 10 to 12 weeks after the last 4 lbs. per 1,000 sq. ft. rate (174 lbs. per acre).

Use caution to confine product to intended application area, as the active ingredients may stain concrete, plastic or other materials.

Do not apply this product to golf course putting greens, dichondra, or newly sprigged areas of Bermudagrass, St. Augustinegrass, Centipedegrass, or Zoysiagrass.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application.

APPLICATION INSTRUCTIONS

Broadcast Application Rates:	Cool Season Turfgrasses	Warm Season Turfgrasses
Summer Annual Grass Weeds (Spring Application)	130 to 260 lbs./acre 3 to 6 lbs./1000 sq. ft.	260 lbs./acre 6 lbs./1000 sq. ft.
Winter Annual Grass Weeds (Fall Application)	130 to 174 lbs./acre 3 to 4 lbs./1000 sq. ft.	260 lbs./acre 6 lbs./1000 sq. ft.

Split Applications for Extended Control and Lawn Care Operator Programs:

As an alternative to single application programs, this product may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time.

Apply this product at 3 lbs. plus 3 lbs. per 1,000 sq. ft. (130 lbs. plus 130 lbs. per acre) for Cool Season Turfgrasses and in the northern half of the Warm Season Turfgrass area of the U.S. or 4 lbs. plus 4 lbs. per 1,000 sq. ft. (174 lbs. plus 174 lbs. per acre) in the southern half of the Warm Season Turfgrasses area of the U.S. The second application should be made 8 to 10 weeks following the 3 lbs. per 1,000 sq. ft. rate (130 lbs. per acre) and 10 to 12 weeks following the 4 lbs. per 1,000 sq. ft. rate (174 lbs. per acre).

SUGGESTED SPREADER SETTINGS

[Note: Registrant/Subregistrant may insert appropriate spreader setting information based on desired spreaders and recommended use rates as listed above.]

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

WARRANTY

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Manufacturer, and Buyer assumes the risk of any such use.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

575

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

*Trademark of Dow AgroSciences LLC.

Advertising claims that may be presented on the label

- A Granular Fertilizer Containing TEAM* for the Preemergence Control of Crabgrass, Foxtail, Goosegrass, *Poa Annua* and Other Annual Weed Grasses in Established Turfgrass.

Additional Labeling Requirements:

If the fertilizer guarantee contains iron, the following statement must appear:

This product contains the secondary nutrient iron. Iron may stain concrete and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash off with water.

If the recommended use rate applies less than 1 pound nitrogen per 1,000 sq. ft. per application, the following statement must appear:

Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

For products distributed in the State of Washington, the following statement must appear:

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-lebsea.com>