

NET WEIGHT: 20 LBS. (XXX Kg.)

COVERS: XXXX Sq. Ft.

# LEBANON LAWN FOOD WITH INSECT CONTROL

Contains 1.0% Dursban®

# 19-4-6

Controls Chinch bugs, Sod webworms (Lawn moths), Ants, Brown dog ticks, Crickets, Cutworms, Earwigs, Sowbugs (Pillbugs), Fleas, Chiggers, Clover mites, Grasshoppers, Armyworms and Hyperodes (Turfgrass) weevils infesting home lawns and other ornamental turf areas. Also controls grubs (larvae) of Japanese beetles and European chafer.

**ACTIVE INGREDIENT:**

Chlorpyrifos 0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl)phosphorothioate\*

1.00%

**INERT INGREDIENTS:**

99.00%

**TOTAL:**

100.00%

\*Dursban® - a trademark of DowElanco

(L) (D)

EPA Reg. No. 961-357

EPA Est. 961-PA-01:961-IL-01

Superscript used is first letter of lot number.

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
See Back Panel for Additional Precautionary Statements

MANUFACTURED BY  
Lebanon Chemical Corporation  
Lebanon, PA 17042

**ACCEPTED**  
**APR 18 1995**  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 961-357

Page 1 of 4 Pages

961-357

01/16/95

1995

## FERTILIZER ANALYSIS

Total Nitrogen (N)	19%
Available Phosphoric Acid ( $P_2O_5$ )	4%
Soluble Potash ( $K_2O$ )	6%

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

May be harmful if swallowed.

Do not take internally. Avoid inhaling dust. Avoid eye contact. Wash thoroughly after handling. Wash contaminated clothing before reuse. Keep away from food, feedstuffs and domestic water supplies.

### FIRST AID

**IF SWALLOWED:** Call a physician or Poison Control center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF INHALED:** Remove victim to fresh air if symptoms of cholinesterase inhibition appear, and get medical attention immediately.

**IF ON SKIN:** Wash thoroughly with soap and water.

**IF IN EYES:** Flush with plenty of water. Get medical attention if irritation persists.

**NOTE TO PHYSICIAN:** *Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine, only by injection, is an antidote.*

### ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. **APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL**

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
**APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.**

**NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION OR FOR RESEARCH PURPOSES.**

Feeds your lawn the essential food elements required for good growth, while controlling the following lawn and home pests: Chinch bugs, Sod webworms (Lawn moths), Ants, Brown dog ticks, Crickets, Cutworms, Earwigs, Sowbugs (Pillbugs), Fleas, Chiggers, Clover mites, Grasshoppers, Armyworms and Hyperodes (Turfgrass) weevils. Also for control of larvae of Japanese beetles and European chafer.

For control of all insects named above, except Hyperodes weevils and larvae of Japanese beetle and European chafer, apply at the rate of 2 lbs. 5 oz. per 1,000 sq. ft. to turf or lawn area. Treat when insects first appear, re-treat as needed.

For control of Hyperodes (Turfgrass) weevils, make a double application using 4 lbs. 10 oz. per 1,000 sq. ft. (Use 2 successive applications per table directions.) Treat suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for best time to treat in your area.

For control of larvae of Japanese Beetle and European Chafer, apply at the rate of 4 lbs. 10 oz. per 1,000 sq. ft. (Use 2 successive applications per table directions). Treat when white grubs are young and actively feeding near the soil surface, usually during late July and August, or as recommended by your local Agricultural Extension Service Specialist.

Immediately after all applications, wash granules into the soil with 1/2 to 1 inch of water.

### SUGGESTED SPREADER SETTINGS

	<u>2 Lbs.</u>	<u>4 Lbs.</u>		<u>2 Lbs.</u>	<u>4 Lbs.</u>
Show 'N Tell	5 1/2	6 1/2	Cyclone	4 1/2	5
Scott	4 1/2	6 1/2	Scott Rotary	F	1
Central	4 1/2	6	Ortho	3 1/2	5
Jackson	4 1/2	6	Ortho Rotary	3 1/2	4 1/2
Glorion	4 1/2	5 1/4			

**NOTE:** Spreader settings are approximate. Check your spreader on 100 sq. ft. and adjust to apply 3.7 oz. per 100 sq. ft. Rate of walk, moisture and spreader condition can vary the settings suggested.

## STORAGE AND DISPOSAL

**STORAGE:** Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food, feed, seeds or other pesticides.

**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

## WARRANTY

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use.

Page 4 of 4 Pages

961-357

01/10/95