

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

APR 2 4 2009

Mrs. Sylvia Whitman State Registration Coordinator Lebanon Seaboard Corporation 1600 East Cumberland Street Lebanon, PA 17042

RE: Notification of Alternate Brand Name: Lebanon Pro Brand Fertilizer with Dimension

0.09% Pre & Postemergence Annual Grassy Weed Control

Registration Number: 961-359

Submission Dated: February 23, 2009

Dear Mrs. Whitman:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated, February 23, 2009, for the product, Lebanon Fertilizer with Dimension Crabgrass Control. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely yours,

Linda Arrington

Notifications & Minor Formulations Team Leader Registration Division (7505P)

Office of Pesticide Programs

			ř		. .				2/4
SEPA	U Environmental	nited States	_	ncy	Form App	roved ✓	Registrat Amendm Other	ion	OPP Identifier Number
		Application	on for F	esticid	e - Sect	ion	1		
1. Company/Product Number	961-359			2. EPA P	roduct Mana	reger		3. Pro	posed Classification
4. Company/Product (Name) Lebanon Fertilizer with		ass Control	<u> </u>	PM#	23	•	**************************************		None Restricted
5. Name and Address of App				·			In annual an		FIFRA Section 3(c)(3)
Lebanon Seaboard C 1600 East Cumberlan Lebanon, PA 17042	orporation			(b)(i), m to:		s sim	NOTI	al in cor	mposition and labeling
Check if this	is a new address			Produc	t Name		APR	2 4 20	U9
			Sec	tion - Il	 				
Amendment - Explain Resubmission in resp Notification - Explain Explanation: Use addition Alternate Brand Name: Leba	onse to Agency letter below. nal page(s) if necessar	y. (For sectio			Agency lett "Me Too" A Other - Expl	er dat Applica Isin be	ation. alow.		
			Sect	ion - II					
1. Material This Product Will	Be Packaged In:								
Child-Resistant Packaging Yes No * Certification must be submitted	Unit Packaging Yes No If "Yes" Unit Packaging wgt.	No. per container	Water		No. per container		2. Type of C	ontainer Metal Plastic Glass Paper Other (S	specify)
3. Location of Net Contents		4. Size(s) Re	tail Contai	ner		5. Lo	cation of Labe	Direction	ns
6. Manner in Which Label is	Affixed to Product	Lithor Paper	graph r glued ciled		Other	·			
		land the second		ion - I\	1				
1. Contact Point (Complete	items directly below f	or identification				if nec	essary, to pro	cess this	application.)
Name Sylvia Whitman			Title		egistration				e No. (include Area Code)
I certify that the states I acknowledge that an both under applicable 2. Signature	y knowlinglly false or law.		d all attach						6. Date Application Received (Stamped)
	111-6)		State Re	gistration Ma	anage	r		

5. Date

Sylvia Whitman

Feb. 23, 2009



LEBANON SEABOARD CORPORATION

1600 EAST CUMBERLAND STREET • LEBANON, PENNSYLVANIA 17042

(717) 273-1685

FAX (717) 273-9466

March 5, 2009

Document Processing Desk Office of Pesticide Program (7504P) U.S. Environmental Protection Agency Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, VA 33302-4501

ATTN: Joanne Miller (Team 23)

RE: Alternate Brand Names

- EPA Reg. No. 961-359 Lebanon Pro Brand Fertilizer with Dimension 0.09% Pre
 & Postemergence Annual Grassy Weed Control
- EPA Reg. No. 961-360 Lebanon Pro Brand Fertilizer w/Dimension .13% Pre & Postemergence Annual Grassy Weed Control
- EPA Reg. No. 961-361 Lebanon Pro Brand Fertilizer with Dimension .19% Pre & Postemergence Annual Grassy Weed Control
- EPA Reg. No. 961-362 Lebanon Pro Brand Fertilizer with Barricade 0.43% Preemergence Weed Control
- EPA Reg. No. 961-365 Lebanon Pro Brand Fertilizer with Barricade 0.38% Preemergence Weed Control
- EPA Reg. No 961-367 Lebanon Pro Brand Fertilizer with Barricade 0.29%

Dear Ms. Miller:

Please find enclosed Lebanon's applications for Alternate Brand Name.

- Transmittal Document (this letter)
- EPA Form 8570-1 (6)
- Copy of each referenced label (6)

Should you have any questions, please feel free to contact me at 800/532-0090 ext. 361.

Sincerely.

State Registration Coordinator

enclosures:



FERTILIZER WITH DIMENSION® 0.09%

PRE & POSTEMERGENCE ANNUAL GRASSY WEED CONTROL

NOTIFICATION APR 2 4 2009

GUARANTEED ANALYSIS	
Total Nitrogen (N)	25%
25.0% Urea Nitrogen	
Soluble Potash (K ₂ O)	5%
Derived from: Urea, Muriate of Potash.	
Chlorine (CI) not more than	5.0%
	F699



22-42725

NOTE: Supplemental applications of fertilizer are needed to achieve a minimum 1 lb. Nitrogen per 1,000 sq. ft.

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-info-lebsea.com.

SPREADER SETTINGS	Rate in pounds 3.2	per 1,000 Sq. Ft. 4.7	These settings were calibrated and field
LebanonTurf ProScape® SS/PrizeLAWN® BF-1/SS/CBR III PennMulch® HVO/PrizeLAWN® BF-HVO Earthway Rotary Lely Lesco (letter dial/numeric dial) Scotts AccuPro 2000/SR-2000 Scotts R8A Rotary Scotts R7X Rotary Spyker Rotary Vicon	4½ K K 15 5II I / 17 K L H 4½ 24	5 N N 18 6½II K / 22 O N J 5 30	tested. Hovever, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

0.000/

Dimension® is a selective herbicide that provides control of listed annual grasse and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes.

ACTI \	/C	INIC	DE	ווח	EVI.	T٠
AUIII		IIAC	INE	ווע	CIA.	Ι.

OTHER INGREDIENTS:		<u>99.91%</u>
*This product contains 0.045 lb. of the active ingredient, dithiopyr, per 50 lb. bag.	*Cas N	o. 97886-45-8
Product protected by U.S. Patent No. 4,692,184. Other patents pending.	(L)	(D)

EPA Reg. No. 961-359

in the land of the same of

EPA Est. No. 961-PA-01:961-IL-01

*Dimension is a registered trademark of Dow AgroSciences LLC.

Superscript used is first letter of lot number.

ProScape and Mesa are registered trademarks of Lebanon Seaboard Corporation.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label.

If terms are not acceptable, return the unopened container to the place of purchase at once.

In New York State, this product may only be used by Professional Applicators at no more than 12.8 lbs. per 1000 sq. ft. (0.5 lb. active ingredient per acre) per year; and is prohibited for use in Nassau and Suffolk Counties.

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for additional Precautionary Statements.

Manufactured by:
Leoanonurf

For technical assistance or more information about our products visit www.LebanonTurf.com 1600 East Cumberland St • Lebanon, PA 17042

NET WEIGHT 5 LBS. (22.7 KG)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated dothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause aftergic reaction in some individuals.

FIRST AID

IF IN EYES:
 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 Call a poison control center or doctor for treatment advice.
 IF INHALED:
 Move person to tresh air.

• Note person to restrict.
 • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
 • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Emergency Medical Information call (888) 208-1368. For any other emergency involving this product, call day or night (800) 424-9300.

ENVIRONMENTAL HAZARDS

ENVINUENTIAL TRACARDS

This product is toxic to fish and highly toxic to other aquatic organisms including contents and shrimp. Use with care when applying to turl areas adjacent to any body of water. Drift and runoff from treated turl may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION, OR FOR RESEARCH PURPOSES.

GENERAL INFORMATION

This product is a selective herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and omamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS USE PRECAUTIONS

Turfgrass Safety

1 utilities out to be used on seeded, socided, or sprigged lawns and ornamental turts that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two movings following its seeding, sociding or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turts that are not well-established, or on those that have been weakened by weather, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turt injury.

This product should only be applied to lawns and omamental turts that are composed of the following furlgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following furlgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following furlgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following furlgrass species that have been determined to be tolerant to applications of this product. lowing established turigrasses are tolerant to this product:

Cool-Season Grasses

Entgrass, creeping (Agrostis palustris)
Bluegrass, Kentucky (Poe pratensis)
Fescue, fine" (Festuca ubra)
Fescue, till (Festuca annotinacea)
Ryegrass, perennial (Lolium perenne)

Warm-Season Grasses

Bahiagrass (Paspalum notatum) Bermudagrass (Cynodon dactylon) Buflalograss*** (Buchloe dactyloides) Carpetgrass (Axonopus affinis)

Centipedegrass (Eremochiba ophiuroides) Kikuyugrass (Pennisetum clandestinum) St. Augustinegrass (Stenotaphrum secundatum) Zoysiagrass (Zoysia japonica)

* Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turtgrass injury. Not all varieties of creeping bentgrass have been tested.

DO NOT apply this product to Colonial bentgrass (Agrostis tenuis) varieties.

"Use of this product on certain varieties of fine fescue may result in undesirable furf injury. The following fine fescue varieties have been found to be sensitive to this product 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Taljana', 'Waldorf', and 'Waldina'. Not all varieties of fine fescue have been fested.

"DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

NOTE: It is not possible to evaluate all turfgrass varieties for tolerance to Dimension. It is suggested that the user evaluate the response of turfgrass varieties to this product on a limited area, at recommended use rates, prior to initiating large-scale use. RESEEDING, OVERSEEDING, OR SPRIGGING

Reseeding, overseeding, or springing of treated areas with this product should be delayed until 12 weeks from the time of application. Reseeding, overseeding or springing before 12 weeks after application may prevent establishment of desirable turigrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. If overseeding with perennial ryegrass in the fall, overseed B weeks or later after a single application. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact. OTHER USE PRECAUTIONS

The early postermergence crabgrass control described below is limited. Postermergence applications are only effective on crabgrass if applied before crabgrass has displayed its lifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core, spike, or hydro-aerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

• DO NOT use dippings from treated turl for mulching around vegetables or fruit trees.

• Keep people and pels off treated areas until dust has settled.

APPLICATION DIRECTIONS Control of Crabgrass

Preemergence and Early Postemergence Control
This product provides "preemergence" control of α abgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turis.

It can also provide "early postermergence" control of crabgrass. Postermergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first see the crabgrass plants in the lawn or furf. The practical benefit of this product's early postermergence activity is that it can give the user a 2-4 week longer period of time to make applications to control crabgrass.

Application Frequency and Timing
This product may be applied as a single application or as a split application for crabgrass control in the spring, summer, or fall.

Spring Applications

For single applications and the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the control programs listed in Table 1 below, depending on the user's location, the furgrass mowing height, and whether the use is preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turtgrass competitiveness, and the user's location within a region.

This program should not be used for early postemer.

This program should not be used for early postemer.

- Use Program 1 (given in Table 1, below) for preemergence control at sites where the furfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early posterner gence crabgrass control.
- Use Program 2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early posternergence control of crabgrass.

 Where split fertilizer applications are recommended, a rate in Table 1 may be applied in two applications made 6-10 weeks apart and prior to crabgrass emergence.

	1: Recommended A for Crabgrass of this Product per	Application Use Rates ,* 1,000 Sq. Ft.)
PROGRAM		2
USE	PREEMERGENCE	PREEMERGENCE
(turfgrass cut)	(high)	POSTEMERGENCE
REGION		
North	3.2	6.4
Transition	4.7	9.3
South	6.4	N.R.**
Coastal South	N.R."	N.R."

acaian	M
REGION	Preemergence
North	9.3
Transition	N.R."
South	N.R."
Coastal South	N.R."

NOTIFICATION APR 24 2009

Fall Applications
This product can also be applied in the late summer or early fall (late August through November) at the use rates listed in Table 2 to provide control of crabgrass through the summer. The fall application should be followed by an appropriate spring application to provide season-long control.

**DO NOT apply more than 9.3 lbs. of this product per 1000 sq. ft. per application, and no more than 38.5 lbs. of this product per 1000 sq. ft. per year.

In New York State, this product may only be used by Professional Applicators at no more than 12.8 lbs. per 1000 sq. ft. (0.5 lb. active ingredient per acre) per year; and is prohibited for use in Nassau and Suffolk Counties. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below.

North: All other areas.

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, and coastal areas of CT, NY, and RI.

AL, AR, AZ, CA, CA, LA, MS, NC, NM, NV, OK, SC, TN, and TX.

Coastal South: HI, FL, and southern coastal areas of AL, GA, LA, MS, NC, SC, and TX.

** N.R. - This formulation is not recommended.

Tips for Improved Control

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

When treated lawn or omamental furliareas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result. Use of split (half-rale) applications spaced 6-8 weeks apart may provide improved weed control.

Control of Other Grass and Broadleaf Weeds

Spring Applications
Used as directed for grabgrass control in the spring (Table 1), this product will also control or suppress the following weeds when applied prior to their emergence:

Barnyardgrass Bluegrass (annual, roughstalk) Crowfootgrass Foxtall (yellow & green)*

Goosegrass Kikuyugrass Lespedeza (common) Oxalis (buttercup, creeping & yellow woodsorrel) Purstane (common) Ryegrass (annual & perennial)

Smutgrass Speedweil (corn) Spurge (prostrate & spotted)

* Also controlled at the Program 2 rates.

Fall Applications
Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring (Table 2), this product will also control or suppress the following weeds when applied prior to their emergence:

Bluegrass (annual, roughstalk)

Bittercress Chickweed Geranium (Carolina) Henbit

Pineappleweed Parsley-piert

APPLY this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equip ment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or omamental turd area. A more uniform application can be mad by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

APPLICATION EQUIPMENT INSTRUCTIONS

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal,

Do not contaminate water, took or feet by surveye or depoissal.

PESTICIDE STORAGE: Slore product in its original container in a cool, dry, locked place out of the reach of children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Of only reuse or refill this container. Of the fore container was the container of the contrainer of the container. On one reuse or refill this container. Of the fore container was the container of the container. On the container of t

22-42725

CONDITIONS OF SALE AND WARRANTY

The Manufacturer warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. THE MANUFACTURER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Handling, storage and use of the product by Buyer or User are beyond the control of the Manufacturer and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property or failure to tollow label directions will be assumed by the Buyer or User. IN NO CASE WILL THE MANUFACTURER OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.