96.1	360		9-24-0		red. OMB No. 20	70-0060. 4	oproval expires 2-28-8	
\$EPA	Environme	United States			Registrati	ion <sup>of</sup>	P Identifier Number	
		Applicati	ion for Pestic	ide - Sectio	n I			
. Company/Product N	umber 961-360		2. EPA Product Manager Joanne Miller		if		Proposed Classification     None     Restricted	
. Company/Product (N Lebanon Fertilizer	eme) with Dimension Cra	barass Contro	PM#	23				
Name and Address o	f Applicant <i>(Include Zh</i>	P Code)					RA Section 3(c)(3)	
ebanon Seaboa 600 East Cumbe ebanon, PA 17	erland Street		to:		imilar or identic	-	sition and labeling	
Check	f this is a new address		Prod	uct Name				
· · · · · · · · · · · · · · · · · · ·			Section -	1(				
Amendment - E	plain below.			Final printed la	bels in repsones t	•		
Resubmission in	response to Agency le	itter dated		Agency latter		ŃC	TIFICATION	
Notification - Ex	plain below.			Other - Explain	below.	SE	P 2 9 2003	
	bject to enforcement ac		Section -					
hild-Resistant Packagi			Water Soluble	Packaging	2. Type of Co	nteiner		
Yes	Yes		Yes N		Metal			
No	No		No			Plestic Glass		
Certification mus e submitted	f "Yes" Unit Packaging v	No. per wgt. container	lf "Yes" Package wgt	Na. per container		Paper Other (Speci	M	
Location of Net Cont	ents information	4. Size(s) Re	stail Container	5.	Location of Label	Directions	<u></u>	
	el is Affixed to Produc	t Litho	graph folued biled	Other _	······································			
	-		Section -	V				
Contect Point /Com	whete items directly bek	w for identificati			ecessary, to proc	ees this appl	ication.)	
Name Henry E. Rohland					(Include Area Code)			
•	at any knowlingly false able law.		ation d all attachments ti atement may be pu 3. Title	hereto are true, a Inishable by fine (	ocurete and comp or imprisonment o	inte de la f	Site Application lectived (Stomped)	
Typed Name	Z. Kella	n	EPA R 5. Date	egistration Manag	er	·	to the second	
Henry E. Rohland				<b>SEP</b> 18	2003			

EPA Form 8570-1	(Rev. 3-94) Prev	vious editions ere	obsolete.

# LEBANON FERTILIZER WITH DIMENSION® CRABGRASS CONTROL

216

NOTIFICATION SEP 2 9 2003

Dimension<sup>®</sup> is a selective herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes.

# **ACTIVE INGREDIENTS:**

*Dithiopyr	0.13%
OTHER INGREDIENTS	<u>99.87%</u>
TOTAL	100.00%

\*This product contains 0.065 lb. of the active ingredient, dithiopyr, per 50 lb. bag. Product protected by U.S. Patent No. 4,692, 184. Other patents pending. \*Dimension is a registered trademark of Dow AgroSciences LLC.

EPA Reg. No. 961-360 Cas No. 97886-45-8 (L) (D) EPA Est. 961-PA-01:961-IL-01 Superscript used is first letter of lot number.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS. Before buying this product, read the Warranty Section elsewhere on this label. If terms are not acceptable, return the unopened container to the place of purchase at once.

In New York State, this product may only be used by Professional Applicators at no more than 8.8 lbs. per 1000 sq. ft. (0.5 lb. active ingredient per acre) per year; and is prohibited for use in Nassau and Suffolk Counties.

# KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel for Additional Precautionary Statements.

# **GUARANTEED ANALYSIS**

Total Nitrogen (N)	XX%
X.X% Ammoniacal Nitrogen	
X.X% Water Insoluble Nitrogen	
X.X% Urea Nitrogen	
X.X% Other Water Soluble Nitrogen	
Available Phosphate (P2O5)	XX%
Soluble Potash (K <sub>2</sub> O)	XX%

Nutrient Sources: (Guaranteed Analysis and Nutrient Sources to be supplied based on fertilizer analysis used by manufacturer.)

MANUFACTURED BY:		
LEBANON SEABOARD CORPORATION		7 6 V 6 C V
1600 EAST CUMBERLAND STREET	. • • • •	С
LEBANON, PA 17042		000 C 277 - 02 16 - 16 10
www.lebsea.com	୯୦୦ <b>୦୫୫</b> ୫ ଜୁନ୍ତ୍ର	
	~ •	90 C 9 0 0 9 90
NET WEIGHT: XX LBS. (X.X kg)	1 U	e e e
COVERS: X,XXX Sq. Ft.		0440
		2 UNIOS
Page 1 of 5 Pages		4 h
LEBANON SEABOARD CORPORATION 961-360		του 1. α. Γ

LABEL NOTIFICATION - 9/15/03

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes eve irritation, Harmful if inhaled, Avoid contact with eves or clothing, Avoid breathing dust, Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

# FIRST AID

- IF IN EYES: Hold eve open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
  - · Call a poison control center or doctor for treatment advice.
- IF INHALED: Move person to fresh air.
  - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
  - · Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

In case of an emergency involving this product, call day or night (800) 424-9300.

# **ENVIRONMENTAL HAZARDS**

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

# DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

### NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION, OR FOR RESEARCH PURPOSES.

# STORAGE AND DISPOSAL

STORAGE: Store this product only in its original container in a dry, cool, secured storage area. Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

PRODUCT DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal. State or local procedures.

EMPTY CONTAINER DISPOSAL: Do not reuse bag. Completely empty bag into application equipment. Dispose of emptied bag(s) in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

SPILL: In case of spill, sweep up material and dispose of material according to disposal directions listed above.

# GENERAL INFORMATION

This product is a selective herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

#### Turfgrass Safety

# USE PRECAUTIONS

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, 0000 disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant at this product:

#### Cool-Season Grasses

Bentgrass, creeping\* (Agrostis palustris) Bluegrass, Kentucky (Poa pratensis) Fescue, fine\*\* (Festuca rubra) Fescue, tall (Festuca arundinacea) Ryegrass, perennial (Lolium perenne)

# Warm-Season Grasses Bahiagrass (Paspalum notatum) Bermudagrass (Cynodon dactylon)

Buffalograss\*\*\* (Buchloe dactyloides) Carpetgrass (Axonopus affinis)

Centipedegrass (Eremochioa ophilireides) Kikuyugrass (Penhišetüm clandestinum) St. Augustinegrass (Stenotaphrum Secundatum) Zoysiagrass (Zoysia japonica) 5

6 - 5 B **B B** 

Page 2 of 5 Pages LEBANON SEABOARD CORPORATION 961-360 LABEL NOTIFICATION - 9/15/03

\* Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested.

DO NOT apply this product to Colonial bentgrass (Agrostis tenuis) varieties.

\*\* Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested.

\*\*\* DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

NOTE: It is not possible to evaluate all turfgrass varieties for tolerance to Dimension. It is suggested that the user evaluate the response of turfgrass varieties to this product on a limited area, at recommended use rates, prior to initiating large-scale use.

# RESEEDING, OVERSEEDING, OR SPRIGGING

Reseeding, overseeding, or sprigging of treated areas with this product should be delayed until 12 weeks from the time of application. Reseeding, overseeding or sprigging before 12 weeks after application may prevent establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. If overseeding with perennial ryegrass in the fall, overseed 8 weeks or later after a single application. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

## **OTHER USE PRECAUTIONS**

The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. **DO NOT** apply this product until the grass has recovered from these cultural practices.

• DO NOT use clippings from treated turf for mulching around vegetables or fruit trees.

. Keep people and pets off treated areas until dust has settled.

## **APPLICATION DIRECTIONS**

#### Control of Crabgrass

#### Preemergence and Early Postemergence Control

This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs.

It can also provide "early postemergence" control of crabgrass. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first see the crabgrass plants in the lawn or turf. The practical benefit of this product's early postemergence activity is that it can give the user a 2-4 week longer period of time to make applications to control crabgrass.

#### **Application Frequency and Timing**

This product may be applied as a single application or as a split application for crabgrass control in the spring, summer, or fall.

#### **Spring Applications**

For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

- Use Program 1 (given in Table 1, below) for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.
- Use Program 2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control of crabgrass.
- Where split fertilizer applications are recommended, a rate in Table 1 may be applied in two applications made 6-10 weeks apart and prior to crabgrass emergence.

Page 3 of 5 Pages LEBANON SEABOARD CORPORATION 961-360 LABEL NOTIFICATION - 9/15/03

#### Table 1: Recommended Application Use Rates For Crabgrass\* (Lbs. of this Product per 1,000 Sq. Ft.)

516

1	2
PREEMERGENCE	PREEMERGENCE
(high)	(low)
	POSTEMERGENCE
2.2	4.4
3.2	6.4
4.4	8.8
6.4	8.8**
	(high) <u>2.2</u> <u>3.2</u> <u>4.4</u>

#### Fall Applications

This product can also be applied in the late summer or early fall (late August through November) at the use rates listed in Table 2 to provide control of crabgrass through the summer.

The fall application should be followed by an appropriate spring application to provide season-long control.

#### Table 2: Recommended Application Use Rates For Fall Applications (Lbs. of Product/1000 Sq. Ft.)

REGION	Preemergence
North	6.4
Transition	8.8*
South	8.8*
Coastal South	8.8*

\*DO NOT apply more than 8.8 lbs. of this product per 1000 sq. ft. per application, and no more than 26.4 lbs. of this product per 1000 sq. ft. per year.

In New York State, this product may only be used by Professional Applicators at no more than 8.8 lbs. per 1000 sq. ft. (0.5 lb. active ingredient per acre) per year; and is prohibited for use in Nassau and Suffolk Counties.

Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below.

 North:
 All other areas.

 Transition:
 DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, and coastal areas of CT, NY, and RI.

 South:
 AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, and TX.

 Coastal South:
 HI, FL, and southern coastal areas of AL, GA, LA, MS, NC, SC, and TX.

\*\* May require split or sequential applications for full control.

#### **Tips for Improved Control**

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result.

Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

#### Control of Other Grass and Broadleaf Weeds

#### Spring Applications

Used as directed for crabgrass control in the spring (Table 1), this product will also control or suppress the following weeds when applied prior to their emergence: Purslane, (common) Barnvardorass Goosegrass Smutgrass Bluegrass (annual, roughstalk) Kikuvugrass Lespedeza (common) Speedwell (corn) Crowfootgrass Ryegrass (annual & perennial) Oxalis (buttercup, creeping & yellow woodsorrel) Spurge (prostrate & spotted) Foxtail (yellow & green)\* - is - i, is \*Also controlled at the Program 2 rates. 

> Page 4 of 5 Pages LEBANON SEABOARD CORPORATION 961-360 LABEL NOTIFICATION - 9/15/03

Fall Applications	4	j,
Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring (Table 2), this product will also control	l or	
suppress the following weeds when applied prior to their emergence:		

Bluegrass (annual, roughstalk) Bittercress Chickweed Henbit Geranium (Carolina) Parsley-piert Pineappleweed Shepherdspurse

00000 00000 00000

0 . . .

2 1 . e . e . e

 $\gamma_{\rm e} V$ 

## **APPLICATION EQUIPMENT INSTRUCTIONS**

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or omamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

# SUGGESTED SPREADER SETTINGS

[Note: Registrant/Subregistrant may insert appropriate spreader setting information based on desired spreaders and recommended use rates.]

## CONDITIONS OF SALE AND WARRANTY

The Manufacturer warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. THE MANUFACTURER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Handling, storage and use of the product by Buyer or User are beyond the control of the Manufacturer and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. IN NO CASE WILL THE MANUFACTURER OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.

## Additional Labeling Requirements:

If the fertilizer guarantee contains iron, the following statement must appear:

This product contains the secondary nutrient iron. Iron may stain concrete and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash off with water.

If the recommended use rate applies less than 1 pound nitrogen per 1,000 sq. ft. per application, the following statement must appear:

Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

For products distributed in the State of Washington, the following statement must appear:

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-info-lebsea.com

Page 5 of 5 Pages LEBANON SEABOARD CORPORATION 961-360 LABEL NOTIFICATION - 9/15/03