

961-361

09/30/2005

34 A 6 - 2

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0080, Approval expires 2-28-95



United States  
Environmental Protection Agency  
Washington, DC 20460

 Registration  
 Amendment  
 Other

OPP Identifier Number

### Application for Pesticide - Section I

1. Company/Product Number 961-361	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Lebanon Fertilizer with Dimension Crabgrass Control	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Lebanon Seaboard Corporation 1600 East Cumberland Street Lebanon, PA 17042 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

### Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	<b>NOTIFICATION</b>  SEP 30 2005
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Alternate Brand Name: Lebanon ProScape Fertilizer with Dimension 0.19%  
This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statements to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under section 12 and 14 of FIFRA.

### Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Metal Plastic Glass Paper Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other _____					

### Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Henry E. Rohland	Title EPA Registration Manager	Telephone No. (include Area Code) 717-273-1635
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title EPA Registration Manager	
4. Typed Name Henry E. Rohland	5. Date 16 September 2005	

SEP 30 2005

2/7

LEBANON  
**ProScape** FERTILIZER WITH **DIMENSION® .19%**  
PRE & POSTEMERGENCE ANNUAL  
GRASSY WEED CONTROL

# LEBANON **ProScape**

SPECIMEN LABEL

FERTILIZER WITH  
**DIMENSION® .19%**  
PRE & POSTEMERGENCE ANNUAL GRASSY WEED CONTROL  
**16-4-8**  
**MESA**

GUARANTEED ANALYSIS

16% Nitrogen	16.00
4% Phosphate	4.00
8% Potassium	8.00
1.9% Dimension® .19%	1.90
Water Soluble	16.00
Other	0.00
Total	36.00

53600

SPREADER SETTINGS	Rate in pounds per 1,000 sq. ft.	Other
Product	16.00	1.90
Carrier	11.00	1.10
Water	1.00	0.10
Other	0.00	0.00
Total	28.00	3.10

Dimension® is a selective herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes.

ACTIVE INGREDIENTS: 0.19%

OTHER INGREDIENTS: 99.81%

TOTAL: 100.00%

This product contains 0.020 lb. of the active ingredient (Dimension®) per 50 lb. bag.  
Product protected by U.S. Patent Nos. 4,432,146 (7/19/81) and 4,432,147 (7/19/81).  
EPA Reg. No. 281-1021      EPA Reg. No. 281-1021-0104-01  
Dimension® is a registered trademark of Dow AgroSciences LLC      SuperSorb used as a first letter of lot number

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT.  
USE ONLY ACCORDING TO LABEL INSTRUCTIONS.  
Before using this product, read the literature fact sheet attached to this bag.  
If terms are not acceptable, return the unopened container to the place of purchase at once.

In New York State, this product may only be used by Professional Applicators at no more than 6.1 lbs. per 1,000 sq. ft. (0.5 lb. active ingredient per acre) per year, and is prohibited for use in Nassau and Suffolk Counties.

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**  
See back panel for additional Precautionary Statements.

Lebanon Seaboard Corporation  
1600 E. Cumberland St. • Lebanon, PA 17042  
www.LebanonTurf.com

NET WEIGHT **50** LBS. (22.68 KG)

LEBANON  
**ProScape** FERTILIZER WITH **DIMENSION® .19%**  
PRE & POSTEMERGENCE ANNUAL  
GRASSY WEED CONTROL

3/7

**FERTILIZER  
WITH  
DIMENSION® .19%  
PRE & POSTEMERGENCE ANNUAL GRASSY WEED CONTROL**

**16-4-8**

**MESA**  
P A T E N T E D

**GUARANTEED ANALYSIS**

Total Nitrogen (N).....	16%
4.0% Ammoniacal Nitrogen	
1.2% Water Insoluble Nitrogen	
8.3% Urea Nitrogen	
2.5% Other Water Soluble Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ).....	4%
Soluble Potash (K <sub>2</sub> O).....	8%
Total Sulfur (S).....	2.8%
2.8% Combined Sulfur (S)	
Total Iron (Fe).....	2.0%
0.2% Water Soluble Iron (Fe)	
Nutrient Sources: Ammonium Phosphate, Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash, Ferric Oxide, Ferrous Sulfate.	
Chlorine (Cl) not more than.....	7.0%
*3.7% Slowly Available Nitrogen from Methylene Ureas.	F699



**53600**

NOTICE This product contains the secondary nutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash off with water.  
MESA<sup>1</sup>, PennMulch<sup>2</sup> and ProScape<sup>3</sup> are registered trademarks of Lebanon Seaboard Corporation.

SPREADER SETTINGS	Rate in pounds per 1,000 Sq. Ft.		These settings were calibrated and field tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.
	3	4.5	
ProScape <sup>3</sup> SS/PrizeLAWN <sup>®</sup> BF-1/SS/CBR III	G	L	
PennMulch <sup>2</sup> HVO/PrizeLAWN <sup>®</sup> BF-HVO	G	L	
Earthway Rotary	11	15	
Lesco (letter dial/numeric dial)	D / 10	J / 21	
Scotts AccuPro 2000/SR-2000	G	K	
Scotts R8A Rotary	J	M	
Scotts R7X Rotary	E	I	
Spyker Rotary	3	4 1/2	

**Dimension<sup>3</sup> is a selective herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes.**

PENN MULCH

4/7

**ACTIVE INGREDIENTS:**

\*Dithiopyr ..... 0.19%

**OTHER INGREDIENTS** ..... **99.81%**

**TOTAL** ..... **100.00%**

\*This product contains 0.095 lb. of the active ingredient, dithiopyr, per 50 lb. bag.

Cas No. 97886-45-8

Product protected by U.S. Patent No. 4,692,184. Other patents pending.

(L) (D)

EPA Reg. No. 961-361

EPA Est. 961-PA-01:961-IL-01

\*Dimension is a registered trademark of Dow AgroSciences LLC.

Superscript used is first letter of lot number.

**READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT.**

**USE ONLY ACCORDING TO LABEL INSTRUCTIONS.**

Before buying this product, read the Warranty Section elsewhere on this label.

If terms are not acceptable, return the unopened container to the place of purchase at once.

In New York State, this product may only be used by Professional Applicators at no more than 6.1 lbs. per 1000 sq. ft. (0.5 lb. active ingredient per acre) per year; and is prohibited for use in Nassau and Suffolk Counties.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See back panel for additional Precautionary Statements.

**Lebanon Seaboard Corporation**

1600 E. Cumberland St. • Lebanon, PA 17042

[www.LebanonTurf.com](http://www.LebanonTurf.com)

SPECIMEN LABEL

3 HT **50** LBS. (22.68 KG)

# LEBANON ProScape®

shy  
NOTIFICATION  
SEP 30 2005

## FERTILIZER WITH DIMENSION® .19% PRE & POSTEMERGENCE ANNUAL GRASSY WEED CONTROL

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

#### FIRST AID

- IF IN EYES:**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
  - Call a poison control center or doctor for treatment advice.
- IF INHALED:**
- Move person to fresh air.
  - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
  - Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Emergency Medical Information call (888) 208-1368. For any other emergency involving this product, call day or night (800) 424-9300.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

**NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION, OR FOR RESEARCH PURPOSES.**

### STORAGE AND DISPOSAL

**STORAGE:** Store this product only in its original container in a dry, cool, secured storage area. Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**PRODUCT DISPOSAL:** Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures.

**EMPTY CONTAINER DISPOSAL:** Do not reuse bag. Completely empty bag into application equipment. Dispose of emptied bag(s) in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**SPILL:** In case of spill, sweep up material and dispose of material according to disposal directions listed above.

### GENERAL INFORMATION

This product is a selective herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

### USE PRECAUTIONS

#### Turfgrass Safety

This product may be used on seeded, sodded, or sprung lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or springing before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

#### Cool-Season Grasses

Bentgrass, creeping\* (*Agrostis palustris*)  
Bluegrass, Kentucky (*Poa pratensis*)  
Fescue, fine\*\* (*Festuca rubra*)  
Fescue, tall (*Festuca arundinacea*)  
Ryegrass, perennial (*Lolium perenne*)

#### Warm-Season Grasses

Bahagrass (*Paspalum notatum*)  
Bermudagrass (*Cynodon dactylon*)  
Buffalograss\*\*\* (*Buchloe dactyloides*)  
Carpetgrass (*Axonopus affinis*)  
Centpedegrass (*Eremochloa ophiuroides*)  
Kikuyugrass (*Pennisetum clandestinum*)  
St. Augustinegrass (*Stenotaphrum secundatum*)  
Zoysiagrass (*Zoysia japonica*)

\* Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested.

DO NOT apply this product to Colonial bentgrass (*Agrostis tenuis*) varieties.

\*\* Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bigart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivado', 'Lanestown', 'Koket', 'Maienta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested.

\*\*\* DO NOT use this product on seeding buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

NOTE: It is not possible to evaluate all turfgrass varieties for tolerance to Dimension. It is suggested that the user evaluate the response of turfgrass varieties to this product on a limited area, at recommended use rates, prior to initiating large-scale use.

6/7

**RESEEDING, OVERSEEDING, OR SPRIGGING**

Reseeding, overseeding, or sprigging of treated areas with this product should be delayed until 12 weeks from the time of application. Reseeding, overseeding or sprigging before 12 weeks after application may prevent establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. If overseeding with perennial ryegrass in the fall, overseed 8 weeks or later after a single application. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

**OTHER USE PRECAUTIONS**

The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller. For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. **DO NOT** apply this product until the grass has recovered from these cultural practices.

- **DO NOT** use clippings from treated turf for mulching around vegetables or fruit trees.
- Keep people and pets off treated areas until dust has settled.

**APPLICATION DIRECTIONS**  
**Control of Crabgrass**

**Preemergence and Early Postemergence Control**

This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs. It can also provide "early postemergence" control of crabgrass. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first see the crabgrass plants in the lawn or turf. The practical benefit of this product's early postemergence activity is that it can give the user a 2-4 week longer period of time to make applications to control crabgrass.

**Application Frequency and Timing**

This product may be applied as a single application or as a split application for crabgrass control in the spring, summer, or fall.

**Spring Applications**

For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

- Use Program 1 (given in Table 1, below) for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.
- Use Program 2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control of crabgrass.
- Where split fertilizer applications are recommended, a rate in Table 1 may be applied in two applications made 6-10 weeks apart and prior to crabgrass emergence.

PROGRAM	1		2	
	USE (turfgrass cut)	PREEMERGENCE (high)	PREEMERGENCE (low)	POSTEMERGENCE
<b>REGION</b>				
North		N.R.***		3.0
Transition		2.2		4.5
South		3.0		6.1
Coastal South		4.5		6.1**

REGION	Preemergence
North	4.5
Transition	6.1*
South	6.1*
Coastal South	6.1*

**Fall Applications**

This product can also be applied in the late summer or early fall (late August through November) at the use rates listed in Table 2 to provide control of crabgrass through the summer.

The fall application should be followed by an appropriate spring application to provide season-long control.

\*DO NOT apply more than 6.1 lbs. of this product per 1000 sq. ft. per application, and no more than 18.2 lbs. of this product per 1000 sq. ft. per year.

In New York State, this product may only be used by Professional Applicators at no more than 6.1 lbs. per 1000 sq. ft. (0.5 lb. active ingredient per acre) per year; and is prohibited for use in Nassau and Suffolk Counties.

Postemergence control is limited: see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below.

- North:** All other areas.
- Transition:** DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, and coastal areas of CT, NY, and RI.
- South:** AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, and TX.
- Coastal South:** HI, FL, and southern coastal areas of AL, GA, LA, MS, NC, SC, and TX.

\*\* May require split or sequential applications for full control. \*\*\* N.R. - This formulation is not recommended.

**Tips for Improved Control**

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result.

Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

**Control of Other Grass and Broadleaf Weeds**

**Spring Applications**

Used as directed for crabgrass control in the spring (Table 1), this product will also control or suppress the following weeds when applied prior to their emergence:

Barnyardgrass	Goosegrass	Oxalis (buttercup, creeping & yellow woodsorrel)	Smutgrass
Bluegrass (annual, roughstalk)	Kikuyugrass	Purslane (common)	Speedwell (corn)
Crowfootgrass	Lespedeza (common)	Ryegrass (annual & perennial)	Spurge (prostrate & spotted)
Foxtail (yellow & green)*			

\* Also controlled at the Program 2 rates.

**Fall Applications**

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring (Table 2), this product will also control or suppress the following weeds when applied prior to their emergence:

Bluegrass (annual, roughstalk)	Bittercress	Chickweed	Geranium (Carolina)	Henbit	Parsley-piert	Pineappleweed	Shepherdspurse
--------------------------------	-------------	-----------	---------------------	--------	---------------	---------------	----------------

**APPLICATION EQUIPMENT INSTRUCTIONS**

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

NOTE: Supplemental applications of lawn fertilizer may be needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

SPECIMEN LABEL

**CONDITIONS OF SALE AND WARRANTY**

The Manufacturer warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. THE MANUFACTURER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Handling, storage and use of the product by Buyer or User are beyond the control of the Manufacturer and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. IN NO CASE WILL THE MANUFACTURER OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.

**MANUFACTURED BY:**  
**LEBANON SEABOARD CORPORATION**  
1600 EAST CUMBERLAND STREET  
LEBANON, PA 17042  
[www.LebanonTurf.com](http://www.LebanonTurf.com)

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-lebesa.com>

7/7  
SPECIMEN LABEL