

# UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

APR 2 4 2009

Mrs. Sylvia Whitman State Registration Coordinator Lebanon Seaboard Corporation 1600 East Cumberland Street Lebanon, PA 17042

RE: Notification of Alternate Brand Name: Lebanon Pro Brand Fertilizer with Dimension

.19% Pre & Postemergence Annual Grassy Weed Control

Registration Number: 961-361

Submission Dated: February 23, 2009

Dear Mrs. Whitman:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated, February 23, 2009, for the product, Lebanon Fertilizer with Dimension Crabgrass Control. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely yours,

\_\_\_\_\_

Linda Arrington Notifications & Minor Formulations Team Leader Registration Division (7505P) Office of Pesticide Programs

Please read instructions on reverse before completing for	_	Form Amorova	d 0049 No 20	70 008/	Ampeouslanding 9 99 95
United St Environmental Prot Washington, D	tates ection Agency	Form Approve.  ✓	Registrati Amendm Other	ion	O. Approval expires 2-28-95 OPP Identifier Number
Appli	cation for Pesti	cide - Section	<u> </u>		
1. Company/Product Number 961-361	2. EP	A Product Manager nne Miller		3. Pro	posed Classification
4. Company/Product (Name) Lebanon Fertilizer with Dimension Crabgrass Co	ontrol PM#	23			
5. Name and Address of Applicant (Include ZIP Code)  Lebanon Seaboard Corporation 1600 East Cumberland Street  Lebanon, PA 17042  Check if this is a new address	(b)(i), to: EPA		CATIONIC		FIFRA Section 3(c)(3) mposition and labeling
	Section		1750	170)	149N
Amendment - Explain below.  Resubmission in response to Agency letter dated  Notification - Explain below.  Explanation: Use additional page(s) if necessary. (For Alternate Brand Name: Lebanon Pro Brand Fertilizer with Di This notification is consistent with the provisions of PR Notice labeling or the confidential statement of formula of this produto EPA. I further understand that if this notification is not confidential may be subject to enforcement action and penal	imension .19% Pre & Po e 98-10 and EPA regulat uct. I understand that it is nsistent with the terms of	stemergence Annual tions at 40 CFR 152.4 s a violation of 18 U.S PR Notice 98-10 and	elow.  Grassy Weed Co 6 and no other c C. Sec. 1001 to	ontrol hanges h	make any false statements
	Section -	· 1f1			
1. Material This Product Will Be Packaged In:  Child-Resistant Packaging  Yes  No  No  * Certification must be submitted  Unit Packaging  Yes  Ves  Ves  Ves  Ves  Ves  Ves  Ves		No. per container		ontainer Metal Plastic Glass Paper Other (S	pecify)
3. Location of Net Contents Information 4. Size  Label Container	(s) Retail Container	5. Lo	ocation of Label	Directio	ins
6. Manner in Which Label is Affixed to Product	Lithograph Paper glued Stenciled	Other			
	Section -	IV			
1. Contact Point (Complete items directly below for identity	ification of individual to	be contacted, if ned	essary, to proc	ess this	application.)
Name Sylvia Whitman	Title State Reg	istration Manager	1	elephone 717-273	B-1685
Cer I certify that the statements I have made on this for I acknowledge that any knowlingly false or mislead both under applicable law.					6. Date Application Received (Stamped)
2. Signature	3. Title State Regis	stration Manager	,	,	
4. Typed Name Sylvia Whitman	5. Date February	23, 2009			• • •



### LEBANON SEABOARD CORPORATION

1600 EAST CUMBERLAND STREET • LEBANON, PENNSYLVANIA 17042

(717) 273-1685 FAX (717) 273-9466

March 5, 2009

Document Processing Desk
Office of Pesticide Program (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 33302-4501

ATTN: Joanne Miller (Team 23)

RE: Alternate Brand Names

- EPA Reg. No. 961-359 Lebanon Pro Brand Fertilizer with Dimension 0.09% Pre & Postemergence Annual Grassy Weed Control
- EPA Reg. No. 961-360 Lebanon Pro Brand Fertilizer w/Dimension .13% Pre & Postemergence Annual Grassy Weed Control
- EPA Reg. No. 961-361 Lebanon Pro Brand Fertilizer with Dimension .19% Pre & Postemergence Annual Grassy Weed Control
- EPA Reg. No. 961-362 Lebanon Pro Brand Fertilizer with Barricade 0.43% Preemergence Weed Control
- **EPA Reg. No. 961-365** Lebanon Pro Brand Fertilizer with Barricade 0.38% Preemergence Weed Control
- EPA Reg. No 961-367 Lebanon Pro Brand Fertilizer with Barricade 0.29%

Dear Ms. Miller:

Please find enclosed Lebanon's applications for Alternate Brand Name.

- Transmittal Document (this letter)
- EPA Form 8570-1 (6)
- Copy of each referenced label (6)

Should you have any questions, please feel free to contact me at 800/532-0090 ext. 361.

Sincerely

State Registration Coordinator

enclosures:



# **FERTILIZER WITH DIMENSION® 0.19%**

PRE & POSTEMERGENCE ANNUAL GRASSY WEED CONTROL



### **GUARANTEED ANALYSIS**

Chlorine (Cl) not more than .
\* 7.5% Slowly Available Nitrogen from Sulfur Coated Urea F699 Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-info-lebsea.com. NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum of 1 pound nitrogen per 1,000 q. ft.

Meth-Ex\*, PennMulch\* and ProScape\* are registered trademarks of Lebanon Seaboard Corporation.

0 88685 42739 8
T000

22-42739

SPREADER SETTINGS	Rate in pounds	per 1,000 Sq. Ft.	These settings were
0.112.12211.0211.1100	3	4.5	calibrated and field
LebanonTurf	4 1/2	4 3/4	tested, However, age
ProScape® SS/PrizeLAWN® BF-1/SS/CBR III	K	M	and condition of
PennMulch® HVO/PrizeLAWN® BF-HVO	K	M	spreader, speed of
Earthway Rotary	15	17	operation, and
Lesco (letter dial/numeric dial)	1/17	J/21	evenness of terrain
Scotts AccuPro 2000/SR-2000	Κ.	M	may require slightly
Scotts R8A Rotary	L	M	different settings for
Scotts R7X Rotary	G		desired coverage.
Spyker Rotary	4 1/2	4 3/4	

Dimension® is a selective herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes.

NOTIFICATION

APR 24 2009

ACTIVE INGREDIENTS:	
*Dithiopyr	0.19%
OTHER INGREDIENTS	99.81%

\*This product contains 0.095 lb. of the active ingredient, dithiopyr, per 50 lb. bag. Product protected by U.S. Patent No. 4,692,184. Other patents pending. \*Dimension is a registered trademark of Dow AgroSciences LLC.

EPA Reg. No. 961-361 Cas No. 97886-45-8

(D) EPA Est. 961-PA-01:961-IL-01 Superscript used is first letter of lot number.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS. Before buying this product, read the Warranty Section elsewhere on this label. If terms are not acceptable, return the unopened container to the place of purchase at once.

In New York State, this product may only be used by Professional Applicators at no more than 6.1 lbs, per 1000 sq.ft. (0.5 lb. active ingredient per acre) per year; and is prohibited for use in Nassau and Suffolk Counties.

# KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel for Additional Precautionary Statements.

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

# FIRST AID

IF IN EYES: IF INHALED:

Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 Call a poison control center or doctor for treatment advice.
 Move person to fresh air.
 If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
 Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Emergency Medical Information call (888) 208-1368. For any other emergency involving this product, call day or night (800) 424-9300.

# **ENVIRONMENTAL HAZARDS**

This product is toxic to fish and highly toxic to other aquatic organisms including cysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.



For technical assistance or more information about our products visit www.l.ebanonTurf.com

NET WEIGHT 50 LBS. (22.7 Kg)

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling

NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION, OR FOR RESEARCH PURPOSES.

GENERAL INFORMATION

The standing of course fairways, roughs, and tee boxes. This product will not harm nearby established This product is a selective herbicide that provides control of crabgrass and other annual grasses and broadleaf w omamentals when used according to label directions. Turfgrass Safety **USE PRECAUTIONS** 

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and omamental lurfs that are composed of the following turtgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

Cool-Season Grasses: Bentgrass, creeping\* (Agrostis palustris), Bluegrass, Kentucky (Poa pratensis), Fescue, fine\*\* (Festuca rubra), Fescue, tall (Festuca arundinacea), Ryegrass, perennial (Lolium perenne)

Warm-Season Grasses: Bahiagrass (Paspalum notatum), Centipedegrass (Etemochioa ophiuroides), Bermudagrass (Cynodon dactylon), Kikuyugrass (Pennisetum chandestinum), Bulfalograss: "(Buchole dactyloides), St. Augustinegrass (Stenotaphrum secundatum), Carpetgrass (Axonopus affinis), Zoysiagrass (Zoysia japonica)

\* Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. DO NOT apply this product to Colonial bentgrass (Agrostis tenuis) varieties.

"Use of this product on certain varieties of fine lescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koker', 'Majenta', 'May', 'Pennlawn', Tamara', Tatjana', 'Waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested.
"" DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

NOTE: It is not possible to evaluate all turfgrass varieties for tolerance to Dimension. It is suggested that the user evaluate the response of turfgrass varieties to this product on a limited area, at recommended use rates, prior to initiating large-scale use.

# RESEEDING, OVERSEEDING, OR SPRIGGING

Researding, overseeding, or springing of treated areas with this product should be delayed until 12 weeks from the time of application. Researding, overseeding or springing before 12 weeks after application may prevent establishment of desirable turty assess. When researding or overseeding, proper outtural practices such as soil cultivation, irrigation and fertilization should be followed. If overseeding with perennial lyegrass in the fat, overseed 8 weeks or later after a single application. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

### OTHER USE PRECAUTIONS

The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticulting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

- DO NOT use dippings from treated turf for mulching around vegetables or fruit trees.
   Keep people and pets off treated areas until dust has settled.

#### **APPLICATION DIRECTIONS**

### **Control of Crabgrass**

Preemergence and Early Postemergence Control

This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and omamental turfs. It can also provide "early postemergence" control of crabgrass. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first see the crabgrass plants in the lawn or turf. The practical benefit of this product's early postemergence activity is that it can give the user a 2-4 week longer period of time to make applications to control crabgrass.

#### **Application Frequency and Timing**

This product may be applied as a single application or as a split application for crabgrass control in the spring, summer, or fall.

Spring Applications

For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the control programs listed in Table 1 below, depending on the user's location, the turigrass mowing height, and whether the use is preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turigrass compet-

- Use Program 1 (given in Table 1, below) for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control
- Use Program 2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control of crabgrass.
- . Where split fertilizer applications are recommended, a rate in Table 1 may be applied in two applications made 6-10 weeks apart and prior to crabgrass emergence.

	: Recommended Application for Crabgrass* s. of this Product per 1,00	
PROGRAM	1	2
USE (turfgrass cut)	PREEMERGENCE (high)	PREEMERGENCE (low)
REGION		POSTEMERGENCE
North	3.2	6.4
Transition	4.7	9.3
South	6.4	N.H."
Coastal South	N.R.*	N.H.

Table 2: Recommended Application Use Rates For Fall Applications (Lbs. of Product per 1,000 Sq. Ft.)	
REGION	Preemergence
North	9.3
Transition	N.H.**
South	N.R.**
Coastal South	N.H.

Fall Applications
This product can also be applied in the late summer or early fall (late August through November) at the use rates listed in Table 2 to provide control of crabgrass through the summer.

The fall application should be followed by an appropriate spring application to provide season-long control.

"DO NOT apply more than 9.3 lbs. of this product per 1000 sq. t. per application, and no more than 19.3 lbs. of this product may only be used by Professional Applications and no more than 19.4 lbs. per 1000 sq. tt. per year; and is prohibited for use in Nassau and Suffolk Counties. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas fisted below.

North: All other areas.

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, and coastal areas of CT, NY, and RI.

AL, AR, AZ, CA, CA, LA, MS, NC, NM, NY, OK, SC, TN, and TX.

Coastal South: HI, FL, and southern coastal areas of AL, GA, LA, MS, NC, SC, and TX.

\*\* N.R. - This formulation is not recommended.

Tips for Improved Control

For best results, apply this product within a few days after moving and delay moving again for a few days after application. When treated lawn or ornamental turl areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result. Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

#### Control of Other Grass and Broadleaf Weeds

Spring Applications
Used as directed for crabgrass control in the spring (Table 1), this product will also control or suppress the following weeds when applied prior to their emergence:

Barnyardgrass Bluegrass (annual, roughstalk) Crowlootgrass

Goosegrass Kikuyugrass Lespedeza (common) Oxalis (buttercup, creeping & yellow woodsorrel) Purslane (common) Ryegrass (annual & perennial)

Smutgrass Speedwell (corn) Spurge (prostrate & spotted)

Foxtail (yellow & green)\*
\* Also controlled at the Program 2 rates.

Fall Applications
Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring (Table 2), this product will also control or suppress the following weeds when applied prior to their emergence:
Bluegrass (annual, roughstalk)
Bittercress
Chickweed
Geranium (Carolina)
Henbit
Parsley-piert
Pineappleweed
Shepherdspurse

#### APPLICATION EQUIPMENT INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turt area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

NOTE: Supplemental applications of lawn fertilizer may be needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

#### STORAGE AND DISPOSAL

Do not contaminate water, tood, or feed by storage or disposal.

PESTICIDE STORAGE: Store product in its original container in a cool, dry, locked place out of the reach of children and animals.

PESTICIDE DISPOSAL: Wasles resulting from the use of this product must be disposed of on site or at an approved waste

disposal facility.
CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, I allowed by State and local authorities, by burning. If burned stay out of smoke

CONDITIONS OF SALE AND WARRANTY The Manufacturer warrants that the product conforms to its chemical description and is reasonably fil for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. THE MANUFACTURER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Handling, storage and use of the product by Buyer or User are beyond the control of the Manufacturer and Seller. Risks such as crop injury, inefectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. IN NO CASE WILL THE MANUFACTURER OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.

# MANUFACTURED BY:

# NOTIFICATION

APR 24 2009

LEBANON SEABOARD CORPORATION

1600 EAST CUMBERLAND STREET LEBANON, PA 17042 www.LebanonTurf.com

ProScape\* is a registered trademark of Lebanon Seaboard Corporation Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-info-lebsea.com

22-42739