

PM 23

961-363

10-24-97

10419

OCT 24 1997

Mr. Henry E. Rohland
Lebanon Chemical Corporation
P.O. Box 180, 1600 E. Cumberland Street
Lebanon, Pennsylvania 17042

Dear Mr. Rohland:

Subject: Lebanon Fertilizer with Barricade® Preemergence Weed Control (0.43%)
EPA Registration No. 961-362
Lebanon Fertilizer with Barricade® Preemergence Weed Control (0.65%)
EPA Registration No. 961-363
Lebanon Fertilizer with Barricade® Preemergence Weed Control (0.22%)
EPA Registration No. 961-364
Lebanon Fertilizer with Barricade® Preemergence Weed Control (0.30%)
EPA Registration No. 961-365
Lebanon Fertilizer with Barricade® Preemergence Weed Control (0.29%)
EPA Registration No. 961-367
Applications Dated August 25, 1997, Request To Update Labels To Match the Barricade® F Label for EPA registration No. 55947-144 and Your Letters Dated August 25, 1997

Your request to amend the registrations of the subject pesticide product registrations to reflect the labeling of the me-too product (Barricade F) has been reviewed and found acceptable under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, provided that you:

1. Correct the statement on the front panels that reads: "Barricade® is a registered trademark of Sandoz, Inc." to read: "Barricade® is a registered trademark of Novartis Crop Protection, Inc."

CONCURRENCES

CONCURRENCES								
SYMBOL								
SURNAME								
DATE								

20/19

-2-

2. Alphabetize the list of weeds controlled in the "Weeds Controlled" sections of the labels.
3. On the labels, correct the list of "Tolerant Ornamental Species" by adding "Geranium cinereum Cranesbill". On copies of the labels submitted this entry is blurred and illegible.
4. Submit one (1) printed copy of the final printed label before releasing the product for shipment under the enclosed stamped label.

If these conditions are not complied with, each registration will be subject to cancellation in accordance with FIFRA, Section 6(e). Your release for shipment of the products constitutes acceptance of these conditions.

A stamped copy of each label is enclosed for your records.

Sincerely yours,

Joanne I. Miller
 Product Manager (23)
 Herbicide Branch
 Registration Division (7505C)

S. Wilson: Diskette Prodimine 10-24-97

CONCURRENCES

SYMBOL								
SURNAME								
DATE								

30/19

**LEBANON FERTILIZER WITH BARRICADE 0.65%
PREEMERGENCE WEED CONTROL**

ACTIVE INGREDIENT:

Prodiamine: [N³,N³-Di- η -propyl-2,4-dinitro-6-(trifluoromethyl)- η -phenylenediamine].....0.65%

INERT INGREDIENTS: 99.35%

TOTAL:..... 100.0%

Guaranteed Fertilizer Analysis:

Nitrogen 32 % (N)
Phosphorus 4 % (P₂O₅)
Potassium 6 % (K₂O)

Nitrogen from: Urea formaldehyde, Urea, Ammonia

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

OCT 23 1997

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**

961-363

For selective preemergence control of grass and broadleaf weeds in:

- established turf grasses (excluding golf course putting greens) and lawns
- landscape ornamentals (including established perennials & wildflower plantings)

KEEP OUT OF REACH OF CHILDREN

CAUTION

See the following section, Precautionary Statements, Hazards To Humans and Domestic Animals, for additional information.

Net Weight:

EPA Reg. No. 961-363

EPA Est. No.

Lebanon Chemical Corp.
P.O. Box 180
Lebanon, PA 17042

BARRICADE® is a Registered Trademark of Sandoz, Ltd.

4/19

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

Statements of Practical Treatment

If in eyes: Flush thoroughly with water for several minutes. Get medical attention if irritation persists.

If on skin: Wash with soap and water. Rinse thoroughly. Get medical attention if irritation persists.

If inhaled: Remove victim to fresh air.

Environmental Hazards

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

If material is spilled: Contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE

Store in original container away from other fertilizer, feed, or food stuffs and separated from other pesticides.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Paper containers: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or at an incineration facility, or; if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

6/21/19

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- established turf grasses (excluding golf course putting greens) and lawns
- landscape ornamentals
- established perennials and wildflower plantings

LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not graze or feed livestock forage cut from areas treated with LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL.

Do not apply this product through any type of irrigation system.

Do not apply aerially.

Do not apply to golf course putting greens.

Application Directions


Apply uniformly with suitable, calibrated application equipment.

ESTABLISHED TURF






LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL that may be applied per year is given for each turf grass species in the **Maximum Annual Rates** section of this label.

Most effective weed control in turf grasses will be obtained when LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

CRABGRASS SEED GERMINATION DATES



Approximate Date

-  After May 30
-  After May 10
-  After April 20
-  After March 20
-  January 1 to March 20

Use Precautions

The following precautions apply to the use of LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL in turf grasses and lawns:

Application of LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL may thin emerged annual bluegrass and newly overseeded grasses.

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year.

) Application of LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL to turf stressed by drought, low fertility, or pest damage may result in turf injury.

Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.

Do not apply LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

9/19

Rates of Application

LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL is related to the rate applied.

Maximum Annual Rates

LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Application Rate/Calendar Year of LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL by Turf Grass Species ¹⁾		
Turf Species:	lbs product/A	lbs ai/A
Bermudagrass ²⁾ Bahigrass Centipedegrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type) Zoysia	230	1.5
Buffalograss Kentucky Bluegrass Perennial Ryegrass	154	1.0
Creeping Red Fescue	115	0.75
Creeping Bentgrass	100	0.65

¹⁾ These are the maximum rates per calendar year by species limitations.

²⁾ May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 80 lbs/A. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

- Do not apply more than 230 lbs. of LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL per calendar year.
- Use higher rates of LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the **Maximum Annual Rates Table**.

Weeds Controlled

When used in accordance with this label LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL will provide control of the following weeds:

Barnyardgrass	Carpetweed
Bluegrass, Annual (<i>Poa annua</i>)	Chickweed, Common
Crabgrass (large, smooth)	Chickweed, Mouseear (from seed)
Crowfootgrass	Henbit
Cupgrass, Woolly	Knotweed
Foxtails, Annual	Kochia
Goosegrass ¹	Lambsquarter, Common
Itchgrass	Pigweed
Johnsongrass (from seed)	Purslane, Common
Junglerice	Puslev, Florida
Lovegrass	Shepherd's Purse ²
Panicum (Texas, Fall, Browntop)	Speedwell, Persian
Rescuegrass ³	Spurge, Prostrate
Signalgrass, Broadleaf	Woodsorrel, Yellow (from seed)
Sprangletop	
Witchgrass	

- 1 In many areas a single application of 100 to 230 lbs/A of LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 100 to 154 lbs/A followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rate Table. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rate Table above.
- 2 Applications for this weed should be made in late summer, fall or winter prior to germination.
- 3 Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL Rate (lb/A)		Months Before Overseeding		
Product	ai	North	Transition	South
77	.50	4	4	4
100	.66	5	4	4
115	.74	6	5	5
123	.80	---	6	6
154	1.00	---	7	7
175	1.14	---	---	9
200	1.30	---	---	10
230	1.50	---	---	12

LANDSCAPE ORNAMENTALS

(including Established Perennials and Wildflower Plantings)

LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL may be applied for residual preemergence weed control in ornamentals.

Use Rates

Apply LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL at 100 to 230 lbs/A in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 230 lbs/A.

Application Timing and Information

LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

Established Landscape Ornamentals

LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL may be applied at any time to established ornamentals as a broadcast, over the top or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL in the soil.

Newly Transplanted Landscape Ornamentals

LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL may be applied to newly transplanted ornamentals as a broadcast, over the top or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL. Apply LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

Tolerant Ornamental Species

LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL. Best results will be obtained when LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL may be applied over the top of these species.

When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to LEBANON FERTILIZER WITH BARRICADE 0.65% PREEMERGENCE WEED CONTROL.

Scientific name	Common name
<i>Abies</i> spp.	Fir species * * (Balsam, Fraser, Noble, etc.)
<i>Abelia grandiflora</i>	Abelia: Sherwood
<i>Acer palmatum</i>	Japanese Maple
<i>A. platanoides</i>	Norway Maple
<i>Achillea</i> sp.	Yarrow: King Edward
<i>Actinidia chinensis</i>	Kiwi *
<i>Agapanthus africanus</i>	Lily-of-the-Nile (African Lily)
<i>Agapanthus orientalis</i>	
<i>Akebia quintata</i>	Five-Leaf or Chocolate Vine
<i>Allium cernuum</i>	Lady's Leek, Nodding Onion
<i>Anemone hybrida</i>	Japanese Anemone
<i>Aquilegia</i> sp.	Aquilegia: Red and Gold
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita
<i>Arctotheca calendula</i>	Cape Weed
<i>Artemesia</i> sp.	Wormwood; Silver Mound Castle
<i>Aster</i> sp.	Aster: Bonny Blue, Purple Dome
<i>Aster X Frikartii</i>	
<i>Athrium Filiz-femina</i>	Lady Fern; Fern Lady

<i>Aucuba japonica</i>	Japanese Aucuba
<i>Begonia sp.</i>	Fibrous Begonia: Hardy Grandis
<i>Berberis gladywynensis</i>	Barberry
<i>B. Julianae</i>	Wintergreen Barberry
<i>B. mentorensis</i>	Mentor Barberry
<i>B. Thunbergii</i>	Japanese Barberry
<i>B. verruculosa</i>	Warty Barberry
<i>Bergenia cordifolia</i>	
<i>Boltonia asteroides</i>	Snowbank
<i>Buddleia davidii</i>	Butterfly-Bush (Dwarf Blue); Royal Red

* Do not use on food producing trees, vines, or plants.

** Not for use on container grown plants.

<u>Scientific Name</u>	<u>Common Name</u>
<i>Buxus microphylla</i>	Japanese Boxwood
<i>Callistemon viminalis</i>	Weeping Bottlebrush
<i>Calluna vulgaris</i>	Scotch Heather
<i>Campanula carpatica</i>	Tussock Bellflower, (White Clips)
<i>Campis X Tagliabuana</i>	Trumpet Creeper, Trumpet Flower, Madame Galen
<i>Carpobrotus edulis</i>	Hottentot Fig (Ice Plant)
<i>Cassia artemisooides</i>	Feathery Cassia
<i>Ceanothus rigidus</i>	Wild Lilac
<i>Ceratostigma plumbagonoides</i>	
<i>Chamaecyparis pisifera</i>	False Cypress
<i>Chrysanthemum nipponicum</i>	
<i>Cleyera japonica</i>	Cleyera
<i>Citrus spp.</i>	Citrus species*
<i>Coreopsis sp.</i>	Coreopsis (Calliopsis): Early Sunrise, Moonbeam
<i>Cornus florida</i>	Flowering Dogwood
<i>C. stolonifera</i>	American Dogwood
<i>Cortaderia selloana</i>	Pampas Grass
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster
<i>C. buxifolius</i>	Cotoneaster
<i>C. Dammeri</i>	Bearberry Cotoneaster
<i>C. microphyllus</i>	Rockspray Cotoneaster
<i>Cretaeagus spp.</i>	Hawthorne
<i>Crocasmia sp.</i>	Lucifer
<i>Cupressus sempervirens</i>	Italian Cypress
<i>Delosperma alba</i>	White Trailing Ice Plant
<i>Delosperma sp.</i>	Cooperi Pink
<i>Delphinium sp.</i>	Larkspur: Blue Elf

<i>Dianthus deltoides</i>	Dianthus; Maiden Pinks "Zing"
<i>D. gratianopolitanus</i>	Cheddar Pink
<i>Dodonea viscosa</i>	Hop Bush
<i>Echinacea pupurea</i>	Coneflower, Purple; Magnus
<i>Elaeagnus pungens</i>	Silverberry
<i>Euonymus Fortunei</i>	Wintercreeper
<i>E. japonica</i>	Japanese Spindle Tree (Evergreen Euonymus)
<i>E. kiautschovica</i>	Spreading Euonymus
<i>Fatsia japonica</i>	Japanese Aralia
<i>Forsythia intermedia</i>	Border Forsythia
<i>Forsythis suspensa</i>	Weeping Forsythia
<i>F. viridissima</i>	Greenstem Forsythia

* Do not use on food producing trees, vines, or plants.
 ** Not for use on container grown plants.

<u>Scientific Name</u>	<u>Common Name</u>
<i>Gaillardia sp.</i>	Gaillardia, Blanket Flower: "Goblin"
<i>Gardenia jasminoides</i>	Gardenia, Cape-Jasmine, Radicans
<i>Gaura sp.</i>	
<i>Gentiana dahurica</i>	Gentian
<i>Gladiolus spp.</i>	Gladiolus species**
<i>Gypsophila repens</i>	Baby's Breath
<i>Hedera helix</i>	English Ivy
<i>Hellanthemum sp.</i>	Sunrose
<i>Hemerocallis sp.</i>	Daylily: Aztec Gold, Stella De Oro, Tender Love
<i>Hibiscus</i>	Rose of Sharon**
<i>Hibiscus sp.</i>	Mallow: Disco Belle White
<i>Hibiscus Rosa-sinensis</i>	Chinese Hibiscus
<i>Hosta plantaginea</i>	Hosta, Plantain Lily (Fragrant)
<i>H. Sieboldiana</i>	Hosta, "Searsucker"
<i>Houttuynia cordata var. variegata</i>	
<i>Hydrangea macrophylla</i>	Bigleaf Hydrangea
<i>Ilex Beryi</i>	Holly
<i>Ilex cornuta</i>	Chinese Holly
<i>I. crenata</i>	Japanese Holly; Helli
<i>I. opaca</i>	American Holly
<i>I. vomitoria</i>	Yaupon Holly; Schillings
<i>Inula ensifolia</i>	
<i>Iris ensata</i>	Sword-Leaved Iris; Jodlesong
<i>Iris siberica</i>	Siberian Iris; Cabernet
<i>Iris spp.</i>	Iris species**
<i>Jasminium nudiflorum</i>	Winter Jasmine
<i>Juniperus chinensis</i>	Chinese Juniper; Nick's Compact, Parsennii

<i>J. conferta</i>	Shore Juniper; Blue Pacific
<i>J. davurica</i>	Parsoni
<i>J. horizontalis</i>	Creeping Juniper
<i>Juglans sp.</i>	Walnut *
<i>Justicia brandegeana</i>	Shrimp Plant
<i>Lagerstromia indica</i>	Crape Myrtle
<i>Lagerstromia indica X Fauriei</i>	Crape Myrtle; Tuscarora
<i>Lantana montevidensis</i>	Weeping Lantana
<i>Lavender sp.</i>	Lavender; Munstead
<i>Leontopodium alpinum</i>	Edelweiss
<i>Ligustrum sinense</i>	Chinese Privet; Variegata
<i>Ligustrum amurense</i>	Amur Privet

* Do not use on food producing trees, vines, or plants.

** Not for use on container grown plants.

<u>Scientific Name</u>	<u>Common Name</u>
<i>L. japonicum</i>	Japanese Privet
<i>L. lucidum</i>	Glossy Privet (Waxleaf)
<i>Lillium sp.</i>	Lily: Jazz
<i>Liriope Muscari</i>	Big Blue Lily; Turf; Silver Mound; Evergreen Giant
<i>Liriope Muscari var. variegata</i>	Liriope, Variegated
<i>Liriope spicata</i>	Liriope, Creeping
<i>Lobelia Cardinalis</i>	Cardinal Flower, Indian Pink
<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>L. tatarica</i>	Tatarian Honeysuckle
<i>Loropetalum chinense</i>	Burgundy
<i>Lythrum sp.</i>	Loosestrife; Modern Pink
<i>Magnolia spp.</i>	Magnolia species **
<i>Maleophora luteola</i>	Ice Plant
<i>Malus sp.</i>	Crabapple *
<i>Miscanthus sinensis</i>	Yaku Jima, Silberfeder
<i>Nandina domestica</i>	Heavenly Bamboo
<i>Narcissus spp.</i>	Narcissus species **
<i>Nerium oleander</i>	Oleander
<i>Olea europaea</i>	Olive*
<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Osmanthus heterophyllus</i>	Osmanthus (False Holly): Gulf Tide
<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Oxydendrum arboreum</i>	Sourwood
<i>Paeonia suffruticosa</i>	Tree Peony
<i>Pennisetum setaceum</i>	Fountain Grass (Dwarf)
<i>Perovskia atriplicifolia</i>	
<i>Persea americana</i>	Avocado *
<i>Photinia Fraseri</i>	Fraser's Photinia (Redtip)

Physotegia Virginiana
Picea spp.
Pieris japonica
Pinus brutia
P. canariensis
P. elliotii
P. halepensis
P. nigra
P. palustris
P. radiata
P. Thunbergiana
P. strobus

Dragonhead, False; Vivid
 Spruce species ** (Colorado Blue, Norway, etc.)
 Lily-of-the-Valley Bush
 Calabrian Pine
 Canary Island Pine
 Slash Pine
 Aleppo Pine
 Austrian Pine
 Longleaf Pine
 Monterey Pine
 Japanese Black Pine
 Eastern White Pine

* Do not use on food producing trees, vines, or plants.
 ** Not for use on container grown plants.

Scientific Name

Common Name

P. sylvestris
P. taeda
P. virginiana
Pistachio sp.
Pittosporum rhombifolium
P. tobira
Podocarpus macrophyllus
Prunus Laurocerasus
Prunus sp.

Pseudotsuga Menziesii
Pyracantha coccinea
P. Fortuneana
P. Koidzumii
Pyrus sp.
Quercus Shumardii
Quercus spp.
Raphiolepis indica
Raphiolepis umbellata
Rhododendron
 (including Azalea)

Scotch Pine
 Loblolly Pine
 Virginia Pine
 Pistachio *
 Queensland Pittosporum
 Japanese Pittosporum
 Japanese Yew
 English Laurel
 Almond, Apricot, Nectarine,
 Peach, Plum and Prune *
 Douglas Fir **
 Firethorn (Scarlet)
 Firethorn
 Firethorn
 Bradford Pear sp.
 Oak, Shumard's Red
 Oak species
 Indian Hawthorne
 Yedda Hawthorn
 'Coral Bells'
 'Delaware Valley White'
 'Flame Creeper'
 'Formosa'
 'Girard Crimson'
 'George L. Tabor'
 'Hino-crimson'
 'PJM'
 'Roseum Elegans' *
 'Wakeiebisu'

<i>Rosa Banksiae</i>	'White Gumbo'
<i>Rosmarinus officinalis</i>	Lady Bank's Rose
<i>Rudbeckia sp.</i>	Rosemary*
<i>Santolina virens</i>	Black-Eyed Susan; Goldstrum
<i>Saxifraga sp.</i>	Santolina
<i>Scabiosa sp.</i>	Saxifrage; Purple Dome
<i>Sedum album</i>	Pincushion Flower
<i>Sedum cauticola</i>	Stonecrop
<i>Sedum dasyphyllum</i>	Stonecrop; Lidakense
<i>Sedum spurium</i>	Stonecrop
<i>Spiraea Bumalda</i>	Stonecrop; Dragon's Blood
	Spirea: Anthony Waterer

* Do not use on food producing trees, vines, or plants.
 ** Not for use on container grown plants.

<u>Scientific Name</u>	<u>Common Name</u>
<i>Syzygium paniculatum</i>	Australian Brushcherry
<i>Taxus cuspidata</i>	Japanese Yew
<i>T. media</i>	Yew
<i>Teucrium sp.</i>	Germander
<i>Thalictrum d'pterocarpum</i>	Meadow Rue
<i>Thuja occidentalis</i>	American Arborvitae
<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Tsuga canadensis</i>	Canada Hemlock
<i>Tulipa spp.</i>	Tulip species
<i>Veronica sp.</i>	Veronica, Speedwell; Sunny Border
<i>Viburnum japonicum</i>	Japanese Viburnum
<i>V. odoratissimum</i>	Sweet Viburnum
<i>V. plicatum</i>	Japanese Snowball
<i>V. rigidum</i>	Canary Island Viburnum
<i>V. suspensum</i>	Arrowood Viburnum
<i>V. tinus</i>	Laurustinus
<i>V. trilobium</i>	Cranberry Bush
<i>V. wrightii</i>	Leatherleaf Viburnum
<i>Vinca major</i>	Greater Periwinkle, Vinca
<i>Vinca minor</i>	Common Periwinkle, Vinca
<i>Vitis sp.</i>	Grape*
<i>Weigela florida</i>	Old Fashioned Weigela
<i>Yucca aloifolia</i>	Spanish Bayonet, Yucca
<i>Y. filamentosa</i>	Adam's Needle, Yucca

* Do not use on food producing trees, vines, or plants.
 ** Not for use on container grown plants.

19 of 19

LIMITATION OF WARRANTY AND LIMITATION OF LIABILITY

NOTICE: Read this Limitation of Warranty and Limitation of Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of The Manufacturer or Seller. All such risks shall be assumed by Buyer or User.

The Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, under normal use conditions, subject to the risks described above. **THE MANUFACTURER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.** In no event shall The Manufacturer or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product.

The Manufacturer and Seller offer this product, and Buyer and User accept it, subject to the foregoing limitations of warranty and limitation of liability, which may not be modified by any oral or written agreement.