

961-367

2/9/2009

1/6



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

9 FEB 2009

Richard J. Newmaster, Jr.
Lebanon Seaboard Corporation
1600 East Cumberland Street
Lebanon, PA 17042

Dear Mr. Newmaster:

Subject: Alternate Brand Name
Lebanon Pro Brand Fertilizer with Barricade 0.29% Preemergence Weed Control
EPA Registration Number 961-367
Application Dated November 5, 2008

The label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) as amended is acceptable. If you have any questions, please contact Tracy White by phone at (703) 308-0042 or via email at white.tracy@epa.gov.

Sincerely,

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

Enclosure

Lebanon Pro



FERTILIZER WITH BARRICADE® 0.29% PREEMERGENCE WEED CONTROL

12-3-5



contains **METH-EX**

22-41267

GUARANTEED ANALYSIS

Total Nitrogen (N)	12%
1.2% Ammoniacal Nitrogen	
0.7% Water Insoluble Nitrogen*	
8.8% Urea Nitrogen	
1.3% Other Water Soluble Nitrogen*	
Available Phosphate (P ₂ O ₅)	3%
Soluble Potash (K ₂ O)	5%
Derived from: Ammonium Phosphate, Methylene Ureas, Urea, Muriate of Potash.	
Chlorine (Cl) not more than	5.0%
*2.0% Slowly Available Nitrogen from Methylene Ureas.	F699

SUGGESTED SPREADER SETTINGS

Spreader	5.2# 1000 Sq. Ft. (65# A/Acre)	Spreader	5.2# 1000 Sq. Ft. (65# A/Acre)
Lebanon Turf	5%	Pfizerblam CBR II	6
Cyclone	5%	Pfizerblam LF-1	4%
Ezrae Flow (109)	A4	ProScape SS	N
Gandy	30	Republic Rotary	6
Lely (HR, W 1250, 1500, 2018)	6%II	Scotts AccuPro	0
Leuco (Bellshumard)	K/22	Scotts FDX	J
Ortha Rotary	6	Scotts FBA	N
PennMach HYD	N	Spylar	5%
Pfizerblam DF & CBR II	N	Vicon (all models)	30

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

- For selective preemergence control of grass and broadleaf weeds in:
- established turfgrasses (excluding golf course putting greens) and lawns
 - landscape ornamentals (including established perennials & wildflower plantings)

ACTIVE INGREDIENT:

Pridamine	0.29%
OTHER INGREDIENTS:	99.71%
TOTAL	100.00%

EPA Reg. No. 901-387

EPA Est. No. 901-PA-01901-L-01
Superscript used to list letter of lot number.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.
Before buying this product, read the Warranty Section elsewhere on this label.
If terms are not acceptable, return the unopened container to the place of purchase at once.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye injury (irritation). Handle if labeled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Redwood or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

FIRST AID	
IF IN EYES:	• Hold eye open and draw slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN:	• Take all contaminated clothing off.
IF ON CLOTHING:	• Remove and isolate immediately with a bag of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance. Then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further advice on action.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact (800) 368-5868 for more product safety information.

ENVIRONMENTAL HAZARDS

This product has low toxicity to fish. At the level of toxicity, fish in ponds or streams are not at risk. However, if concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water. Do areas where surface water is present or to water bodies before the same high water mark. Drain equipment from treated areas may be hazardous to aquatic organisms in adjacent areas. Do not contaminate water when disposing of equipment washwaters.

If material is spilled, contain and sweep up material or spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination. Excess quantities should be disposed of in accordance with applicable regulations. Do not dump in drains, subject to pH. Waste in sewer, discharge, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.



For technical assistance or more information about our products visit www.LebanonTurf.com

Manufactured by:
Lebanon Seaboard Corporation
1600 E. Cumberland St. • Lebanon, PA 17042
Made in the U.S.A.
www.LebanonTurf.com

NET WT. 50 LBS. (22.7 kg)

9 FEB 2009
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

961-367

FERTILIZER WITH BARRICADE® 0.29% PREEMERGENCE WEED CONTROL

12-3-5



contains **METH-EX®**

22-41267


GUARANTEED ANALYSIS

Total Nitrogen (N)	12%
1.2% Ammoniacal Nitrogen	
0.7% Water Insoluble Nitrogen*	
8.8% Urea Nitrogen	
1.3% Other Water Soluble Nitrogen*	
Available Phosphate (P ₂ O ₅)	3%
Soluble Potash (K ₂ O)	5%
Derived from: Ammonium Phosphate, Methylene Ureas, Urea, Muriate of Potash.	
Chlorine (Cl) not more than	5.0%
*2.0% Slowly Available Nitrogen from Methylene Ureas.	F699

SUGGESTED SPREADER SETTINGS

Spreader	5.2#/1000 Sq. Ft. (.65# AI/Acre)	Spreader	5.2#/1000 Sq. Ft. (.65# AI/Acre)
Lebanon Turf	5½	Prizelawn CBR II	6
Cyclone	5½	Prizelawn LF-1	4½
Ezee Flow (109)	A4	ProScape SS	N
Gandy	30	Republic Rotary	6
Lely (HR, W 1250, 1500, 2010)	6½/11	Scotts AccuPro	O
Lesco (letter/numeric)	K/22	Scotts R7X	J
Ortho Rotary	6	Scotts R8A	N
PennMulch HVO	N	Spyker	5½
Prizelawn BF & CBR III	N	Vicon (all models)	30

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.



For selective preemergence control of grass and broadleaf weeds in:

- established turfgrasses (excluding golf course putting greens) and lawns
- landscape ornamentals (including established perennials & wildflower plantings)

ACTIVE INGREDIENT:

Prodiamine 0.29%

OTHER INGREDIENTS: 99.71%

TOTAL 100.00%

EPA Reg. No. 961-367
 Barricade is a registered trademark of a Syngenta Group Company.

(L) (D)
 EPA Est. No. 961-PA-01:961-IL-01
 Superscript used is first letter of lot number.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label.

If terms are not acceptable, return the unopened container to the place of purchase at once.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

FIRST AID

- | | |
|-------------------------------------|---|
| IF IN EYES: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| IF ON SKIN:
OR CLOTHING: | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| IF INHALED: | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice. |

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact (888) 209-1368 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

If material is spilled: Contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.



For technical assistance or more information about our products visit www.LebanonTurf.com

Manufactured by:

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042

Made in the U.S.A.

www.LebanonTurf.com



NET WT. 50 LBS. (22.7 kg)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turfgrasses (excluding golf course putting greens) and lawns, landscape ornamentals, established perennials and wildflower plantings.

This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not graze or feed livestock forage cut from areas treated with this product.

Do not apply this product through any type of irrigation system.

Do not apply aerially.

Do not apply to golf course putting greens.

APPLICATION DIRECTIONS

Apply uniformly with suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turfgrasses and lawns. The maximum amount of this product that may be applied per year is given for each turfgrass species in the Maximum Annual Rates section of this label.

Most effective weed control in turfgrasses will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

CRABGRASS SEED GERMINATION DATES



- Approximate Date
- After May 30
 - After May 10
 - After April 20
 - After March 20
 - January 1 to March 20

USE PRECAUTIONS

The following precautions apply to the use of this product in turfgrasses and lawns:

Application of this product may thin emerged annual bluegrass and newly overseeded grasses.

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year.

Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.

Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.

Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES

This product is recommended for use on the turfgrass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

MAXIMUM APPLICATION RATE/CALENDAR YEAR of LEBANON PROSCAPE FERTILIZER WITH BARRICADE 0.29% PREEMERGENCE WEED CONTROL by Turfgrass Species¹⁾

Turf Species:	Lbs. Product/Acre	Lbs. AI/Acre	Lbs./1,000 Sq. Ft.
Bermudagrass ²⁾ , Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type), Zoysia	518	1.5	11.6
Buffalograss, Kentucky Bluegrass, Perennial Ryegrass	345	1.0	7.8
Creeping Red Fescue	260	0.75	6.0
Creeping Bentgrass	225	0.65	5.2

¹⁾ These are the maximum rates per calendar year by species limitations.

²⁾ May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 179 lbs./Acre. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

• Do not apply more than 518 lbs. of this product per calendar year.

• Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates Table.

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

Barnyardgrass	Chickweed, Mouseear (from seed)	Foxtails, Annual	Johnsongrass (from seed)	Lambsquarters, Common	Purslane, Common	Speedwell, Peter
Bluegrass, Annual (<i>Poa annua</i>)	Crabgrass (large, smooth)	Goosegrass ¹⁾	Junglerice	Lovegrass	Pusley, Florida	Sprangletop
Carpetweed	Crowfootgrass	Herbit	Kochia	Panicum (Texas, Fall, Browntop)	Rescuegrass ²⁾	Spurge, Prostrate
Chickweed, Common	Cupgrass, Woolly	Itchgrass	Knotweed	Pigweed	Shepherd's Purse ³⁾	Witchgrass
					Signalgrass, Broadleaf	Woodsorel, Yellow (from seed)

¹⁾ In many areas a single application of 225 to 518 lbs./Acre of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 225 to 345 lbs./Acre followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rate Table. Do not exceed the maximum rate for turfgrass species listed in the Maximum Annual Rate Table above.

²⁾ Applications for this weed should be made in late summer, fall or winter prior to germination.

³⁾ Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

PREEMERGENCE WEED CONTROL RATE (lb./A)

Rate (lb./A)	North	Months Before Transplanting	South
172	4	4	4
225	5	4	4
260	6	5	5
275	---	6	6
345	---	7	7
393	---	---	9
448	---	---	10
518	---	---	12

LANDSCAPE ORNAMENTALS (Including Established Perennials and Wildflower Plantings)

This product may be applied for residual preemergence weed control in ornamentals.

USE RATES

Apply this product at 225 to 518 lbs./Acre in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 518 lbs./Acre.

APPLICATION TIMING AND INFORMATION

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

ESTABLISHED LANDSCAPE ORNAMENTALS

This product may be applied at any time to established ornamentals as a broadcast, over the top or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

NEWLY TRANSPLANTED LANDSCAPE ORNAMENTALS

This product may be applied to newly transplanted ornamentals as a broadcast, over the top or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species.

When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

Fir species** (Balsam, Fraser, Noble, etc.)	Feathery Cassia	Gaura sp.	Edelweiss	Canary Island Pine	'White Gumbo'
Abelia: Sherwood	Wild Lilac	Genlian	Chinese Privet: Variegata	Slash Pine	Lady Bank's Rose
Japanese Maple	Ceratostigma plumbagonoides	Cranesbill	Amur Privet	Alleppo Pine	Rosemary
Norway Maple	False Cypress	Gladiolus species**	Japanese Privet	Austrian Pine	Black-Eyed Susan; Goldstrum
Yarrow: King Edward	<i>Chrysanthemum nipponicum</i>	Baby's Breath	Glossy Privet (Waxleaf)	Longleaf Pine	Santolina
Kiwi*	Clayera	English Ivy	Lily Jazz	Monteary Pine	Saxifrage; Purple Dome
Lily-of-the-Nile (African Lily)	Citrus species*	Sunrose	Big Blue Lily; Turf; Silver Mound;	Japanese Black Pine	Pincushion Flower
<i>Agapanthus orientalis</i>	Coreopsis (Calliopsis): Early	Daylily: Aztec Gold, Stella De Oro,	Evergreen Giant	Eastern White Pine	Stoncrop (<i>Sedum album</i>)
Five-Leaf or Chocolate Vine	Sunrise, Moonbeam	Tender Love	Liriope, Variegated	Scotch Pine	Stoncrop; Lidakense
Lady's Leek, Nodding Onion	Flowering Dogwood	Rose of Sharon**	Liriope, Creeping	Loblolly Pine	Stoncrop (<i>Sedum dasycyllum</i>)
Japanese Anemone	American Dogwood	Mallow: Disco Belle White	Cardinal Flower, Indian Pink	Virginia Pine	Stoncrop; Dragon's Blood
Aquilegia: Red and Gold	Pampas Grass	Chinese Hibiscus	Japanese Honeysuckle	Pistachio*	Spiraea: Anthony Waterer
Vine Hill Manzanita	Cranberry Cotoneaster	Hosta, Plantain Lily (Fragrant)	Tatarian Honeysuckle	Queensland Pittosporum	Australian Brushcherry
Cape Weed	Cotoneaster	Hosta, "Seasucker"	Burgundy	Japanese Pittosporum	Japanese Yew
Wormwood, Silver Mound, Castle	Boaberry Cotoneaster	<i>Houttuynia cordata var. variegata</i>	Loosestrife, Modern Pink	Japanese Yew	Yew
Aster: Bonny Blue, Purple Dome	Rockspray Cotoneaster	Bigleaf Hydrangea	Magnolia species**	English Laurel	Gemmander
Aster X <i>Frikarti</i>	Hawthorne	Holly	Ice Plant	Almond, Apricot, Nectarine, Peach,	Meadow Rue
Lady Fern; Fern Lady	Ludlow	Chinese Holly	Crabapple*	Plum and Prune*	American Arborvitae
Japanese Aucuba	Italian Cypress	Japanese Holly; Helleri	Yaku Jima, Silberfeder	Douglas Fir**	Star Jasmine
Fibrous Begonia: Hardy Grandis	White Trailing Ice Plant	American Holly	Heavenly Bamboo	Firethorn (Scarlet)	Canada Hemlock
Barberry	Cooperi Pink	Yaupon Holly; Schottings	Narcissus species**	Firethorn (<i>P. Fortuneana</i>)	Tulip species
Wintergreen Barberry	Larkspur: Blue Elf	<i>Inula ensifolia</i>	Oleander	Firethorn (<i>P. Koizumii</i>)	Veronica, Speedwell; Sunny Border
Mentor Barberry	Dianthus: Maiden Pinks "Zing"	Sword-Leaved Iris; Jodlesong	Olive*	Bradford Pear sp.	Japanese Viburnum
Japanese Barberry	Cheddar Pink	Siberian Iris; Cabernet	Mondo Grass	Oak, Shumard's Red	Sweet Viburnum
Warty Barberry	Hop Bush	Iris species**	Osmanthus (False Holly): Gulf Tide	Oak species	Japanese Snowball
<i>Bergenia cordifolia</i>	Coneflower, Purple; Magnus	Winter Jasmine	Trailing African Daisy	Indian Hawthorne	Canary Island Viburnum
Snowbank	Silverberry	Chinese Juniper; Nick's Compact,	Sourwood	Yedda Hawthorn	Arowood Viburnum
Butterfly-Bush (Dwarf Blue);	Wintercreeper	Parsonii	Tree Peony	'Coral Bells'	Laurustinus
Royal Red	Japanese Spindle Tree (Evergreen	Shore Juniper; Blue Pacific	Fountain Grass (Dwarf)	'Delaware Valley White'	Craberry Bush
Japanese Boxwood	Eunonymus)	Parsoni	<i>Perovskia atriplicifolia</i>	'Flame Creeper'	Leatherleaf Viburnum
Weeping Bottlebrush	Spreading Eunonymus	Creeping Juniper	Avocado*	'Fomosa'	Greater Periwinkle, Vinca
Scotch Heather	Japanese Aralia	Walnut*	Fraser's Photinia (Redtip)	'Girard Crimson'	Common Periwinkle, Vinca
Tussock Bellflower, (White Clips)	Border Forsythia	Shrimp Plant	Dragonhead, False, Vivid	'George L. Tabor'	Grape-Old Fashioned Weigela
Trumpet Creeper, Trumpet Flower,	Weeping Forsythia	Crape Myrtle	Spruce species** (Colorado Blue,	'Hino-crimson'	Spanish Bayonet, Yucca
Madame Galen	Greenstem Forsythia	Crape Myrtle; Tuscarora	'Norway, etc.)	'PJM'	Adam's Needle, Yucca
Hottentot Fig (Ice Plant)	Gaillardia, Blanket Flower: "Goblin"	Weeping Lantana	Lily-of-the-Valley Bush	'Roseum Elegans'	
	Gardenia, Cape-Jasmine, Padiicans	Lavender; Munstead	Calabrian Pine	'Wakelebisu'	

* Do not use on food producing trees, vines, or plants. ** Not for use on container grown plants.

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum of 1 pound nitrogen per 1,000 sq. ft.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store product in its original container in a cool, dry, locked place out of the reach of children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

LIMITATION OF WARRANTY AND LIMITATION OF LIABILITY

NOTICE: Read this Limitation of Warranty and Limitation of Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of The Manufacturer or Seller. All such risks shall be assumed by Buyer or User.

The Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, under normal use conditions, subject to the risks described above. THE MANUFACTURER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. In no event shall The Manufacturer or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product.

The Manufacturer and Seller offer this product, and Buyer and User accept it, subject to the foregoing limitations of warranty and limitation of liability, which may not be modified by any oral or written agreement.

MANUFACTURED BY:

LEBANON SEABOARD CORPORATION

1600 EAST CUMBERLAND STREET
LEBANON, PA 17042

For technical assistance or more information about our products visit www.LebanonTurf.com

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-lebsea.com>