

U.S. ENVIRONMENTAL ENCIRECTION AGENCY Office of Perticide Programs Registration Civision (7505C) 401 TMT St., S.W. Washinton, D.G. 20460

NOTICE OF PESTICIDE:

X Registration
Reregistration

(under FIFRA, as amended)

EPA Req

Clark of Instance

961-391

JAN 12 2001

Term of Tas Annel

Conditional

Name of Festicide Froduct:

Lebanon Turf Fertilizer with Confront and Team

Name and Address of Registrant (include 219 Code):

Lebanon Seaboard Corporation

1600 E. Cumberland Street

Lebanon, Pennsylvania 17042

Note: Changes in labeling differing in surstance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Submit/cite all data required for the registration/reregistration of your product when the Agency requires all registrants of similar products to submit data.
 - 2. Make the following label changes:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. 961-391".
 - b. Under the subheading entitled "Hazards to Humans and Domestic Animals," revise the first sentence of the paragraph to read as "Causes moderate eye irritation".
 - c. Under the heading entitled "DIRECTIONS FOR USE" within the section subtitled "General Use Precautions and Restrictions," incorporate the following application and entry restrictions:

Application Restriction: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Signature of Approving Official:

Date:

JAN 12 2001

EPA Form 857**0**−6

MINISON

page 2 EPA Reg. No. 961-391

Entry Restrictions: Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by water-in, do not enter or allow others to enter the treated area until the surface is dry after water-in.

3. Submit (2) two copies of the revised final printed label for the record.

A stamped copy of the label is enclosed for your records.

If these conditions are not complied with, the registration will be subject to concellation in accordance with FIFRA sec. 6(e). Your release for shipment fo the product constitutes acceptance of these conditions

Joanne I. Miller Product Manager (23) Herbicide Branch Registration Division (7505C)

Enclosures

LEBANON TURF FERTILIZER WITH CONFRONT* AND TEAM*

Active Ingredients:	
triclopyr: 3,5,6 trichloro-2-pyridinyloxyacetic acid, triethylamine salt	0.50%
clopyralid: 3,6-dichloro-2-pyridinyloxyacetic acid, triethylamine salt	0.18%
benefin: N-butyl-N-ethyl-α,α,α-triffuoro-2,6-dinitro-p-toluidine	0.82%
trifluralin: (α,α,α-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine)	0.43%
Inert Ingredients	
Total	
a campana a a cala a campana	

Acid Equivalent: triclopyr - 0.36% clopyralid - 0.12%

EPA Reg. No. 961-XXX

EPA Est. 961-PA-01: 961-IL-01
Superscript used is first letter of lot number

*Trademark of Dow AgroSciences LLC

GUARANTEED ANALYSIS			
Total Nitrogen	X.X%		
X.X% Ammoniacal Nitrogen			
X.X% Water Insoluble Nitrogen			
X.X% Urea Nitrogen			
X.X% Other Water Soluble Nitrogen			
Available Phosphate (P2O5)	X.X%		
Soluble Potash (K2O)			

Nutrient Sources: (Guaranteed Analysis and Nutrient Sources to be supplied based on fertilizer analysis used by manufacturer.)

KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel for Additional Precautionary Statements.

MANUFACTURED BY:
LEBANON SEABOARD CORPORATION
1600 EAST CUMBER LAND STREET
LEBANON, PA 17042
www.lebses.com

ACCEPTED with COMMENTS In EPA Letter Dated

JAN 12 2001

Under the Federal Insecticide, Fundicide, and Redenticide Act as amended, for the posticide registered under EPA Reg. No. 96/-39/ NET WEIGHT: XX LBS. (X.XX kg) COVERS: X,XXX sq ft Lebanon Turf Fertilizer with Confront and Team / Proposed Section 3 / 02-28-00

Page 2

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals CAUTION

Causes substantial but temporary eye injury. Harmful if absorbed through the skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause silergic reaction in some individuals.

Personal Protective Equipment:

Long-sleeved shirt and long pants • Chemical-resistant gloves • Shoes plus socks • Protective eyewear (goggles, face shield or safety glasses)

User Safety Requirements: Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the WPS (40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

- Users should wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the tollet.
- Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If on eldin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including strimp and cyster. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate ground water which may be used for imigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an over limestone bedrock, severely fractured surfaces, and substrates which could allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

(Back Panel of Label)

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use before applying.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container only in a clean, dry area away from extremes of heat and cold and out of direct sunlight.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

Lebanon Turf Fertilizer with Confront* herbicide and Team* herbicide provides broad-spectrum control of broadleaf weeds and annual grasses in established cool season or warm season turfgrasses. Confront herbicide is particularly effective against established broadleaf weeds such as dandelions and Team herbicide prevents the establishment of annual grass weeds, including crabgrass. This product is not effective against established grass weeds.

Lebanon Turf Fertilizer with Confront and Team is recommended for the following turf species:

Established Cool Season Turfgrasses	Established Warm Season Turf grasses †
Kentucky bluegrass	Bahiagrass (Argentine or Pensacola)
chewings fescue	Common Bermudagrass
creeping red fescue	Buffalograss
sheeps fescues	Centipedegrass
tall fescue	Zoysiagrass (Zoysia japonica, Zoysia tenuifolia)
perennial ryegrass	Tall Fescue (growing in warm season area).

[†] Do not treat warm season turfgrasses with Lebanon Turf Fertilizer with Confront and Team when mowing height is less than 1/2 inch. Do not use Lebanon Turf Fertilizer with Confront and Team on got course putting greens and tees.

General Use Precautions and Restrictions

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

\Box	Do not apply this product to bentgrass; hybrid bermudagrass varieties such as Tifway,
	Tifdwarf, and Tifgreen; St. Augustinegrass; and ground covers such as lippia.
	This product may discolor and/or stunt turf that is not well established or is stressed or weakened due
	to unfavorable climatic conditions, temperature extremes, drought, nematodes, or other factors which
	damage or weaken turf. Apply only to healthy, well-established turf that has a well-anchored root
	system.
	Common harmydaerana, cantinadaerana, zawiaerana, and habiaerana may ha tamparanily discolared

Common bermudagrass, centipedegrass, zoysiagrass, and bahiagrass may be temporarily discolored by this product.

- On bahiagrass and centipedegrass, do not apply before mid-March or until after the second mowing of the year. In the fall, do not treat centipedegrass after the first of October or bahiagrass after the first of November.
- Do not apply this product to newly seeded turf until turfgrass is fully established. Do not seed, sod, or sprig after application. Do not reseed until 12 weeks after application. Delay sodding or sprigging until 4 weeks after application.
- Do not use grass clippings from turf treated with this product for mulch. Do not use compost containing grass clippings from turf treated with this product in the growing season of application. Do not graze treated areas or feed clippings to livestock.
- Do not apply this product directly to, or otherwise permit it to come into direct contact with vegetables, flowers, fruit or ornamental frees, cotton, grapes, tobacco, or other nontarget broadleaf plants. Do not apply when weather conditions favor drift from target areas. Drift or runoff may adversely affect nontarget plants.
- Do not apply this product to exposed roots of shallow rooted trees and shrubs or allow contact with exposed suckers or injury may occur. Do not apply at greater than recommended rates, especially under the branch spread (dripline) of trees.
- Do not contaminate water used for irrigation or domestic purposes. Do not apply on ditches used to transport irrigation water. Do not apply where runoff or irrigation water may flow onto susceptible crops as injury may result.

Weeds Controlled

When applied as directed, Lebanon Turf Fertilizer with Confront and Team will control the following weeds:

Annual Grass Weeds	Established Broadleaf Weeds		
Annual Grass Weeds crabgrass (smooth and hairy) goosegrass barnyardgrass foxtail annual bluegrass (Poa annua)	black medic bristly hawksbeard broadleaf plantain burdock Canada thistle catsear (coast dandelion) coffee weed common chickweed common cocklebur common dandelion common ragweed common vetch common yellow woodsorrel	curly dock henbit dollarweed dwarf beggarweed English lawn daisy false dandelion goldenrod ground ivy hawkweed hop clover lambsquarters lespedeza matchweed musk thistle	old world diamond flower poison ivy prostrate spurge red clover roundleaf mailow sheep sorrel shepherdspurse slender speedwell smartweed spurweed Virginia buttonweed Virginia pepperweed white clover wild buckwheat wild violet
	creeping beggarweed creeping woodsorrel	narrow leaf plantain (buckhorn)	

Note: Moist foliage application will provide quicker results on broadleaf weeds. For best herbicide effect when applying to dry foliage, apply when good soil moisture conditions are present.

A repeat application may be needed for harder to control weeds in 4 to 6 weeks.

APPLICATION DIRECTIONS

How to Apply Lebanon Turf Fertilizer with Confront and Team

Apply Lebanon Turf Fertilizer with Confront and Team at a recommended rate of 3.6 lb per 1,000 sq ft (157 lb/acre). This application rate is equivalent to 0.75 lb ae/acre of Confront and 2 lb ai/acre of Team. Make application using equipment that will ensure uniform coverage. For postemergence control of broadleafs, application should be made when weeds are actively growing.

Good soil moisture conditions should be present before application to achieve optimal results. Application under drought conditions may provide less than desirable results.

Broadleaf weed species germinate at different times. Emerged broadleaf weeds present at time of application are controlled by Confront, unemerged grassy weeds present at time of application are controlled by Team.

Note: Additional applications should not be made less than 4 weeks apart to minimize the potential for grass injury. Newly seeded turf should be moved 4 times before being treated.

Suggested Spreader Settings †

[Editor's Note: Manufacturer to supply a listing of spreaders and suggested settings to apply this product at a rate of 3.6 lbs. per 1,000 sq ft.]

[†] The above spreader settings are approximate. These settings were calibrated and field-tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

 	-		
WA	R	RANTY	

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to Manufacturer and Buyer assumes the risks of any such use.

*Trademark of	Dow AgroSciences	LLC
EPA-accepted		