



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg.
Number:

961-395

Date of Issuance:

JAN 25 2005

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Confront III + Fertilizer

Name and Address of Registrant (include ZIP Code):

Lebanon Seaboard Corporation
1600 East Cumberland Street
Lebanon, PA 17042

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label change:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. 961-395".
 - b. Remove and delete all references to use on sod farms.
 - c. Under the headings entitled "PRECAUTIONARY STATEMENTS" on the commercial and residential product labeling, please revise the precautionary statements to read as the following:

"Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long-sleeved shirt and long pants, socks, shoes, chemical-resistant gloves (such as Natural Rubber, Selection Category A) and protective eyewear (goggles, face shield or safety glasses) when applying. Remove and wash contaminated clothing before reuse."

Signature of Approving Official:

Joanne J. Miller

Date:

JAN 25 2005

d. Within the box entitled "User Safety Recommendations" on the commercial and residential product labeling, please include the following statement: "Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing."

3. Submit, within one (1) year of the date of this pesticide registration notice, the following product chemistry data: one year storage stability study (830-6317) and corrosion characteristics study (830-6320).

4. Submit one (1) copy of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosures

CONFRONT* III + FERTILIZER

For the postemergence control of annual and perennial broadleaf weeds in established ornamental turf, such as lawns, golf courses, roadsides and cemeteries.

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-D†	1.037%
Triethylamine salt of triclopyr.....	0.303%
Methylheptyl ester of fluroxypyr.....	0.207%

OTHER INGREDIENTS	98.453%
TOTAL	100.000%

Acid Equivalents:

2,4-dichlorophenoxyacetic acid – 0.860% 17.20 lb/ton	2, 4-D CAS. No. 94-75-7
triclopyr: 3,5,6-trichloro-2-pyridyloxyacetic acid – 0.217% 4.34 lb/ton	Triclopyr CAS. No. 55335-06-3
fluroxypyr: 4-amino-3,5-dichloro-6-fluoro-2-pyridyloxyacetic acid – 0.143% 2.86 lb/ton	Fluroxypyr CAS. No. 69377-81-7

†Isomer Specific by AOAC Method No. 6.275-6.279 (13th Ed.)

EPA File Symbol 961-XXX

(L) (D)
EPA Est. 961-PA-01: 961-IL-01

*Trademark of Dow AgroSciences LLC.

Superscript used is first letter of lot number.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See back panel for additional Precautionary Statements

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label.

If terms are not acceptable, return the unopened container to the place of purchase at once.

MANUFACTURED BY:
LEBANON SEABOARD CORPORATION
1600 EAST CUMBERLAND STREET
LEBANON, PA 17042
www.LebanonTurf.com

GUARANTEED ANALYSIS

Total Nitrogen (N).....	X.X%
X.X% Ammoniacal Nitrogen	
X.X% Water Insoluble Nitrogen	
X.X% Urea Nitrogen	
X.X% Other Water Insoluble Nitrogen	
Available Phosphate (P ₂ O ₅).....	X.X%
Soluble Potash (K ₂ O).....	X.X%

Nutrient Sources: (Guaranteed Analysis and Nutrient Sources to be supplied based on fertilizer analysis used by manufacturer.)

NET WEIGHT: XX.XX LBS. (XX.XX kg)

COVERS: X,XXX Sq. Ft.

ACCEPTED
with COMMENTS
In EPA Letter Dated:
JAN 25 2005
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

961-395

4/11

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye irritation. Harmful if swallowed or absorbed through the skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long-sleeved shirt, chemical-resistant gloves, shoes plus socks, and protective eyewear (goggles, face shield or safety glasses) when applying.

User Safety Recommendations

Users should:

- Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-888-208-1368 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Drift or runoff from treated areas may be hazardous to aquatic organisms and non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION OR FOR RESEARCH PURPOSES.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store this product only in its original container in a cool, dry, secured storage area out of the reach of children. Do not store in or around the home. In case of spill, carefully sweep up material and use as directed, or dispose of material according to pesticide disposal directions below.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

Confront III + Fertilizer provides broad-spectrum postemergence control of annual and perennial broadleaf weeds in established turf, including residential lawns, golf courses, sports fields, sod farms, roadsides, cemeteries and around commercial buildings and other commercial and residential turf areas. Apply during warm weather when weeds are actively growing. Application under drought conditions may provide less than desirable results. Applications to moist foliage helps granules adhere to weed surface and will improve control.

USE PRECAUTIONS AND RESTRICTIONS

Be sure that use of this product conforms to all applicable regulations.

Restrictions on Retreatment: Do not apply more than 2 broadcast applications per year per treatment site. To minimize the potential for grass injury, additional applications should not be made within 4 weeks of a previous application.

Do not use Confront III + Fertilizer on golf course putting greens or tees.

Do not use on grass species, such as bentgrass or St. Augustinegrass, unless injury can be tolerated.

Avoid off target application. Applications should be made so as to avoid spreading material off target. Avoid getting granules in shrub bed or other areas where they may seriously injure susceptible plants.

Do not apply to exposed suckers or roots of shallow rooted trees or shrubs.

Do not apply Confront III + Fertilizer directly to, or otherwise permit it to come into direct contact with cotton, grapes, tobacco, vegetable crops, flowers, fruit or ornamental trees, or other desirable broadleaf plants.

Do not apply under conditions that may have a potential to run-off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no-till, limited till and contour plowing; these methods also reduce pesticide run-off. Use of vegetation filter strips is recommended along rivers, creeks, streams, wetlands, etc.

Do not apply on ditches used to transport irrigation water. Do not apply where runoff or irrigation water may flow onto susceptible crops as injury may result.

Confront III + Fertilizer is recommended for use on the following established turfgrass species:

Common Name	Scientific Name
bentgrass ⁽¹⁾	<i>Agrostis species</i>
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, chewing	<i>Festuca rubra var. commutata</i>
fescue, creeping red	<i>Festuca rubra</i>
fescue, sheeps	<i>Festuca ovina</i>
fescue, tall	<i>Festuca arundinaceae</i>
ryegrass, perennial	<i>Lolium perenne</i>

Established Warm Season Turf Grasses ⁽¹⁾

Common Name	Scientific Name
Bahiagrass	<i>Paspalum notatum var. sauræ parodi</i>
Bermudagrass	<i>Cynodon dactylon</i>
Centipedegrass	<i>Eremochloa ophiuroides</i>
Zoysiagrass	<i>Zoysia japonica</i>
Zoysiagrass	<i>Zoysia tenuifolia</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
fescue, tall (growing in warm season areas)	<i>Festuca arundinaceae</i>

⁽¹⁾ Use Confront III + Fertilizer on these species only at the 150 lbs. per acre (3.5 lbs. per 1,000 sq. ft.) rate and only if some injury can be tolerated.

For use of Confront III + Fertilizer on a turf species not recommended on this label, the user may determine the suitability for such uses by treating a small area at a recommended rate. Prior to treatment of larger areas, the treated area should be observed for any sign of herbicidal injury during 30 days of normal growing conditions to determine if the treatment is safe to the target turf species. The user assumes the responsibility for any plant damage or other liability resulting from use of Confront III + Fertilizer on turf species not recommended on this label.

6/11

WEEDS CONTROLLED AND APPLICATION RATES

Weeds Controlled		Rate/Acre	Rate/1000 Sq. Ft.
bedstraw black medic buttonweed, Virginia carpet weed catchweed catnip chamise chickweeds chicory cinquefoil clovers cocklebur coffee weed cornflower creeping beggarweed dandelion dayflower deadnettle, purple docks dwarf beggarweed field bindweed goldenrod	henbit hop clover knawel lambsquarters lespedeza little starwort mallow matchweed plantain purslane speedwell smartweed sowthistle spiderwort spotted catsear spurweed velvetleaf vetch woodsorrel, common yarrow	150 lb.	3.5 lb.
burdock buttercup knotweed mustard	poison ivy poison oak wild carrot	174 lb.	4.0 lb.
comspeedwell ground ivy oxalis (stricta and comiculata) veronica species (suppression only)	parsley-piert prostrate spurge wild violet	174 lb. (2 applications, 4 to 6 weeks apart; apply to moist foliage)	4.0 lb. (2 applications, 4 to 6 weeks apart; apply to moist foliage)

LABELED USES

TURF

Broadcast Treatment of Ornamental Turf

Apply 150 to 174 lbs. per acre (3.5 to 4 lbs. per 1,000 sq. ft.) of Confront III + Fertilizer to control broadleaf weeds growing in established turf. Apply from early spring through fall when weeds are actively growing. Broadleaf weed species germinate at different times. Only emerged weeds present at time of application are controlled. Additional applications should be made four weeks apart to minimize grass injury. Newly seeded turf should be mowed two or three times before being treated. Do not water for 24 hours after application. Do not reseed for 3 weeks after application.

APPLICATION INSTRUCTIONS

For best results, apply this product evenly and uniformly over the turfgrass area, avoiding streaking, skips or overlaps. Before each application, calibrate the spreader according to the equipment manufacturer's directions. Adjust the spreader so the spreader delivers the appropriate application rate as recommended above.

SUGGESTED SPREADER SETTINGS

Spreader 3.5 lbs./1,000 Sq. Ft. 4.0 lbs./1,000 Sq. Ft.

(Editor's Note: Manufacture to supply a listing of spreaders and suggested settings to apply the product at the rates listed above.)

These settings were calibrated and field-tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

WARRANTY

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Manufacturer, and Buyer assumes the risk of any such use.

Additional Labeling Requirements

If the fertilizer guarantee contains iron, the following statement must appear:

This product contains the secondary nutrient iron. Iron may stain concrete and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash off with water.

If the recommended use rate applies less than 1 lb. nitrogen per 1,000 sq. ft. per application, the following statement must appear:

Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

For products distributed in the State of Washington, the following statement must appear:

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-lebsea.com>

CONFRONT* III + FERTILIZER

For the postemergence control of annual and perennial broadleaf weeds in established turf.

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-D†	1.037%
Triethylamine salt of triclopyr.....	0.303%
Methylheptyl ester of fluroxypyr.....	0.207%
OTHER INGREDIENTS	98.453%
TOTAL	100.000%

Acid Equivalents:

2,4-dichlorophenoxyacetic acid – 0.860% 17.20 lb/ton	2, 4-D CAS. No. 94-75-7
triclopyr: 3,5,6-trichloro-2-pyridyloxyacetic acid – 0.217% 4.34 lb/ton	Triclopyr CAS. No. 55335-06-3
fluroxypyr: 4-amino-3,5-dichloro-6-fluoro-2-pyridyloxyacetic acid – 0.143% 2.86 lb/ton	Fluroxypyr CAS. No. 69377-81-7

†Isomer Specific by AOAC Method No. 6.275-6.279 (13th Ed.)

EPA File Symbol 961-XXX

(L) (D)
EPA Est. 961-PA-01: 961-IL-01

*Trademark of Dow AgroSciences LLC.

Superscript used is first letter of lot number.

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for additional Precautionary Statements

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label.

If terms are not acceptable, return the unopened container to the place of purchase at once.

MANUFACTURED BY:
LEBANON SEABOARD CORPORATION
1600 EAST CUMBERLAND STREET
LEBANON, PA 17042
www.lebsea.com

ACCEPTED
with COMMENTS
In EPA Letter Dated:

JAN 25 2005

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No

961-395

GUARANTEED ANALYSIS

Total Nitrogen (N).....	X.X%
X.X% Ammoniacal Nitrogen	
X.X% Water Insoluble Nitrogen	
X.X% Urea Nitrogen	
X.X% Other Water Insoluble Nitrogen	
Available Phosphate (P ₂ O ₅).....	X.X%
Soluble Potash (K ₂ O).....	X.X%

Nutrient Sources: (Guaranteed Analysis and Nutrient Sources to be supplied based on fertilizer analysis used by manufacturer.)

NET WEIGHT: XX.XX LBS. (XX.XX kg)
COVERS: X,XXX Sq. Ft.

9/11

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye irritation. Harmful if swallowed or absorbed through the skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long-sleeved shirt, chemical-resistant gloves, shoes plus socks, and protective eyewear (goggles, face shield or safety glasses) when applying.

User Safety Recommendations

Users should:

- Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Have this product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-888-208-1368 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Drift or runoff from treated areas may be hazardous to aquatic organisms and non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION, OR FOR RESEARCH PURPOSES.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store this product only in its original container in a cool, dry, secured storage area out of the reach of children. In case of spill, carefully sweep up material and use as directed, or dispose of material according to disposal directions below.

DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any interior or outdoor drain.

GENERAL INFORMATION

Confront III — Fertilizer provides control of most annual and perennial broadleaf weeds commonly found in established lawns, including dandelions, clover, chickweed and thistle. Confront III + Fertilizer also contains a long-lasting fertilizer that has all the nutrients your lawn needs for 2-3 months, depending on grass variety.

Confront III — Fertilizer should be applied to wet grass during warm weather when weeds are actively growing. Application under drought conditions may provide less than desirable results.

10/11

USE PRECAUTIONS AND RESTRICTIONS

Restrictions on Retreatment: Do not apply more than 2 broadcast applications per year per treatment site. To minimize the potential for grass injury, additional applications should not be made within 4 weeks of a previous application.

Avoid off target application. Applications should be made so as to avoid getting granules on desirable plants or into shrub, fruit or vegetable gardens or other areas where they may seriously injure susceptible plants.

Do not apply to exposed suckers or roots of shallow rooted trees or shrubs.

Do not apply when windy or if product might run-off into surface water or adjacent land.

Confront III + Fertilizer is recommended for use on the following established turfgrass species:

Established Cool Season Turf Grasses - Bentgrass*, Kentucky bluegrass, Chewing Fescue, Creeping Red Fescue, Sheeps Fescue, Tall Fescue, Perennial Ryegrass

Established Warm Season Turf Grasses* - Bahiagrass, Bermudagrass, Centipedegrass, Zoysiagrass, St. Augustinegrass, Tall Fescue (growing in warm season areas).

*Use Confront III + Fertilizer on these species only if some injury can be tolerated.

For use of Confront III + Fertilizer on a turf species not recommended on this label, the user may determine the suitability for such uses by treating a small area at the recommended rate. Prior to treatment of larger areas, the treated area should be observed for any sign of herbicidal injury during 30 days of normal growing conditions to determine if the treatment is safe to the target turf species. The user assumes the responsibility for any plant damage or other liability resulting from use of Confront III + Fertilizer on turf species not recommended on this label.

APPLICATION INSTRUCTIONS

Confront III + Fertilizer can be applied any time from spring through fall when weeds are actively growing. Broadleaf weed species germinate at different times. Only emerged weeds present at the time of application are controlled. Spring applications will kill the most weeds if you delay applications until the first dandelions flower.

- This product needs to stick to the weeds and should be applied when the grass is wet from dew, rain or watering; poor results will occur if it is applied to dry lawns.
- Apply this product evenly and uniformly over the lawn area, avoiding streaking, skips or overlaps.
- Do not water or disturb the lawn for 24 hours after application.
- Delay mowing for two days after application.
- A second application may be made in 4-6 weeks if needed for hard to control weeds.

Newly seeded turf should be mowed two or three times before being treated. Do not reseed for 3 weeks after application.

SUGGESTED SPREADER SETTINGS

<u>Spreader</u>	<u>3.5 lbs./1,000 Sq. Ft.</u>
-----------------	-------------------------------

(Editor's Note: Manufacture to supply a listing of spreaders and suggested settings to apply the product at 3.5 lbs. per 1,000 Sq. Ft.)

These settings were calibrated and field-tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

WARRANTY

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Manufacturer, and Buyer assumes the risk of any such use.

11/11

Advertising Claims that may be presented on the label

- Eliminates difficult, hard to kill weeds like clover, chickweed, mustard and thistle that other products miss.
- Eliminates dandelions, clover, chickweed, thistle, ground ivy, henbit and other troublesome lawn weeds.
- Combination of three weed control technologies kills more weeds than most other brands.
- Now with Confront III for improved control of clover.
- Unlike most weed & feed products, won't harm St. Augustinegrass and bentgrass.
- Kills poison ivy and poison oak missed by the leading weed & feed brands and leading weed killer sprays.
- A Work Saver Product for the Joy of Weed-Free Lawns.
- Just one easy application kills weeds missed by competitive brands and fertilizes into summer!
- Better weed control and fertilizer than other leading brands.
- Extended time release feeding of nitrogen lasts into summer.
- Effective weed control leads to greener, thicker lawns.
- Apply anytime when weeds are actively growing.
- All the fertilizer nutrients your lawn needs for spring.
- Feeds, greens, strengthens, and thickens your lawn for up to 3 months – longer than leading competitive brands.
- Apply once for the whole season.
- Feeds and strengthens root systems without burning your lawn.
- Fertilize Once for Whole Season – Feeds and greens with slow release fertilizer for up to three months.
- Fertilizes to fill in bare spots and provide plush green lawns.

Additional Labeling Requirements

If the fertilizer guarantee contains iron, the following statement must appear:

This product contains the secondary nutrient iron. Iron may stain concrete and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash off with water.

If the recommended use rate applies less than 1 lb. nitrogen per 1,000 sq. ft. per application, the following statement must appear:

Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

For products distributed in the State of Washington, the following statement must appear:

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-lebsea.com>