	U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460	EPA Reg. Number: 961-406	Date of Issuance: SEP 26 2006
	NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration <small>(under FIFRA, as amended)</small>	Term of Issuance: Conditional	Name of Pesticide Product: Confront IV + Fertilizer
Name and Address of Registrant (include ZIP Code): Lebanon Seaboard Corporation 1600 East Cumberland Street Lebanon, PA 17042			
Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.			
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) and (B) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data. 2. Submit within one year of the date of this registration notice, storage stability (830.6317) and corrosion characteristics (830.6320) studies. 3. Make the following label changes: <ol style="list-style-type: none"> a. Under the Storage and Disposal section for both, commercial and homeowner labels, the first bullet must read "Pesticide Storage". Under the same section for the homeowner use/ residential label, it must read Pesticide Storage and Pesticide Disposal: If empty:" b. The Pesticide Disposal Statement (for the Commercial label) must have the following statement "Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility." c. Also under "Storage and Disposal" replace Container Disposal bullet with "Paper and Plastic Bags:" d. Under the Homeowner Use label title, add "For use on residential lawns". e. Revise the EPA Registration Number to read, EPA Reg. No. 961-406. 			
Signature of Approving Official: Joanne I. Miller Product Manager 23 Herbicide Branch Registration Division (7505P)		Date: SEP 26 2006	

2 7 12

4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

3 7 12

CONFRONT* IV + FERTILIZER

For the postemergence control of annual and perennial broadleaf weeds, and preemergence control of annual grassy weeds and labeled broadleaf weeds, in established ornamental turf, such as lawns, golf courses, roadsides and cemeteries.

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-D [†]	0.819%
Triethylamine salt of triclopyr	0.303%
Methylheptyl ester of fluroxypyr	0.177%
Dithiopyr	0.145%
OTHER INGREDIENTS	98.556%
TOTAL	100.000%

Acid Equivalents:

2,4-dichlorophenoxyacetic acid - 0.680% 13.60 lb/ton	2,4-D CAS. No. 94-75-7
triclopyr: 3,5,6-trichloro-2-pyridyloxyacetic acid - 0.217% 4.34 lb/ton	Triclopyr CAS. No. 55335-06-3
fluroxypyr: 4-amino-3,5-dichloro-6-fluoro-2-pyridyloxyacetic acid - 0.123% 2.46 lb/ton	Fluroxypyr CAS. No. 69377-81-7
dithiopyr: - 0.145% 2.9 lb/ton	Dithiopyr CAS No. 97886-45-8

[†]Isomer Specific by AOAC Method No. 6.275-6.279 (13th Ed.)

EPA File Symbol 961-XXX

*Trademark of Dow AgroSciences LLC.

(L) (D)

EPA Est. 961-PA-01: 961-IL-01

Superscript used is first letter of lot number.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See back panel for additional Precautionary Statements.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label.

If terms are not acceptable, return the unopened container to the place of purchase at once.

In New York State: This product may be applied only by commercial applicators at no more than 241 pounds (0.35 lb. of active ingredient of dithiopyr) per acre (5.5 lb. per 1,000 sq. ft.) per year. Use of this product in Nassau and Suffolk Counties is prohibited.

MANUFACTURED BY:
LEBANON SEABOARD CORPORATION
1600 EAST CUMBERLAND STREET
LEBANON, PA 17042
www.LebanonTurf.com

GUARANTEED ANALYSIS

Total Nitrogen (N)

X.X% Ammoniacal Nitrogen

X.X% Water Insoluble Nitrogen

X.X% Urea Nitrogen

X.X% Other Water Soluble Nitrogen

Available Phosphate (P₂O₅)

Soluble Potash (K₂O)

Nutrient Sources: (Guaranteed Analysis and Nutrient Sources to be supplied based on fertilizer analysis used by manufacturer.)

ACCEPTED
with COMMENTS:
In EPA Letter Dated:
SEP 26 2006

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

961-406

X.X%

X.X%

X.X%

NET WEIGHT: XX LBS. (X.XX kg)

COVERS: X,XXX Sq. Ft.

Page 1 of 10 Pages

CONFRONT IV + FERTILIZER

EPA File Symbol 961-XXX

NEW REGISTRATION LABEL - 3/15/06

4 8 12

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long-sleeved shirt and long pants, socks, shoes, chemical-resistant gloves (such as Natural Rubber, Selection Category A) and protective eyewear (goggles, face shield or safety glasses) when applying. Remove and wash contaminated clothing before reuse.

User Safety Recommendations

- Users should:
- Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
 - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

- IF IN EYES:**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.
- IF ON SKIN**
- Take off contaminated clothing.
- OR CLOTHING:**
- Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.
- SWALLOWED:**
- Call a poison control center or doctor immediately for treatment advice.
 - Have person sip a glass of water if able to swallow.
 - Do not induce vomiting unless told to do so by a poison control center or doctor.
 - Do not give anything by mouth to an unconscious person.
- IF INHALED:**
- Move person to fresh air.
 - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
 - Call a poison control center or doctor for treatment advice.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-888-208-1368 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift or runoff from treated areas may be hazardous to aquatic organisms and non-target plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION OR FOR RESEARCH PURPOSES.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store this product only in its original container in a cool, dry, secured storage area out of the reach of children. Do not store in or around the home. In case of spill, carefully sweep up material and use as directed, or dispose of material according to pesticide disposal directions below.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

5 7 12

Lebanon Confront* IV + Fertilizer provides broad-spectrum postemergence control of actively growing annual and perennial broadleaf weeds, preemergence or early postemergence control of certain annual grassy weeds, and preemergence control or suppression of certain broadleaf weeds in established turf, including residential lawns, golf courses, sports fields, roadsides, cemeteries and around commercial buildings and other commercial and residential turf areas. Apply when broadleaf weeds are actively growing and prior to germination or tillering of target annual grassy weeds. Application under drought conditions may provide less than desirable results. Applications to moist foliage helps granules adhere to weed surface and will improve postemergence control.

Control of Crabgrass Preemergence and Early Postemergence Control

This product provides preemergent control of crabgrass in established turfgrasses (including the large, smooth, and southern species) when applied prior to germination and emergence of crabgrass. In addition, it will control or suppress annual bluegrass, barnyardgrass, crowfootgrass, foxtail (green and yellow), goosegrass, kikuyugrass, ryegrass (annual and perennial) and smutgrass.

This product also provides early postemergent control of crabgrass if applied prior to the 3-leaf stage of growth or first tiller as described below. This stage of growth generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. The practical benefit of early postemer- gent crabgrass control is a 2-4 week longer period of time (window of application) during which crabgrass control can be achieved.

USE PRECAUTIONS AND RESTRICTIONS

Be sure that use of this product conforms to all applicable regulations.

Do not use on grass species such as bentgrass or St. Augustinegrass unless injury can be tolerated. Use of this product on certain varieties of fine fescue may result in undesirable turf injury.

Do not use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Do not use Lebanon Confront* IV + Fertilizer on golf course putting greens or tees.

Avoid off target application. Applications should be made so as to avoid spreading material off target. Do not use this product where desirable clovers are present or apply (spread) the granules onto flowers, vegetables or ornamental shrubbery. Avoid getting granules in shrub bed or other areas where they may seriously injure susceptible plants. Do not apply to exposed suckers or roots of shallow rooted trees or shrubs. Do not exceed specified dosage for any area; be particularly careful within dripline of trees and other ornamental species.

Do not contaminate domestic or irrigation waters. Do not apply on ditches used to transport irrigation water. Do not apply under conditions that may have a potential to run-off to surface water or adjacent land. Do not apply where runoff or irrigation water may flow onto susceptible crops as injury may result.

Where possible, use methods that reduce soil erosion, such as slice seeding; these methods also reduce pesticide run-off. Use of vegetation filter strips is recommended along rivers, creeks, streams, wetlands, etc.

Do not use clippings from treated areas for mulching vegetables or fruit trees.

Maximum Use Rate Restrictions: In New York State: This product may be applied only by commercial applicators at no more than 241 pounds (0.35 lb. of active ingredient of dithiopyr) per acre (5.5 lb. per 1,000 sq. ft.) per year. Use of this product in Nassau and Suffolk Counties is prohibited.

Re-entry Restrictions: Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

Seeding Restrictions: Do not reseed for 10 weeks after application.

Retreatment Restrictions: Do not apply more than 2 broadcast applications per year per treatment site. To minimize the potential for grass injury, additional applications should not be made within 5 weeks of a previous application.

on Confront IV + Fertilizer is recommended for use on the following established turfgrass species:

Established Cool Season Turf Grasses

Common Name	Scientific Name
bentgrass (1)	<i>Agrostis</i> species
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, chewing	<i>Festuca rubra</i> var. <i>commutata</i>
fescue, creeping red	<i>Festuca rubra</i>
fescue, sheeps	<i>Festuca ovina</i>
fescue, tall	<i>Festuca arundinaceae</i>
ryegrass, perennial	<i>Lolium perenne</i>

Established Warm Season Turf Grasses⁽¹⁾

Common Name	Scientific Name
Bahiagrass	<i>Paspalum notatum</i> var. <i>saurae</i> parody
Bermudagrass	<i>Cynodon dactylon</i>
Buffalograss	<i>Buchloe dactyloides</i>
Centipedegrass	<i>Eremochloa ophiuroides</i>
Zoysiagrass	<i>Zoysia japonica</i>
Zoysiagrass	<i>Zoysia tenuifolia</i>
St. Augustinegrass (except *Floritam)	<i>Stenotaphrum secundatum</i>
fescue, tall (growing in warm season areas)	<i>Festuca arundinaceae</i>

(1) Use Lebanon Confront* IV + Fertilizer on this species only if some injury can be tolerated.

WHEN TO APPLY

Established Lawns: Apply when broadleaf weeds are actively growing and prior to germination or tillering of annual grasses. Lebanon Confront* IV + Fertilizer may be applied as a single application, as a split application or as a sequential application in the spring, summer or fall as described in the following use programs. Use of split applications 5-10 weeks apart may provide improved weed control.

Newly Seeded Areas: When establishing a new lawn of Kentucky bluegrass, perennial ryegrass and/or tall fescue, this product may be applied 2 weeks after the first sign of grass germination. For other turfgrass varieties, wait until the grass is well established (3 mowings) before applying this product.

HOW TO APPLY

6 8 12

For optimum broadleaf weed control, apply to recently mowed turf when foliage is moist. When possible, delay mowing or irrigation for several days after treatment. The dithiopyr preemergent component in this product is not effective until activated by rainfall or irrigation. Performance is improved by 1/2 inch of rainfall or sprinkler irrigation within a 30 day period. Erratic weed control may result if not activated by rainfall or irrigation.

Uniform coverage is essential for best results. Before each application, calibrate the spreader according to the equipment manufacturer's directions. Avoid streaking, skips or overlaps. Adjust the spreader so the spreader delivers the appropriate application rate as recommended below.

APPLICATION RATES

Apply 152 to 218 lbs. per acre (3.5 to 5 lbs. per 1,000 sq. ft.) of Lebanon Confront* IV + Fertilizer to control actively growing broadleaf weeds and prevent germination of labeled annual grasses and broadleaf weeds growing in established turf. Broadleaf and grassy weed species germinate at different times. Only emerged weeds present at time of application are controlled.

Specific rate recommendations are based on the user's location, the mowing height of the turfgrass, and whether the application timing is considered to be preemergence or early postemergence. The duration of residual weed control provided by this product is directly related to application rate, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the users' location within a region. The following charts shows application rates and some of the common broadleaf weeds controlled. The rates vary based on region which are described as follows:

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southeastern areas of IL, IN, OH, and coastal areas of CT & RI

South: AL, AR, GA, LA, MS, NC, SC, and TX

Coastal South: HI, FL, and southern coastal areas of AL, GA, LA, MS, NC, SC, and TX

West: AZ, CA, and NV

POSTEMERGENCE & PREEMERGENCE PROGRAMS WITH APPLICATION RATES & WEEDS CONTROLLED

LB/ACRE, 3.5 LB/1,000 SQ. FT.	Fluroxypyr AE/Acre	Triclopyr AE/Acre	2,4-D AE/Acre	Dithiopyr AI/Acre
	0.187	0.330	1.034	0.2204

Postemergent control of the following broadleaf weeds and 3-5 months preemergent control, as well as early postemergence control of crabgrass up to the 3 leaf or first tiller stage, at sites with high cut turf* in all states except NY, parts of states not included in the transition, south, coastal south, or west.

bedstraw	chamise	cornflower	field bindweed	lespedeza	shepherdspurse	velvetleaf
bittercress	chickweeds	creeping beggarweed	geranium, Carolina	little starwort	speedwell	vetch
black medic	chicory	dandelion	goldenrod	mallow	smartweed	woodsorrel
buttonweed, Virginia	cinquefoil	dayflower	henbit	matchweed	sowthistle	yarrow
carpet weed	clovers	deadnettle, purple	hop clover	pineappleweed	spiderwort	
catchweed	cocklebur	docks	knawel	plantain	spotted catsear	
catnip	coffeeweed	dwarf beggarweed	lambquarters	purslane	spurweed	

174 LB/ACRE, 4.0 LB/1,000 SQ. FT.	Fluroxypyr AE/Acre	Triclopyr AE/Acre	2,4-D AE/Acre	Dithiopyr AI/Acre
	0.214	0.378	1.183	0.252

es postemergent control to include the additional weeds listed below and provides 4-6 months of preemergent control on low cut turf** where weed control has been conducted during the previous year. Also provides early postemergent control of crabgrass up to the 3 leaf or first tiller stage on high cut turf*. All states and parts of states not included in the south, coastal south, or west. Recommend split applications of 4.0 + 4.0 Lbs. on high cut turf* in the coastal south and low cut turf in the south.

burdock	Cornspeedwell ⁺	knotweed	oxalis (stricta and comiculata) ⁺	poison ivy	parsley-pier ⁺	wild carrot	veronica species ⁺
buttercup	ground ivy ⁺	mustard		poison oak	prostrate spurge ⁺	wild violet ⁺	(suppression only)

⁺Recommend 2 applications, 5 to 6 weeks apart; apply to moist foliage.

196 LB/ACRE, 4.5 LB/1,000 SQ. FT.	Fluroxypyr AE/Acre	Triclopyr AE/Acre	2,4-D AE/Acre	Dithiopyr AI/Acre
	0.241	0.425	1.333	0.284

Improves postemergent control and provides 3-5 months of preemergent control on high cut turf* and low cut turf** where weed control has been conducted during the previous year. Provides early postemergent control of crabgrass up to the 3 leaf or first tiller stage on high cut turf. Includes the West use higher rate in local areas with longer crabgrass seasons, see below. Note precautions for warm season turf species.

218 LB/ACRE, 5.0 LB/1,000 SQ. FT.	Fluroxypyr AE/Acre	Triclopyr AE/Acre	2,4-D AE/Acre	Dithiopyr AI/Acre
	0.268	0.473	1.482	0.316

Improves postemergent control and provides 4-6 months of preemergent control on low cut turf** where weed control has been conducted during the previous year in the coastal south and west. For coastal south split applications of 5.0 + 5.0 Lbs are recommended. Note precautions for warm season turf species.

*High cut turf such as home lawns and commercial sites.

**Low cut turf such as golf fairways.

7 7 12

SUGGESTED SPREADER SETTINGS

Spreader 3.5 lbs./1,000 Sq. Ft. 4 lbs./1,000 Sq. Ft. 4.5 lbs./1,000 Sq. Ft. 5 lbs./1,000 Sq. Ft.

(Editor's Note: Manufacturer to supply a listing of spreaders and suggested settings to apply this product at a rate of 3.5, 4, 4.5 and 5 lbs. per 1,000 sq. ft.)

These settings were calibrated and field-tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

WARRANTY

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Manufacturer, and Buyer assumes the risk of any such use.

Additional Labeling Requirements:

If the fertilizer guarantee contains iron, the following statement must appear:

This product contains the secondary nutrient iron. Iron may stain concrete and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash off with water.

If the recommended use rate applies less than 1 pound nitrogen per 1,000 sq. ft. per application, the following statement must appear:

Additional applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

For products distributed in the State of Washington, the following statement must appear:

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-lebsea.com>

CONFRONT* IV + FERTILIZER

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-D [†]	0.819%
Triethylamine salt of triclopyr	0.303%
Methylheptyl ester of fluroxypyr	0.177%
Dithiopyr	0.145%
OTHER INGREDIENTS	98.556%
TOTAL	100.000%

Acid Equivalents:

2,4-dichlorophenoxyacetic acid - 0.680% 13.60 lb/ton	2,4-D CAS. No. 94-75-7
triclopyr: 3,5,6-trichloro-2-pyridyloxyacetic acid - 0.217% 4.34 lb/ton	Triclopyr CAS. No. 55335-06-3
fluroxypyr: 4-amino-3,5-dichloro-6-fluoro-2-pyridyloxyacetic acid - 0.123% 2.46 lb/ton	Fluroxypyr CAS. No. 69377-81-7
dithiopyr: - 0.145% 2.9 lb/ton	Dithiopyr CAS No. 97886-45-8

[†]Isomer Specific by AOAC Method No. 6.275-6.279 (13th Ed.)

EPA File Symbol 961-XXX

*Trademark of Dow AgroSciences LLC.

(L) (D)

EPA Est. 961-PA-01: 961-IL-01

Superscript used is first letter of lot number.

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for additional Precautionary Statements.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label.

If terms are not acceptable, return the unopened container to the place of purchase at once.

In New York State: Do not exceed 3.5 lbs. of product per 1,000 sq. ft. (0.22 lb. active ingredient of dithiopyr per acre) per year. Do not apply more than once per year to the same area.

MANUFACTURED BY:
LEBANON SEABOARD CORPORATION
1600 EAST CUMBERLAND STREET
LEBANON, PA 17042
www.lebsea.com

ACCEPTED
with COMMENTS
In EPA Letter Dated:
SEP 26 2006

GUARANTEED ANALYSIS

Total Nitrogen (N)	X.X%
X.X% Ammoniacal Nitrogen	
X.X% Water Insoluble Nitrogen	
X.X% Urea Nitrogen	
X.X% Other Water Soluble Nitrogen	
Available Phosphate (P ₂ O ₅)	X.X%
Soluble Potash (K ₂ O)	X.X%

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

961-406

Nutrient Sources: (Guaranteed Analysis and Nutrient Sources to be supplied based on fertilizer analysis used by manufacturer.)

NET WEIGHT: XX LBS. (X.XX kg)
COVERS: X,XXX Sq. Ft.

9 8 12

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long-sleeved shirt and long pants, socks, shoes, chemical-resistant gloves (such as or made out of any waterproof material) and protective eyewear (goggles, face shield or safety glasses) when applying. Remove and wash contaminated clothing before reuse.

User Safety Recommendations

- Users should:
- Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
 - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

- IF IN EYES:**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.
- IF ON SKIN**
- Take off contaminated clothing.
- OR CLOTHING:**
- Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.
- IF SWALLOWED:**
- Call a poison control center or doctor immediately for treatment advice.
 - Have person sip a glass of water if able to swallow.
 - Do not induce vomiting unless told to do so by a poison control center or doctor.
 - Do not give anything by mouth to an unconscious person.
- IF INHALED:**
- Move person to fresh air.
 - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
 - Call a poison control center or doctor for treatment advice.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-888-208-1368 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift or runoff from treated areas may be hazardous to aquatic organisms and non-target plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION OR FOR RESEARCH PURPOSES.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store this product only in its original container in a cool, dry, secured storage area out of the reach of children. In case of spill, carefully sweep up material and use as directed, or dispose of material according to disposal directions below.

DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

GENERAL INFORMATION

10 8 12

Lebanon Confront* IV + Fertilizer will control all your lawn's weeds in one application. It will kill over 50 lawn weeds commonly found in home lawns including clover, dandelion, plantain and chickweed and will prevent crabgrass and other undesirable annual grasses. Lebanon Confront* IV + Fertilizer also contains the slow release nitrogen your lawn needs for 2-3 months, depending on grass variety and weather conditions.

Lebanon Confront IV + Fertilizer is recommended for use on the following established turfgrass species:

Established Cool Season Turf Grasses — Bentgrass*, Kentucky Bluegrass, Fine and Tall Fescues, Perennial Ryegrass

Established Warm Season Turf Grasses* — Bahiagrass, Bermudagrass, Buffalograss, Centipedegrass, Zoysiagrass, St. Augustinegrass (except 'Floritam'), Tall Fescue (growing in warm season areas).

*Use on these grass species only if some injury can be tolerated.

If the lawn has been seeded recently, wait until after the third mowing before using this product. Do not seed, overseed or sprig the treated area for 10 weeks after application.

USE PRECAUTIONS AND RESTRICTIONS

Retreatment Restrictions: Do not apply more than 2 broadcast applications per year per treatment site. To minimize the potential for grass injury, additional applications should not be made within 6 weeks of a previous application.

Avoid off target application. Applications should be made so as to avoid getting granules on desirable plants or into shrub, fruit or vegetable gardens or other areas where they may seriously injure susceptible plants.

Do not apply to exposed suckers or roots of shallow rooted trees or shrubs.

Do not apply when windy or if product might run-off into surface water or adjacent land.

Do not use clippings from treated areas for mulching vegetables, vines, or fruit trees.

Keep people and pets off treated areas until dust has settled.

WEEDS CONTROLLED AND/OR SUPPRESSED

Annual Grass Weeds:

Annual Bluegrass	Barnyardgrass	Crowfootgrass	Goosegrass	Perennial Ryegrass
Annual Ryegrass	Crabgrass	Foxtail (green and yellow)	Kikuyugrass	Smutgrass

Established Broadleaf Weeds:

Bedstraw	Chamise	Cornflower	Field Bindweed	Lespedeza	Plantain	Spotted Catsear
Bittercress	Chickweeds	Creeping Beggarweed	Geranium, Carolina	Little Starwort	Purslane	Spurge
Black Medic	Chicory	Dandelion	Goldenrod	Mallow	Shepherdspurse	Spurweed
Buttonweed, Virginia	Cinquefoil	Dayflower	Henbit	Matchweed	Speedwell	Velvetleaf
Carpet Weed	Clovers	Deadnettle, purple	Hop Clover	Oxalis	Smartweed	Vetch
Catchweed	Cocklebur	Docks	Knawel	Parsley-piert	Sowthistle	Woodsorrel
Catnip	Coffeeweed	Dwarf Beggarweed	Lambsquarters	Pineappleweed	Spiderwort	Yarrow

WHEN TO APPLY

Lebanon Confront* IV + Fertilizer can be used in spring and fall. It should be used when the grass and weeds are actively growing and the soil is moist. Do not apply under drought conditions.

In the spring, the ideal time to apply Lebanon Confront* IV + Fertilizer is after the forsythia has bloomed or when the first dandelion flowers appear. In southern areas (see map), make a second application in 6-8 weeks for best weed and crabgrass control and for full season fertilization.

In the fall, apply Lebanon Confront* IV + Fertilizer when the weather cools and both the grass and weeds are actively growing, generally around Labor Day.


HOW TO APPLY

Lebanon Confront* IV + Fertilizer needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain; poor results will occur if it is applied to dry lawns. If possible, do not water or mow the lawn for 3 days after application. After 3 days and any time up to the 30th day after treatment, water the top 1-2 inches of soil to fully activate the product.

Apply this product when the wind is calm. Do not apply this product onto desirable plants or shrubs as damage may result. If the product gets onto other plants, immediately brush or wash it off the foliage.

Apply 3.5 lbs. of Lebanon Confront* IV + Fertilizer per 1,000 sq. ft. of turfgrass. Use a drop or rotary spreader designed to apply granular fertilizers. Apply evenly and uniformly over the lawn area for best results. Avoid streaking, skips or overlaps as poor weed control or turfgrass injury may occur.

USE REGIONS WITHIN THE UNITED STATES



SUGGESTED SPREADER SETTINGS

11 8 12

(Editor's Note: Manufacturer to supply a listing of spreaders and suggested settings to apply this product at a rate of 3.5 lbs. per 1,000 sq. ft.)


These settings are approximate. Rate of walk, moisture and spreader condition can vary the spreader setting necessary to apply the correct amount. Before each application, calibrate your spreader according to manufacturer's directions. Adjust to apply at the rate of 1 lb. of product per 286 sq. ft. Check equipment frequently to verify calibration.

Broadcast Spreader

Settings

Optional Spreading Instructions:

[Apply in a single pass at a rate of 3.5 lbs. per 1,000 sq. ft. Start on the outside border and apply in a circular path working toward the center. Overlap the coverage slightly, and shut off spreader before making hard turns.]




Graphic Optional

Drop Spreader

Settings

Optional Spreading Instructions:

[Apply in a single pass at a rate of 3.5 lbs. per 1,000 sq. ft. Start on the outside border and apply in a rectangular path working back and forth across the length of the lawn. Overlap the coverage slightly, and shut off spreader before making hard turns.]



Graphic Optional

If a spreader model is not listed (or does not appear above), go to our website at www.lebsea.com (or www.preen.com), or call toll-free (800) 233-1067, 8 A.M. - 5 P.M. EDT, Monday - Friday, for additional model settings.

WARRANTY

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Manufacturer, and Buyer assumes the risk of any such use.

Additional Labeling Requirements:

If the fertilizer guarantee contains iron, the following statement must appear:

This product contains the secondary nutrient iron. Iron may stain concrete and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash off with water.

If the recommended use rate applies less than 1 pound nitrogen per 1,000 sq. ft. per application, the following statement must appear:

Additional applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

For products distributed in the State of Washington, the following statement must appear:

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-lebsea.com>

Advertising claims that may be presented on the label

For the postemergence control of annual and perennial broadleaf weeds, and preemergence control of annual grassy weeds and labeled broadleaf weeds, in established ornamental turf.

- Eliminates difficult, hard to kill weeds like clover, dandelion, chickweed and plantain that other products miss
- Eliminates dandelions, clover, chickweed, goldenrod, henbit and other troublesome lawn weeds
- Combination of four weed control technologies kills more weeds than most other brands
- Season long weed preventer
- Now with Confront IV for improved control of broadleaf and grassy weeds
- Unlike most weed & feed products, won't harm many warm season turf grasses
- One product kills broadleaf weeds and prevents crabgrass
- Apply once in spring when weeds are growing
- Stops more weeds in one application
- Works season long

Advertising claims that may be presented on the label (continued)

12 8 12

- Controls tough grasses and weeds [such as clover, crabgrass, goldenrod, henbit, etc.]
- Apply at any time [spring, summer, fall]
- Use for a [beautiful] weed free lawn
- *Easy to use*
- Just one application kills weeds missed by competitive brands and fertilizes into summer!
- Better weed control and fertilizer than other leading brands
- Extended, time release feeding of nitrogen lasts into summer
- Effective weed control leads to greener, thicker lawns
- Apply anytime when weeds are actively growing
- All the fertilizer nutrients your lawn needs for spring
- Feeds, greens, strengthens, and thickens your lawn for up to 3 months – longer than leading competitive brands
- Apply once for the whole season
- Feeds and strengthens root systems without burning your lawn
- Fertilize Once for Whole Season - Feeds and greens with slow release fertilizer for up to three months
- Fertilizes to fill in bare spots and provide plush green lawns
- Prevents weeds & fertilizes the lawn
- Fertilizes to provide rich color and vigorous growth
- Effective weed control - prevents weeds all season long
- New and improved
- Season long weed preventer and fertilizer
 - Controls weeds longer
 - Fertilizes longer

PREEN® GUARANTEE

Lebanon Seaboard Corporation guarantees you will be satisfied with the performance of this product when used as directed, or your money back. If not satisfied, describe the nature of your problem, and send with proof of purchase to: Lebanon Seaboard Corporation, 1600 East Cumberland Street, Lebanon, PA 17042.

Preen® is a registered trademark of Lebanon Seaboard Corporation.