

961-406

11-9-2007

1/13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Mr. Henry E. Rohland
Product Registration
Lebanon Seaboard Corporation
1600 East Cumberland Street
Lebanon, PA 17042

NOV 9 2007

SUBJECT: Application for Pesticide Notification (PRN 98-10)
Request Storage & Disposal Changes and Marketing Claims
EPA Reg. No. 961-406
Application Dated October 9, 2007

Dear Mr. Rohland:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 10/09/07 the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

313



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 961-406	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Confront IV + Fertilizer	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Lebanon Seaboard Corporation 1600 East Cumberland Street Lebanon, PA 17042 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Response to PR Notice 2007-1 and 2 Marketing statements. Deletions in RED Additions in BLUE
 This notification is consistent with the guidance in PR Notice 2007-1 and the requirements of EPA's regulations at 40 CFR 156.10 and 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the guidance of PR Notice 2007-1 and the requirements of 40 CFR 156.10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA."

Section - III

1. Material This Product Will Be Packaged In:					
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name Henry E. Rohland		Title EPA Registration Manager		Telephone No. (Include Area Code) _____	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.					6. Date Application Received (Stamped)
2. Signature 		3. Title EPA Registration Manager			
4. Typed Name Henry E. Rohland		5. Date 9 Oct 2007			

CONFRONT* IV + FERTILIZER

For the postemergence control of annual and perennial broadleaf weeds,
and preemergence control of annual grassy weeds and labeled broadleaf weeds,
in established ornamental turf, such as lawns, golf courses, roadsides and cemeteries.

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-D [†]	0.819%
Triethylamine salt of triclopyr	0.303%
Methylheptyl ester of fluroxypyr	0.177%
Dithiopyr	0.145%
OTHER INGREDIENTS	98.556%
TOTAL	100.000%

Acid Equivalents:

2,4-dichlorophenoxyacetic acid - 0.680% 13.60 lb/ton	2,4-D CAS. No. 94-75-7
triclopyr: 3,5,6-trichloro-2-pyridyloxyacetic acid - 0.217% 4.34 lb/ton	Triclopyr CAS. No. 55335-06-3
fluroxypyr: 4-amino-3,5-dichloro-6-fluoro-2-pyridyloxyacetic acid - 0.123% 2.46 lb/ton	Fluroxypyr CAS. No. 69377-81-7
dithiopyr: - 0.145% 2.9 lb/ton	Dithiopyr CAS No. 97886-45-8

[†]Isomer Specific by AOAC Method No. 6.275-6.279 (13th Ed.)

EPA File Symbol 961-406

*Trademark of Dow AgroSciences LLC.

(L) (D)

EPA Est. 961-PA-01: 961-IL-01

Superscript used is first letter of lot number.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See back panel for additional Precautionary Statements.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label.

If terms are not acceptable, return the unopened container to the place of purchase at once.

In New York State: This product may be applied only by commercial applicators at no more than 241 pounds (0.35 lb. of active ingredient of dithiopyr) per acre (5.5 lb. per 1,000 sq. ft.) per year. Use of this product in Nassau and Suffolk Counties is prohibited.

MANUFACTURED BY:
LEBANON SEABOARD CORPORATION
1600 EAST CUMBERLAND STREET
LEBANON, PA 17042
www.LebanonTurf.com

GUARANTEED ANALYSIS

Total Nitrogen (N)	X.X%
X.X% Ammoniacal Nitrogen	
X.X% Water Insoluble Nitrogen	
X.X% Urea Nitrogen	
X.X% Other Water Soluble Nitrogen	
Available Phosphate (P ₂ O ₅)	X.X%
Soluble Potash (K ₂ O)	X.X%
Nutrient Sources: (Guaranteed Analysis and Nutrient Sources to be supplied based on fertilizer analysis used by manufacturer.)	

NET WEIGHT: XX LBS. (X.XX kg)

COVERS: X,XXX Sq. Ft.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long-sleeved shirt and long pants, socks, shoes, chemical-resistant gloves (such as Natural Rubber, Selection Category A) and protective eyewear (goggles, face shield or safety glasses) when applying. Remove and wash contaminated clothing before reuse.

User Safety Recommendations

- Users should:
- Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
 - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

- IF IN EYES:**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.
- IF ON SKIN OR CLOTHING:**
- Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.
- IF SWALLOWED:**
- Call a poison control center or doctor immediately for treatment advice.
 - Have person sip a glass of water if able to swallow.
 - Do not induce vomiting unless told to do so by a poison control center or doctor.
 - Do not give anything by mouth to an unconscious person.
- IF INHALED:**
- Move person to fresh air.
 - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
 - Call a poison control center or doctor for treatment advice.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-888-208-1368 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift or runoff from treated areas may be hazardous to aquatic organisms and non-target plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION OR FOR RESEARCH PURPOSES.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store this product only in its original container in a cool, dry, locked place out of the reach of children and animals. ~~secured storage area out of the reach of children. Do not store in or around the home. In case of spill, carefully sweep up material and use as directed, or dispose of material according to pesticide disposal directions below.~~

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. ~~Improper disposal of excess pesticide or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.~~

PAPER AND PLASTIC BAG CONTAINER DISPOSAL: Completely empty bag container into application equipment. Then dispose of empty bag container in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

4/13

Lebanon Confront* IV + Fertilizer provides broad-spectrum postemergence control of actively growing annual and perennial broadleaf weeds, preemergence or early postemergence control of certain annual grassy weeds, and preemergence control or suppression of certain broadleaf weeds in established turf, including residential lawns, golf courses, sports fields, roadsides, cemeteries and around commercial buildings and other commercial and residential turf areas. Apply when broadleaf weeds are actively growing and prior to germination or tillering of target annual grassy weeds. Application under drought conditions may provide less than desirable results. Applications to moist foliage helps granules adhere to weed surface and will improve postemergence control.

Control of Crabgrass Preemergence and Early Postemergence Control

This product provides preemergent control of crabgrass in established turfgrasses (including the large, smooth, and southern species) when applied prior to germination and emergence of crabgrass. In addition, it will control or suppress annual bluegrass, barnyardgrass, crowfootgrass, foxtail (green and yellow), goosegrass, kikuyugrass, ryegrass (annual and perennial) and smutgrass.

This product also provides early postemergent control of crabgrass if applied prior to the 3-leaf stage of growth or first tiller as described below. This stage of growth generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. The practical benefit of early postemergent crabgrass control is a 2-4 week longer period of time (window of application) during which crabgrass control can be achieved.

USE PRECAUTIONS AND RESTRICTIONS

Be sure that use of this product conforms to all applicable regulations.

Do not use on grass species such as bentgrass or St. Augustinegrass unless injury can be tolerated. Use of this product on certain varieties of fine fescue may result in undesirable turf injury.

Do not use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Do not use Lebanon Confront* IV + Fertilizer on golf course putting greens or tees.

Avoid off target application. Applications should be made so as to avoid spreading material off target. Do not use this product where desirable clovers are present or apply (spread) the granules onto flowers, vegetables or ornamental shrubbery. Avoid getting granules in shrub bed or other areas where they may seriously injure susceptible plants. Do not apply to exposed suckers or roots of shallow rooted trees or shrubs. Do not exceed specified dosage for any area; be particularly careful within dripline of trees and other ornamental species.

Do not contaminate domestic or irrigation waters. Do not apply on ditches used to transport irrigation water. Do not apply under conditions that may have a potential to run-off to surface water or adjacent land. Do not apply where runoff or irrigation water may flow onto susceptible crops as injury may result. Where possible, use methods that reduce soil erosion, such as slice seeding; these methods also reduce pesticide run-off. Use of vegetation filter strips is recommended along rivers, creeks, streams, wetlands, etc.

Do not use clippings from treated areas for mulching vegetables or fruit trees.

Maximum Use Rate Restrictions: In New York State: This product may be applied only by commercial applicators at no more than 241 pounds (0.35 lb. of active ingredient of dithiopyr) per acre (5.5 lb. per 1,000 sq. ft.) per year. Use of this product in Nassau and Suffolk Counties is prohibited.

Re-entry Restrictions: Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

Seedling Restrictions: Do not reseed for 10 weeks after application.

Retreatment Restrictions: Do not apply more than 2 broadcast applications per year per treatment site. To minimize the potential for grass injury, additional applications should not be made within 5 weeks of a previous application.

Lebanon Confront IV + Fertilizer is recommended for use on the following established turfgrass species:

Established Cool Season Turf Grasses

Common Name	Scientific Name
bentgrass (1)	<i>Agrostis species</i>
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, chewing	<i>Festuca rubra var. commutata</i>
fescue, creeping red	<i>Festuca rubra</i>
fescue, sheeps	<i>Festuca ovina</i>
fescue, tall	<i>Festuca arundinaceae</i>
ryegrass, perennial	<i>Lolium perenne</i>

Established Warm Season Turf Grasses(1)

Common Name	Scientific Name
Bahiagrass	<i>Paspalum notatum var. sauræ parody</i>
Bermudagrass	<i>Cynodon dactylon</i>
Buffalograss	<i>Buchloe dactyloides</i>
Centipedegrass	<i>Eremochloa ophiuroides</i>
Zoysiagrass	<i>Zoysia japonica</i>
Zoysiagrass	<i>Zoysia tenuifolia</i>
St. Augustinegrass (except "Florata")	<i>Stenotaphrum secundatum</i>
fescue, tall (growing in warm season areas)	<i>Festuca arundinaceae</i>

(1) Use Lebanon Confront* IV + Fertilizer on this species only if some injury can be tolerated.

WHEN TO APPLY

Established Lawns: Apply when broadleaf weeds are actively growing and prior to germination or tillering of annual grasses. Lebanon Confront* IV + Fertilizer may be applied as a single application, as a split application or as a sequential application in the spring, summer or fall as described in the following use programs. Use of split applications 5-10 weeks apart may provide improved weed control.

Newly Seeded Areas: When establishing a new lawn of Kentucky bluegrass, perennial ryegrass and/or tall fescue, this product may be applied 2 weeks after the first sign of grass germination. For other turfgrass varieties, wait until the grass is well established (3 mowings) before applying this product.

HOW TO APPLY

7/13

For optimum broadleaf weed control, apply to recently mowed turf when foliage is moist. When possible, delay mowing or irrigation for several days after treatment. The dithiopyr preemergent component in this product is not effective until activated by rainfall or irrigation. Performance is improved by 1/2 inch of rainfall or sprinkler irrigation within a 30 day period. Erratic weed control may result if not activated by rainfall or irrigation.

Uniform coverage is essential for best results. Before each application, calibrate the spreader according to the equipment manufacturer's directions. Avoid streaking, skips or overlaps. Adjust the spreader so the spreader delivers the appropriate application rate as recommended below.

APPLICATION RATES

Apply 152 to 218 lbs. per acre (3.5 to 5 lbs. per 1,000 sq. ft.) of Lebanon Confront[®] IV + Fertilizer to control actively growing broadleaf weeds and prevent germination of labeled annual grasses and broadleaf weeds growing in established turf. Broadleaf and grassy weed species germinate at different times. Only emerged weeds present at time of application are controlled.

Specific rate recommendations are based on the user's location, the mowing height of the turfgrass, and whether the application timing is considered to be preemergence or early postemergence. The duration of residual weed control provided by this product is directly related to application rate, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the users' location within a region. The following charts shows application rates and some of the common broadleaf weeds controlled. The rates vary based on region which are described as follows:

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southeastern areas of IL, IN, OH, and coastal areas of CT & RI

South: AL, AR, GA, LA, MS, NC, SC, and TX

Coastal South: HI, FL, and southern coastal areas of AL, GA, LA, MS, NC, SC, and TX

West: AZ, CA, and NV

POSTEMERGENCE & PREEMERGENCE PROGRAMS WITH APPLICATION RATES & WEEDS CONTROLLED

152 LB/ACRE, 3.5 LB/1,000 SQ. FT.	Fluroxypyr AE/Acre	Triclopyr AE/Acre	2,4-D AE/Acre	Dithiopyr AI/Acre
	0.187	0.330	1.034	0.2204

Postemergent control of the following broadleaf weeds and 3-5 months preemergent control, as well as early postemergence control of crabgrass up to the 3 leaf or first tiller stage, at sites with high cut turf* in all states except NY, parts of states not included in the transition, south, coastal south, or west.

bedstraw	chamise	cornflower	field bindweed	lespedeza	shepherdspurse	velvetleaf
bittercress	chickweeds	creeping beggarweed	geranium, Carolina	little starwort	speedwell	vetch
black medic	chicory	dandelion	goldenrod	mallow	smartweed	woodsorrel
buttonweed, Virginia	cinquefoil	dayflower	henbit	matchweed	sowthistle	yarrow
carpet weed	clovers	deadnettle, purple	hop clover	pineappleweed	spiderwort	
catchweed	cocklebur	docks	knawel	plantain	spotted catsear	
catnip	coffeeweed	dwarf beggarweed	lambquarters	purslane	spurweed	

174 LB/ACRE, 4.0 LB/1,000 SQ. FT.	Fluroxypyr AE/Acre	Triclopyr AE/Acre	2,4-D AE/Acre	Dithiopyr AI/Acre
	0.214	0.378	1.183	0.252

Increases postemergent control to include the additional weeds listed below and provides 4-6 months of preemergent control on low cut turf** where weed control has been conducted during the previous year. Also provides early postemergent control of crabgrass up to the 3 leaf or first tiller stage on high cut turf*. All states and parts of states not included in the south, coastal south, or west. Recommend split applications of 4.0 + 4.0 Lbs. on high cut turf* in the coastal south and low cut turf in the south.

burdock	Cornspeedwell ⁺	knotweed	oxalis (stricta and comiculata) ⁺	poison ivy	parsley-piert ⁺	wild carrot	veronica species ⁺
buttercup	ground ivy ⁺	mustard		poison oak	prostrate spurge ⁺	wild violet ⁺	(suppression only)

*Recommend 2 applications, 5 to 6 weeks apart; apply to moist foliage.

196 LB/ACRE, 4.5 LB/1,000 SQ. FT.	Fluroxypyr AE/Acre	Triclopyr AE/Acre	2,4-D AE/Acre	Dithiopyr AI/Acre
	0.241	0.425	1.333	0.284

Improves postemergent control and provides 3-5 months of preemergent control on high cut turf* and low cut turf** where weed control has been conducted during the previous year. Provides early postemergent control of crabgrass up to the 3 leaf or first tiller stage on high cut turf. Includes the West use higher rate in local areas with longer crabgrass seasons, see below. Note precautions for warm season turf species.

218 LB/ACRE, 5.0 LB/1,000 SQ. FT.	Fluroxypyr AE/Acre	Triclopyr AE/Acre	2,4-D AE/Acre	Dithiopyr AI/Acre
	0.268	0.473	1.482	0.316

Improves postemergent control and provides 4-6 months of preemergent control on low cut turf** where weed control has been conducted during the previous year in the coastal south and west. For coastal south split applications of 5.0 + 5.0 Lbs are recommended. Note precautions for warm season turf species.

*High cut turf such as home lawns and commercial sites.

**Low cut turf such as golf fairways.

9/13

SUGGESTED SPREADER SETTINGS

Spreader 3.5 lbs./1,000 Sq. Ft. 4 lbs./1,000 Sq. Ft. 4.5 lbs./1,000 Sq. Ft. 5 lbs./1,000 Sq. Ft.

(Editor's Note: Manufacturer to supply a listing of spreaders and suggested settings to apply this product at a rate of 3.5, 4, 4.5 and 5 lbs. per 1,000 sq. ft.)

These settings were calibrated and field-tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

WARRANTY

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Manufacturer, and Buyer assumes the risk of any such use.

Additional Labeling Requirements:

If the fertilizer guarantee contains iron, the following statement must appear:

This product contains the secondary nutrient iron. Iron may stain concrete and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash off with water.

If the recommended use rate applies less than 1 pound nitrogen per 1,000 sq. ft. per application, the following statement must appear:

Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

For products distributed in the State of Washington, the following statement must appear:

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-lebsea.com>

Advertising claims that may be stated on the label:

PRACTICAL SOLUTIONS FOR THE SERIOUS TURF PROFESSIONALS

PRACTICAL SOLUTIONS FOR THE TURF PROFESSIONALS

9/13

CONFRONT* IV + FERTILIZER

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-D [†]	0.819%
Triethylamine salt of triclopyr	0.303%
Methylheptyl ester of fluroxypyr	0.177%
Dithiopyr	0.145%
OTHER INGREDIENTS	98.556%
TOTAL	100.000%

Acid Equivalents:

2,4-dichlorophenoxyacetic acid – 0.680% 13.60 lb/ton	2,4-D CAS. No. 94-75-7
triclopyr: 3,5,6-trichloro-2-pyridyloxyacetic acid – 0.217% 4.34 lb/ton	Triclopyr CAS. No. 55335-06-3
fluroxypyr: 4-amino-3,5-dichloro-6-fluoro-2-pyridyloxyacetic acid – 0.123% 2.46 lb/ton	Fluroxypyr CAS. No. 69377-81-7
dithiopyr: — 0.145% 2.9 lb/ton	Dithiopyr CAS No. 97886-45-8

[†]Isomer Specific by AOAC Method No. 6.275-6.279 (13th Ed.)

(L) (D)

EPA File Symbol 961-406

EPA Est. 961-PA-01: 961-IL-01

*Trademark of Dow AgroSciences LLC.

Superscript used is first letter of lot number.

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for additional Precautionary Statements.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label.

If terms are not acceptable, return the unopened container to the place of purchase at once.

In New York State: Do not exceed 3.5 lbs. of product per 1,000 sq. ft. (0.22 lb. active ingredient of dithiopyr per acre) per year. Do not apply more than once per year to the same area.

MANUFACTURED BY:
LEBANON SEABOARD CORPORATION
1600 EAST CUMBERLAND STREET
LEBANON, PA 17042
www.lebsea.com

GUARANTEED ANALYSIS

Total Nitrogen (N)	X.X%
X.X% Ammoniacal Nitrogen	
X.X% Water Insoluble Nitrogen	
X.X% Urea Nitrogen	
X.X% Other Water Soluble Nitrogen	

Available Phosphate (P ₂ O ₅)	X.X%
Soluble Potash (K ₂ O)	X.X%

Nutrient Sources: (Guaranteed Analysis and Nutrient Sources to be supplied based on fertilizer analysis used by manufacturer.)

NET WEIGHT: XX LBS. (X.XX kg)
COVERS: X,XXX Sq. Ft.

10/13

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long-sleeved shirt and long pants, socks, shoes, chemical-resistant gloves (such as or made out of any waterproof material) and protective eyewear (goggles, face shield or safety glasses) when applying. Remove and wash contaminated clothing before reuse.

User Safety Recommendations

- Users should:
- Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
 - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

- IF IN EYES:**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.
- IF ON SKIN OR CLOTHING:**
- Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.
- IF SWALLOWED:**
- Call a poison control center or doctor immediately for treatment advice.
 - Have person sip a glass of water if able to swallow.
 - Do not induce vomiting unless told to do so by a poison control center or doctor.
 - Do not give anything by mouth to an unconscious person.
- IF INHALED:**
- Move person to fresh air.
 - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
 - Call a poison control center or doctor for treatment advice.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-888-208-1368 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift or runoff from treated areas may be hazardous to aquatic organisms and non-target plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION OR FOR RESEARCH PURPOSES.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store this product only in its original container in a cool, dry, secured storage area out of the reach of children. In case of spill, carefully sweep up material and use as directed, or dispose of material according to disposal directions below. locked place out of the reach of children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

PESTICIDE-CONTAINER DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused material down any indoor or outdoor drain. Use as directed or call your local solid waste agency for disposal instructions.

GENERAL INFORMATION

Lebanon Confront* IV + Fertilizer will control all your lawn's weeds in one application. It will kill over 50 lawn weeds commonly found in home lawns including clover, dandelion, plantain and chickweed and will prevent crabgrass and other undesirable annual grasses. Lebanon Confront* IV + Fertilizer also contains the slow release nitrogen your lawn needs for 2-3 months, depending on grass variety and weather conditions.

Lebanon Confront IV + Fertilizer is recommended for use on the following established turfgrass species:

Established Cool Season Turf Grasses — Bentgrass*, Kentucky Bluegrass, Fine and Tall Fescues, Perennial Ryegrass

Established Warm Season Turf Grasses* — Bahiagrass, Bermudagrass, Buffalograss, Centipedegrass, Zoysiagrass, St. Augustinegrass (except 'Florata'), Tall Fescue (growing in warm season areas).

*Use on these grass species only if some injury can be tolerated.

If the lawn has been seeded recently, wait until after the third mowing before using this product. Do not seed, overseed or sprig the treated area for 10 weeks after application.

USE PRECAUTIONS AND RESTRICTIONS

Retreatment Restrictions: Do not apply more than 2 broadcast applications per year per treatment site. To minimize the potential for grass injury, additional applications should not be made within 6 weeks of a previous application.

Avoid off target application. Applications should be made so as to avoid getting granules on desirable plants or into shrub, fruit or vegetable gardens or other areas where they may seriously injure susceptible plants.

Do not apply to exposed suckers or roots of shallow rooted trees or shrubs.

Do not apply when windy or if product might run-off into surface water or adjacent land.

Do not use clippings from treated areas for mulching vegetables, vines, or fruit trees.

Keep people and pets off treated areas until dust has settled.

WEEDS CONTROLLED AND/OR SUPPRESSED

Annual Grass Weeds:

Annual Bluegrass
Annual Ryegrass

Barnyardgrass
Crabgrass

Crowfootgrass
Foxtail (green and yellow)

Goosegrass
Kikuyugrass

Perennial Ryegrass
Smutgrass

Established Broadleaf Weeds:

Bedstraw
Bittercress
Black Medic
Buttonweed, Virginia
Carpet Weed
Catchweed
Catnip
Chamise
Chickweeds
Chicory
Cinquefoil
Clovers
Cocklebur
Coffeeweed

Cornflower
Creeping Beggarweed
Dandelion
Dayflower
Deadnettle, purple
Docks
Dwarf Beggarweed

Field Bindweed
Geranium, Carolina
Goldenrod
Henbit
Hop Clover
Knawel
Lambsquarters

Lespedeza
Little Starwort
Mallow
Matchweed
Oxalis
Parsley-piert
Pineappleweed

Plantain
Purslane
Shepherdspurge
Speedwell
Smartweed
Sowthistle
Spiderwort

Spotted Catsear
Spurge
Spurweed
Velvetleaf
Vetch
Woodsorrel
Yarrow

WHEN TO APPLY

Lebanon Confront* IV + Fertilizer can be used in spring and fall. It should be used when the grass and weeds are actively growing and the soil is moist. Do not apply under drought conditions.

In the spring, the ideal time to apply Lebanon Confront* IV + Fertilizer is after the forsythia has bloomed or when the first dandelion flowers appear. In southern areas (see map), make a second application in 6-8 weeks for best weed and crabgrass control and for full season fertilization.

In the fall, apply Lebanon Confront* IV + Fertilizer when the weather cools and both the grass and weeds are actively growing, generally around Labor Day.


HOW TO APPLY

Lebanon Confront* IV + Fertilizer needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain; poor results will occur if it is applied to dry lawns. If possible, do not water or mow the lawn for 3 days after application. After 3 days and any time up to the 30th day after treatment, water the top 1-2 inches of soil to fully activate the product.

Apply this product when the wind is calm. Do not apply this product onto desirable plants or shrubs as damage may result. If the product gets onto other plants, immediately brush or wash it off the foliage.

Apply 3.5 lbs. of Lebanon Confront* IV + Fertilizer per 1,000 sq. ft. of turfgrass. Use a drop or rotary spreader designed to apply granular fertilizers. Apply evenly and uniformly over the lawn area for best results. Avoid streaking, skips or overlaps as poor weed control or turfgrass injury may occur.

USE REGIONS WITHIN THE UNITED STATES



North
 South

SUGGESTED SPREADER SETTINGS

(Editor's Note: Manufacturer to supply a listing of spreaders and suggested settings to apply this product at a rate of 3.5 lbs. per 1,000 sq. ft.)

4/13


These settings are approximate. Rate of walk, moisture and spreader condition can vary the spreader setting necessary to apply the correct amount. Before each application, calibrate your spreader according to manufacturer's directions. Adjust to apply at the rate of 1 lb. of product per 286 sq. ft. Check equipment frequently to verify calibration.

Broadcast Spreader

Settings

Optional Spreading Instructions:

[Apply in a single pass at a rate of 3.5 lbs. per 1,000 sq. ft. Start on the outside border and apply in a circular path working toward the center. Overlap the coverage slightly, and shut off spreader before making hard turns.]




Graphic Optional

Drop Spreader

Settings

Optional Spreading Instructions:

[Apply in a single pass at a rate of 3.5 lbs. per 1,000 sq. ft. Start on the outside border and apply in a rectangular path working back and forth across the length of the lawn. Overlap the coverage slightly, and shut off spreader before making hard turns.]



Graphic Optional

If your spreader model is not listed (or does not appear above), go to our website at www.lebsea.com (or www.preen.com), or call toll-free (800) 233-1067, 8 A.M. - 5 P.M. EDT, Monday - Friday, for additional model settings.

WARRANTY

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Manufacturer, and Buyer assumes the risk of any such use.

Additional Labeling Requirements:

If the fertilizer guarantee contains iron, the following statement must appear:

This product contains the secondary nutrient iron. Iron may stain concrete and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash off with water.

If the recommended use rate applies less than 1 pound nitrogen per 1,000 sq. ft. per application, the following statement must appear:

Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

For products distributed in the State of Washington, the following statement must appear:

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-lebsea.com>

Advertising claims that may be presented on the label

For the postemergence control of annual and perennial broadleaf weeds, and preemergence control of annual grassy weeds and labeled broadleaf weeds, in established ornamental turf.

- Eliminates difficult, hard to kill weeds like clover, dandelion, chickweed and plantain that other products miss
- Eliminates dandelions, clover, chickweed, goldenrod, henbit and other troublesome lawn weeds
- Combination of four weed control technologies kills more weeds than most other brands
- Season long weed preventer
- Now with Confront IV for improved control of broadleaf and grassy weeds
- Unlike most weed & feed products, won't harm many warm season turf grasses
- One product kills broadleaf weeds and prevents crabgrass
- Apply once in spring when weeds are growing
- Stops more weeds in one application
- Works season long

Advertising claims that may be presented on the label (continued)

13/13

- Controls tough grasses and weeds [such as clover, crabgrass, goldenrod, henbit, etc.]
- Apply at any time [spring, summer, fall]
- Use for a [beautiful] weed free lawn
- Easy to use
- Just one application kills weeds missed by competitive brands and fertilizes into summer!
- Better weed control and fertilizer than other leading brands
- Extended, time release feeding of nitrogen lasts into summer
- Effective weed control leads to greener, thicker lawns
- Apply anytime when weeds are actively growing
- All the fertilizer nutrients your lawn needs for spring
- Feeds, greens, strengthens, and thickens your lawn for up to 3 months – longer than leading competitive brands
- Apply once for the whole season
- Feeds and strengthens root systems without burning your lawn
- Fertilize Once for Whole Season - Feeds and greens with slow release fertilizer for up to three months
- Fertilizes to fill in bare spots and provide plush green lawns
- Prevents weeds & fertilizes the lawn
- Fertilizes to provide rich color and vigorous growth
- Effective weed control - prevents weeds all season long
- New and improved
- Season long weed preventer and fertilizer
- Controls weeds longer
- Fertilizes longer

PREEN® GUARANTEE

Lebanon Seaboard Corporation guarantees you will be satisfied with the performance of this product when used as directed, or your money back. If not satisfied, describe the nature of your problem, and send with proof of purchase to: Lebanon Seaboard Corporation, 1600 East Cumberland Street, Lebanon, PA 17042.

Preen® is a registered trademark of Lebanon Seaboard Corporation.