

DIRECTIONS

LACCO MAGIC SULPHUR is prepared for use against certain pest and diseases of crops as listed below. Unless otherwise stated, dosages are given in terms of pounds of this product per acre. Do not apply if temperatures during or within 3 days after application are expected to exceed 90 F. in the area to be treated. Use as directed only when the pests or diseases appear.

BEANS: Powdery Mildew, Spider Mite, 15 to 20 pounds. Do not apply to varieties susceptible to sulphur injury.

MELONS: Powdery Mildew, 15 to 20 pounds, Spider Mite, 20 to 25 pounds. Do not apply to varieties susceptible to sulphur injury.

CUCUMBER, SQUASH: Powdery Mildew, 15 to 20 pounds. Do not apply to varieties susceptible to sulphur injury.

ONION, GARLIC: Petrobia Mite, 25 to 30 pounds.

SUGAR BEET: Spider Mite, 45 to 50 pounds.

TOMATO: Russet Mite, 30 to 40 pounds. If the crop is being grown with the aid of stakes or poles, apply 10 to 13 pounds per acre per foot of average plant height.

CARROT: Petrobia Mite, 30 to 40 pounds.

APPLE: Scab, 30 to 35 pounds at green-tip and again at pink-bud. Pink-bud treatments also control mildew. Do not apply during or shortly before frost conditions and/or the use of captan.

BUSHBERRIES: Powdery Mildew, 15 to 20 pounds during the pre-harvest season.

CHERRY: Brown Rot, 40 to 50 pounds 2 to 3 weeks pre-harvest.

GRAPE: Powdery Mildew, 5 to 10 pounds when new shoots are 6, 12, and 18 inches long. Thereafter, repeat each 10 to 14 days to harvest. Do not apply if temperatures during or shortly after treatment are expected to exceed 100 F. When applying sulphur to grapes, do not allow workers to re-enter the field within 1 day (24 hours) of application.

FIG: Fig Rust Mite, 40 pounds in the spring and again in the early summer.



ACTIVE INGREDIENT:	%% BY WT.
Sulphur	98.0%
INERT INGREDIENTS	2.0%

CAUTION
KEEP OUT OF REACH OF CHILDREN

DESTROY THIS CONTAINER WHEN EMPTY
KEEP CONTAINER CLOSED. DO NOT LEAVE IN SUNSHINE.

LOT NUMBER. NET WEIGHT
50 POUNDS

CTTTT1071 MANUFACTURED BY

LOS ANGELES CHEMICAL COMPANY

ACCEPTED
JUN 7 1972 962 73

42-73
ACCEPTED
5/11/75

DIRECTIONS—Continued

PEACH, NECTARINE: Mildew, 50 pounds at petal-fall, 2 weeks later and again at beginning of pit-hardening. Rust, mildew, 50 pounds after pit-hardening. Peach Silver Mite, 25 to 35 pounds after thinning. Brown Rot, 50 to 60 pounds 4 weeks pre-harvest and again 2 weeks later. When applying sulphur to peaches and nectarines, do not allow workers to re-enter the field within 1 day (24 hours) of application.

PLUM, PRUNE: Brown Rot, Prune Rust, 30 to 35 pounds 5 weeks before harvest. Repeat each 7 days if wet weather prevails during the period prior to harvest.

STRAWBERRY: Mildew, 10 to 20 pounds pre-harvest.

ALFALFA (grown for seed): Atlantic Mite, 15 to 25 pounds when first observed in flower heads and before defoliation of more than the bottom leaves had occurred. Repeat as needed throughout seed-set. May cause some flower burn.

COTTON: Atlantic Mite 45 to 50 pounds when first observed and before defoliation has occurred. Repeat as needed.

CITRUS: Six-spotted Mite, Citrus Rust Mite, 80 pounds from November to June. Citrus Flat Mite, 100 pounds from October to June. Yuma Spider Mite, 100 pounds from October to March 15. Citrus Thrips, 100 pounds at mid-March, petal-fall and again 2 to 4 weeks later. Do not apply within 2 months of oil sprays.

ORNAMENTALS: Powdery Mildew of Alder Azalea, Bittersweet, Boston Ivy, Buttonbush, Catalpa, Chrysanthemums, Dahlia, Delphinium, Dogweed, Honeysuckle, Horse Chestnut, Hydrangea, Lilac, Linden, Magnolia, Matrimony Vine, Phlox, Rhododendron, Rose, Staghorn Sumac, Sweet Pea, Sycamore, Trumpet Vine, Witch-Hazel, Zinnia; Rust of Aster, Carnation, Chrysanthemum, Rose, Snapdragon, 20 to 40 pounds.

NOTICE: Recommendations for the use of this product are based upon information believed to be reliable at time of printing. The use of this product being beyond the control of LOS ANGELES CHEMICAL COMPANY, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe and sound practice. THE BUYER must assume all responsibility including injury and/or damage, resulting from its misuse as such or in combination with other products.

