

Toxic to wildlife, do not contaminate streams, lakes or ponds. Do not apply to water. Do not apply to treated areas may be killed. Do not apply when water is present. Do not apply to treated areas. Do not contaminate water by cleaning of equipment or disposal of product only as specified on the label. NOTICE: Before using this product in the State of California, the user must obtain a permit from the State Agricultural Commissioner in the County where such use is to be made.

DESTROY CONTAINER WHEN EMPTY STOP ONLY IN ORIGINAL CONTAINER KEEP CONTAINER CLOSED

HERBICIDE DIRECTIONS

LACCO CALCIUM ARSENATE is prepared for use as a selective herbicide in ornamental lawn and turf as listed below. Dosages are given in pounds of this product per 1000 square feet. When used as an insecticide use sufficient water to provide uniform coverage. Unless otherwise noted wash treatment into the soil immediately after application. Keep children and pets off treated areas until this product is washed into soil and grass has dried. Do not use near bee hives. Do not use high phosphorus fertilizers the first year of treatment.

HAIKY CRABGRASS, SMOOTH CRABGRASS in established Kentucky bluegrass or ryegrass in sandy or infertile soil, 8-12 lbs. early spring before germination or late fall. In clay or fertile soil, 12 lbs. early spring before germination or late fall.

ANNUAL BLUEGRASS, in established Kentucky bluegrass or ryegrass, 6-8 lbs. in early spring and again in late fall.

MOUSEEAR CHICKWEED, SMOOTH CHICKWEED, in established Kentucky bluegrass or ryegrass, 12-16 lbs. in early spring. Apply when weeds are damp, do not rinse foliage, some temporary leaf scorch will occur.

Maintenance application of 3-6 lbs., in spring or fall will be needed to suppress the problem. Any of the above named treatments also controls Japanese beetles and Asiatic beetle.

CRABGRASS, ANNUAL BLUEGRASS in established bentgrass, bermuda grass, and fescue, 3-4 lbs. in fall and early spring until control is obtained.

INSECTICIDE DIRECTIONS

LACCO CALCIUM ARSENATE is prepared for use as a concentrate for formulation into a bait. Thoroughly blend as indicated, the following ingredients: Bran, 80% by wt., LACCO CALCIUM ARSENATE, 10% by wt. and Molasses, 10% by wt. Apply to the following crops at the rate of 40-70 lbs. per acre to the soil for the control of slugs, snails, sowbugs, cutworm and armyworm. Care should be taken when applying, to keep the bait off the foliage or other plant parts.

TREE CROPS:

Almond, apple, apricot, avocado, cashew, cherry, chestnut, date fig, filbert, grapefruit, guava, hazelnut, hickory nut, lemon, lime, mango, nectarine, olive, orange, papaya, passion fruit, peach, pear, pecan, persimmon, plum, quince, tangerine, tangelo, walnut.

VEGETABLE CROPS:

Asparagus, beans, blackeyed peas, broccoli, brussel sprouts, cabbage, carrot, cauliflower, celery, collards, corn, cowpeas, cucumber, eggplant, kale, kohlrabi, peppers, pumpkin, rutabaga, spinach, squash, tomato, turnip.

FLYLARVAE CONTROL: 2 1/4 lb. per 1000 square feet of poultry droppings as a coarse wet low pressure spray to the surface of the droppings under birds in cages or on wire floor. Use sufficient water to provide uniform coverage.

NOTICE: Recommendations for the use of this product are based upon information believed to be reliable at time of printing. The use of this product being beyond the control of LOS ANGELES CHEMICAL COMPANY, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe and sound practice. THE BUYER must assume all responsibility including injury and/or damage, resulting from its misuse as such or in combination with other products.

LOT NO. BTTTB670 MANUFACTURED BY NET WEIGHT 50 POUNDS

