

BACK PANEL

DIRECTIONS - TREE & VINE CROPS

LACCO COPRO 53 is prepared for use against certain diseases of crops as listed below. Unless otherwise noted, dosages are given in pounds of this product per 100 gallons of water to be applied as full coverage sprays with conventional spray equipment. Maximum dosages refer to pounds of this product per acre per application.

ALMOND: Brown rot, 4 to $4\frac{1}{2}$ lbs. at pop corn. Do not apply after pop corn or exceed 30 lbs. per acre.

APPLE: Fire blight, $\frac{1}{4}$ lb at full bloom, petal fall and late secondary bloom. Do not exceed $6\frac{1}{2}$ lbs per acre.

APRICOT: Brown rot, 4 to $4\frac{1}{2}$ lbs at red bud, early bloom and again at full bloom. Shot hole disease 4 to $4\frac{1}{2}$ lbs. November 15 to December 15. Do not apply after petal fall during the growing season or exceed 30 lbs. per acre.

BLACKBERRY, RASPBERRY: Leaf and cane spot, yellow rust, $1\text{-}3/4$ to $1\text{-}7/8$ lbs when new laterals leaf out and again when flowers begin to open. After harvest, prune and remove old fruit wood and before fall rains apply $3\frac{1}{2}$ to $3\text{-}3/4$ lbs. Do not exceed $5\text{-}3/4$ lbs. per acre.

CHERRY: Brown rot, leaf spot, 4 to $4\frac{1}{2}$ lbs at pop corn, full bloom and again at petal fall. Do not apply after petal fall or exceed 25 lbs per acre.

CITRUS: Brown rot, $\frac{1}{2}$ to $1\frac{1}{2}$ lbs. as a directed spray applying 6 gallons per tree to the lower 3 to 4 feet of tree skirts and 2 to 4 gallons per tree to the trunk and soil or litter beneath the tree in November or December and again in January or February depending upon the amount of rain during this period. The addition of $\frac{1}{2}$ to $1\frac{1}{2}$ lbs of hydrated lime plus 4 oz of sticker adjuvant may increase the effectiveness of this operation. Do not apply in areas where copper injury is known to occur or where fumigation with hydrogen cyanide gas is practiced. Do not apply after fruit is $3/4$ inch in diameter.

PEACH, NECTARINE: Peach leaf curl, peach blight, 4 to $4\frac{1}{2}$ lbs November 15 to December 15, if this application cannot be made, make application just before bud swell. Peach blight, brown rot, 4 to $4\frac{1}{2}$ lbs. at pink bud. Do not apply after pink bud during the growing season. Do not exceed 18 lbs per acre on peach or 25 lbs. per acre on nectarine.

PEAR: Fire blight, $\frac{1}{2}$ lb. at 10% bloom, repeat each 5 days until all late bloom is over. If rain occurs during this period, shorten intervals to 3 days. If rain is forecast, and no spray was applied within the last 72 hours, apply immediately. Do not apply to Anjou or Comice varieties or exceed $5\text{-}5/8$ lbs. per acre.

PLUM: Brown rot, $5/8$ to $7/8$ lb. at pop corn and full bloom. Do not apply after petal fall or exceed $4\text{-}1/8$ lbs per acre.

WALNUT: Walnut blight, $2\text{-}3/8$ to $4\text{-}3/4$ lbs at 1% pistillate bloom and again at 10 to 20% pistillate bloom. Do not exceed 45 lbs. per acre.

FIELD & ROW CROPS

Dosages are given in pounds of this product in 100 gallons of water per acre uniformly applied when conditions favoring the disease occur.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER: Downy mildew, $3\text{-}3/4$ lbs.

CELERY, POTATO, TOMATO: Late blight, $4\text{-}3/4$ to $9\frac{1}{2}$ lbs each 7 days.

MELON, SPINACH: Downy mildew, $5\frac{1}{2}$ lbs.

SUGAR BEETS: Cercospora Leaf Spot, 3 to 4 pounds per acre in sufficient water to insure uniform coverage at 10 to 15 day intervals from about July 25 to about September 12.

ACCEPTED