

DIRECTIONS

Poisonous if swallowed, inhaled or absorbed through skin or eyes. Do not swallow, inhale or allow contact with skin, eyes or clothing. Wash thoroughly after handling and before eating or smoking. Do not contaminate food or feed with this product. Do not apply to or allow this product to drift onto crop or non-crop lands not specifically being treated. Do not apply to any crop not stated on the label. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label. Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes. LACCO BASIC LEAD ARSENATE is prepared for use against certain pests of crops listed below. Dosages are given in pounds of this product in 100 gallons of water to be applied as full coverage sprays with conventional equipment. Maximum dosages refer to pounds of this product per acre per application. This product may be toxic to honeybees. To protect bees or other pollinators apply this product early in the morning or late in the evening or when bees are not actively working the field.



BASIC LEAD ARSENATE

DANGER - POISON

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: % BY WT.
*Basic Lead Arsenate . . . 96.0%
INERT INGREDIENTS: . . . 4.0%

*Total lead as metallic, 66.82% by weight this product. Total arsenic as metallic, 14.5% by weight this product. Total water soluble arsenic as metallic, 0.35% by weight this product.

Not for use or storage in or around the home.

Toxic to fish and wildlife, do not contaminate streams, lakes or ponds with this product. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

APRICOT: Peach twig borer, 3 to 4 lbs. just before bloom and at petal fall. Do not apply within 30 days of harvest. Do not exceed 30 lbs. per acre. Do not graze livestock on treated forage. Remove excess residues at time of harvest.

GRAPE: Leafroller, leaf folder, 3 lbs. just before bloom. Do not apply after edible parts start to form nor exceed 15 lbs. per acre.

PLUM AND PRUNE: Peach twig borer, 3 to 4 lbs. just before bloom and again at petal fall. Eye-spotted budmoth, 3 lbs. as tips of leaves are pushing out from buds, repeat in 7 to 10 days. Do not apply within 30 days of harvest nor exceed 30 lbs. per acre. Do not graze livestock on treated area. Remove excess residues at time of harvest.

FIRST AID


If swallowed give victim 1 tablespoonful salt in glass of water. Repeat until vomit fluid is clear. After, give 2 tablespoonfuls Epsom salt or milk of magnesia and force fluids. Call a physician in all cases of suspected poisoning.

NOTICE

Recommendations for the use of this product are based upon information believed to be reliable at time of printing. The use of this product being beyond the control of LOS ANGELES CHEMICAL COMPANY, no guarantee expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe and sound practice. The Buyer must assume all responsibility including injury and or damage, resulting from its misuse as such or in combination with other products.



962-300



Net Wt. 50 lbs.

Lct No. 5114

LOS ANGELES CHEMICAL COMPANY

4415 ARDINE STREET • SOUTH GATE, CALIFORNIA 90280 • (213) 583-4700 • TELEX 151111