

IF SWALLOWED and victim is awake (conscious) make him vomit quickly. Induce vomiting by sticking finger down throat or by giving soapy or strong salty water to drink. Repeat until vomit is clear. Never give anything by mouth to an unconscious person. Have victim lie down and keep quiet. See doctor immediately.

ATROPINE IS AN ANTIDOTE. CONSULT PHYSICIAN FOR EMERGENCY SUPPLY.

If symptoms or signs of poisoning include blurred vision, abdominal cramps, and tightness in the chest, do not wait for a doctor but give two atropine tablets (each 1/100 grain or 0.65 milligrams) at once. (One tablet to children under five years of age)

NOTE TO PHYSICIAN

Antidote — administer atropine sulfate in large doses. Two to Four mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 10 minute intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. **DO NOT GIVE MORPHINE OR TRANQUILIZERS.** Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement. **VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.**

WARNING NOTICE

Balco Chemicals, Inc. warrants that this material conforms to the chemical description of the label and is reasonable fit for the purpose referred to in the directions for use. This product is sold subject to the understanding that the buyer assumes all risks of use or handling which may result in loss or damage which are beyond the control of the seller, such as for example incompatibility with other products, the manner of its use or application, or the presence of other products or materials in or on the soil or crop. Balco Chemicals, Inc. does not make any express or implied warranty of the fitness or merchantability. The exclusive remedy of the user is the return of the product to the seller at the option of the user and the limit of the liability of Balco Chemicals, Inc. is the purchase price paid by the user or buyer of the product. The user or buyer is responsible for any and all losses, injuries or damages resulting from the use or handling of this product. The user or buyer is responsible for the safe use of this product. The user or buyer is responsible for the safe use of this product. The user or buyer is responsible for the safe use of this product.

DIRECTIONS FOR USE

Read the precautionary statements before using.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT, OR OPEN FLAME

KEEP CONTAINER CLOSE. LEAKING PACKAGES SHOULD BE REMOVED TO A SAFE PLACE

Do not heat shell of drum or any part of contents above 200 degrees F. because of violent decomposition. Do not store at below freezing temperatures to avoid cry stabilization.

DO NOT USE IN UNDILUTED FORM NOT FOR USE OR STORAGE IN OR AROUND THE HOME

USAGE CAUTION: DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROP OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH DIRECTIONS BECAUSE PLANT INJURY, EXCESSIVE RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

Add the concentrate to the spray tank while filling with water, and mix thoroughly either by means of a tank agitator or pump by pass. For best results, thoroughly cover all surfaces to be treated with spray. Rates of application given below should not be exceeded.

POISON

See side panels for antidote and precautions

STOP! READ THE LABEL



CAN KILL YOU

"6-3"

PARATHION-METHYL PARATHION Emulsifiable Concentrate

ACTIVE INGREDIENTS:

Parathion (0,0-diethyl 0-p-nitrophenyl phosphorothioate)	58.1%
0,0 dimethyl 0-p-nitrophenyl phosphorothioate*	29.1%
Xylene	7.2%
INERT INGREDIENTS	5.6%

* Methyl parathion

CONTAINS 6 POUNDS PARATHION AND 3 POUNDS METHYL PARATHION PER GALLON

EPA Reg. No. 960-180

NET CONTENTS..... U. S. GAL.

DANGER—POISON—PRECAUTIONS



CAN KILL YOU IF SWALLOWED

This product can kill you if swallowed even in small amounts. Spray mist or dust may be fatal if swallowed.



CAN KILL YOU BY SKIN CONTACT

This product can kill you if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes).



CAN KILL YOU IF BREATHED

This product can kill you if vapors, spray mist or dust are breathed.


POISONOUS TO FISH AND WILDLIFE

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

NOT FOR HOME USE

DANGER

Keep out of reach of children



		per 100 Gals of Spray	
Apples (California Only)	Aphids, Codling Moths, Pear Tree Borer	1/4 - 1/2 pint / 100 gal of spray	Do not apply within 14 days of harvest. See restrictions under plums.
Apples	Aphids, Codling Moths, Pear Tree Borer	1/4 - 1/2 pint / 100 gal of spray	Do not apply within 14 days of harvest. See restrictions under plums.
Beans (Dry)	Aphids, Armyworms, Leafhoppers, Leaf Miners, Leaf Beetles, Mexican Bean Beetles, Red Spider Mites	1/2	
Beans (Green)	Aphids, Armyworms, Leafhoppers, Leaf Miners, Leaf Beetles, Mexican Bean Beetles, Red Spider Mites	1/2	Do not apply within 21 days of harvest.
Beets (Red)	Aphids, Blister Beetles, Weevils	1/2	Do not apply within 15 days of harvest. 21 days if tips are to be used as food or feed.
Cabbage and Cole Crops (Brussels, Brussels Sprouts, Cauliflower, Kohlrabi)	Aphids, Vegetable Weevils, Imported Cabbageworms, Cabbage Looper, Armyworms, Leaf Beetles	1/2 - 1/3	Do not apply within 21 days of harvest.
Carrots	Aphids, Vegetable Weevils, Leafhoppers, Petiole Borer	1/2	Do not use tips for food or feed. Do not apply within 15 days of harvest.
Celery	Aphids, Celery Leaf Ties, Cabbageworms, Spider Mites, Turnip Plant Bugs	1/2	Do not apply within 21 days of harvest.
Cauliflowers, Kale, Mustard Greens, Spinach	Aphids, Imported Cabbageworms, Leaf Miners, Leaf Beetles, Weevils, Cabbage Looper, Leafhoppers, Spider Mites, Turnip Plant Bugs	1/2	Do not apply within 21 days of harvest.
Corn (Sweet & Pop)	Aphids, Armyworms, European Corn Borer, Corn Earworm, Corn Rootworm, Spider Mites, Cabbage Looper, Leafhoppers, Leaf Beetles, Weevils, Cabbage Looper, Leafhoppers, Spider Mites, Turnip Plant Bugs	1/2	Apply when 75% of the plants show shoot hole feeding and follow with two additional sprays at 5 to 7 day intervals. Use sufficient water to provide complete coverage and be certain whorls of plants are well treated. Do not apply within 12 days of harvest, cutting or forage use.
Cotton	Aphids, Armyworms, European Corn Borer, Cotton Leafworms, Flea hoppers, Lygus Bugs, Red Spider Mites, Thrips, Garden Webworms, Cabbage Looper, Boll weevil, Bollworms, Budworms	1/2	For European Corn Borer control, consult your State Agricultural Experiment Station for proper timing of treatment.
Cucumbers	Aphids, Two Spotted Mites	1/2	Do not apply within 15 days of harvest.
Grapes	Aphids, Grape Leafhoppers	1/2 - 3/4 pint per acre	Do not apply within 15 days of harvest.
Hot Peppers	Aphids, Spider Mites	1/2 - 3/4 pint per acre	Do not apply within 15 days of harvest.
Lettuces, Endive, Escarole, Spinach	Aphids, Armyworms, Cabbage Looper, Imported Cabbageworm	1/2	Do not apply within 15 days of harvest.
Melons (Cantaloupe)	Aphids, Thrips	1/2	Do not apply within 15 days of harvest.
Onions	Onion Thrips	1/2	Do not apply within 15 days of harvest.
Peas (Garden and Blackeye)	Aphids, Pea Weevils, Armyworms	1/2	Do not apply within 15 days of harvest.
Peppers	Aphids, Leaf Miners	1/2	Do not apply within 15 days of harvest.
Peaches	Aphids, Peach Tree Borer, Plum Curculio, Oriental Fruit Moth	1/2 - 3/4 pint per 100 gal of spray	Do not apply within 14 days of harvest. See restrictions under plums.
Pears & Nectarines (California Only)	Aphids, Peach Tree Borer, Plum Curculio, Oriental Fruit Moth	1/2 pint per 100 gal of spray	Do not apply within 14 days of harvest. See restrictions under plums.
Plums	Aphids, Peach Tree Borer, Plum Curculio	1/2 pint per 100 gal of spray	Do not apply within 14 days of harvest. See restrictions under plums.
Potatoes	Aphids, Blister Beetles, Leafhoppers, Leaf Miners, Plant Bugs, Flea beetles, Thrips, Vegetable Weevils	1/2 - 1/3	Do not apply within 14 days of harvest.
Pumpkins	Aphids	1/2	Do not apply within 14 days of harvest.
Squash	Aphids, Armyworms, Flea Beetles, Leafhoppers, Leaf Miners, Leaf Beetles, Mexican Bean Beetles, Red Spider Mites	1/2	Do not apply within 14 days of harvest.
Strawberries	Aphids	1/2	Do not apply within 14 days of harvest.
Sugar Beets	Aphids, Armyworms, Flea Beetles, Leafhoppers, Leaf Miners, Leaf Beetles, Mexican Bean Beetles, Red Spider Mites	1/2	Do not apply within 14 days of harvest.
Sweet Potatoes	Aphids, Spider Mites, Leafhoppers	1/2 - 3/4	Do not apply within 15 days of harvest.
Tomatoes	Aphids, Armyworms, Flea Beetles, Leafhoppers, Leaf Miners, Leaf Beetles, Mexican Bean Beetles, Red Spider Mites	1/2	Do not apply within 15 days of harvest. 21 days if tips are to be used as food or feed.
Turnips	Aphids, Cabbage Weevils, Vegetable Weevils	1/2 - 1/3	Do not apply within 15 days of harvest. 21 days if tips are to be used as food or feed.