

	
ACTIVE INGREDIENTS:	% BY WT.
*Sulfur	95.0
INERT INGREDIENTS	5.0
*95% 325 Mesh or Smaller	

CAUTION - KEEP OUT OF REACH OF CHILDREN
(DESTROY THIS CONTAINER WHEN EMPTY)

DIRECTIONS

LACCO WETTABLE SULFUR is prepared for use against certain diseases and/or pests of crops and ornamental plants as listed below. Unless otherwise stated, dosages are in terms of pounds of this product per 100 gallons of water to be applied as a full coverage spray. Maximum dosages refer to pounds per each application. Do not apply with or within 30 days of oil sprays. Do not apply if temperatures during or within 3 days of application are expected to exceed 90° F. Use as directed only when the diseases and/or pests appear.

ALMONDS: Brown Rot, Shot Hole, apply at 2-6 lbs. per 100 gallons water. Begin when disease threatens or in pre-bloom stage.

GRAPE: Powdery Mildew, Rust (physopella), apply at 2-6 lbs. per 100 gallons of water. Begin when new shoots are 6 inches long. Repeat when blossoms are open and continue as necessary. Concord and other Labrusca type grapes may be injured by sulfur. When applying sulfur to grapes do not allow workers to re-enter the field within 1 day (24 hr.) of sulfur application.

APPLE: Apple Cedar Rust, Black Rot, Brown Rot, Frog Eye Leaf Spot, Powdery Mildew, Quince Rust, Scab, Sooty Blotch, apply 3½ to 8 lbs. per 100 gallons water from pre-pink to cover. MacIntosh, Golden Delicious, Jonathan and certain other varieties may be injured by sulfur under certain conditions.

CITRUS: Six Spotted Mite, Citrus Rust Mite, apply 4-8 lbs. of Wettable Sulfur per 100 gallons of water. Apply a minimum of 85 lbs. sulfur per acre. For Citrus Flat Mite and Yuma Spider Mite apply 4-8 lbs. Wettable Sulfur per 100 gallons of water. Apply a minimum of 105 lbs. per acre. Do not apply within 60 days of oil sprays.

PEACH, NECTARINE: Brown Rot, 6 lbs. at pink-bud and full-bloom, repeat using 10 lbs. 4 weeks before harvest and again 2 weeks later. **On cling varieties in the Sacramento Valley, only:** If rain occurs during the last 3 weeks before harvest, apply 5 lbs. plus 2 gallons LACCO LIQUID LIME SULFUR as directed. Caution, severe phytotoxicity may occur in some areas, following this use of lime sulfur. Powdery Mildew, 5 lbs. or 5 lbs. plus ½ to 1 gallon of LACCO LIQUID LIME SULFUR as directed at early petal fall. Rust, Peach Silver Mite, 5 lbs. or 5 lbs. plus ½ gallon LACCO LIQUID LIME SULFUR as directed after pit-hardening. When applying sulfur to peaches and nectarines, do not allow workers to re-enter the field within 1 day (24 hours) of applications.

PEAR: Scab (ALL varieties), 4 lbs. plus 2 gallons LACCO LIQUID LIME SULFUR as directed at green-tip. Repeat at cluster-bud on Bartlett Pear Scab, 4 lbs. at petal-fall and again 5 to 10 days later.

PLUM & PRUNE: Brown Rot, Prune Rust, 4 lbs. 5 weeks pre-harvest. Repeat each 5 days until harvest if wet weather prevails.

STRAWBERRY: Powdery Mildew, 1 to 5 lbs. per acre in sufficient water to assure uniform coverage. Begin at first sign of disease and repeat as necessary. May cause leaf burn in hot weather.

The Food and Drug Administration have established residue tolerances on most pesticides. Directions should be carefully followed to avoid excessive residue at time of harvest.

NOTICE: Recommendations for the use of this product are based upon information believed to be reliable at time of printing. The use of this product being beyond the control of LOS ANGELES CHEMICAL COMPANY, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe and sound practice. THE BUYER must assume all responsibility including injury and/or damage, resulting from its misuse as such or in combination with other products.

LOT NUMBER

NET WEIGHT
50 POUNDS

REF 10772

800-028

LOS ANGELES CHEMICAL COMPANY