ACCEPTED
Sept 4, 1974
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC PURPOSES REGISTERED UNDER NO. 962-377
TO ATTACHED LABEL

CAUTION

962-

Harmful if swallowed!
Avoid breathing dust.
Avoid contact with skin.
Wash thoroughly after use.
Avoid contamination of feed and foodstuffs.

KEEP CONTAINER CLOSED.

DO NOT LEAVE IN SUNSHINE.

DESTROY THIS CONTAINER WHEN EMPTY.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

DIRECTIONS

LACCO MALATHION 5 DUST is prepared for use against certain pests of crops, livestock and poultry as listed below. Unless otherwise stated, dosages are given in terms of pounds of this product per acre per each application. Use as directed only when the pests appear. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

BEANS: Mexican Bean Beetles, Japanese Beetles, Cucumber Beetles, Leafhoppers, Spider Mites, Aphids, 30-35 lbs. per acre. Two or more applications as needed. Do not apply within one day of harvest.

CELERY: Aphid, Whitefly, Leafhopper, 24 pounds. Apply as needed. Do not apply within 7 days of harvest.

CHERRY: Cherry Slug, 30-40 pounds. Apply 2 to 3 weeks pre-harvest and again post-harvest. May injure foliage on some varieties. Do not apply within 3 days of harvest.

REG. NO. 962-377-
EPA EST. NO. 962-CA-1

Lacco

MALATHION 5 DUST

ACTIVE INGREDIENT:	
* Malathion	5.0%
INERT INGREDIENTS	95.0%

* O, O-dimethyl dithiophosphate of diethyl mercap osuccinate.

CAUTION

KEEP OUT OF REACH OF CHILDREN

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely o occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NOTICE: Recommendations for the use of this product are based upon information believed to be reliable at time of printing. The use of this product being beyond the control of LOS ANGELES CHEMICAL COMPANY, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe and sound practice. The BUYER must assume all responsibility including injury and or damage resulting from its misuse as such or in combination with other products.

LOT NUMBER **NET WEIGHT**

CTTTB774 **MANUFACTURED BY**

DIRECTIONS (Continued)

COTTON: Cotton Aphid, 30-40 pounds. Apply before honeydew formation begins. May be applied up to harvest.

GRAPE: Variegated Grape Leafhopper, 20 pounds. Apply 14 days after appearance and again post-bloom to pre-harvest at 20-25 lbs. In the Coachella Valley, also apply post-harvest at 12-16 pounds. Grape Leafhopper, apply 16-20 pounds after growth starts to pre-bloom and again post-bloom to pre-harvest at 24 to 28 pounds. Vinegar or Pomace Fly, 20-28 pounds, apply near harvest. Use only when temperature is over 80 F. Do not apply within 3 days of harvest.

CUCUMBERS, SQUASH, MELONS: Aphids, Spider Mites, Pickleworms, 30-35 lbs. per acre. Cantaloupe Leafhoppers, 30-35 lbs. per acre as necessary. Do not apply unless plants are dry nor within 1 day of harvest.

LETTUCE: Loopers, apply 25-30 lbs. per acre. Do not apply within 14 days of harvest of leaf lettuce and 7 days on head lettuce.

POTATO: Leafhopper, apply 25-30 pounds during dormant, pre-harvest and harvest seasons. Fleabeetle, apply 40 pounds during dormant, pre-harvest and harvest seasons. Lygus Bug, Strawberry Whitefly, apply 40 pounds during the harvest season. Do not apply within 3 days of picking.

DAIRY CATTLE (Lactating): Hornfly, 3 tablespoons per animal. Sprinkle on the back and neck during the warm season. Repeat each 10 to 14 days. Do not contaminate milk, utensils, feed or water. Do not apply within 5 hours of milking.

TURKEYS: Northern Fowl Mite, Lice, apply to areas of vent, fluff and under wings or apply by using a dust-box or dusting wallow at the rate of 12 pounds per 100 birds. Do not contaminate eggs, feed or water.

800-024

LOS ANGELES CHEMICAL COMPANY