

BEST DOCUMENT AVAILABLE



E.P.A. REG. NO. 962-402

E.P.A. EST. NO. 962-CA-1

DIAZINON AG 500 INSECTICIDE

For Agricultural/Commercial Use Only

ACTIVE INGREDIENTS:

0,0-Diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl phosphorothioate)	47.5%
XYLENE	39.4%
INERT INGREDIENTS:	13.1%
Total	100.0%

LACCO DIAZINON AG 500 CONTAINS 4 LBS. DIAZINON PER GALLON

KEEP OUT OF REACH OF CHILDREN WARNING

See side panel for additional precautionary statements.

PRACTICAL TREATMENT: (FIRST AID)

IF SWALLOWED Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. **DO NOT INDUCE VOMITING** unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED Remove victim to fresh air. Give artificial respiration if victim stops breathing. Get medical attention immediately.

IF ON SKIN Remove contaminated clothing and wash contaminated skin with soap and water.

IF IN EYES Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

LOT NO.

NET CONTENTS:
1 GALLON

LOS ANGELES CHEMICAL COMPANY
4545 Ardine St., South Gate, California 90280

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Keep out of reach of children. May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly before smoking or eating and after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not contaminate ornamental fishponds. Do not use on humans, household pets or livestock. Keep children and pets out of treated areas until spray has dried.

PHYSICAL OR CHEMICAL HAZARDS

Do not use near spill or where near heat or open flame.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife feeding in treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes, and estuaries (except for purposes from treated waterways). Do not apply to water that will be used for recreational purposes and human and livestock consumption. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, crab and other aquatic life are important resources. This product must not be used in areas where impact on threatened endangered species is likely. Contact your State Fish and Game Agency before applying this product. Apply this product only as specified on the label.

GENERAL INFORMATION

LACCO Diazinon AG 500 is an emulsifiable solution which when diluted with water according to the directions, is intended for the control of insect pests of agricultural crops. Do not use in barns or dwellings. Do not use on livestock.

It is the manufacturer's intent that best control is obtained when application is made at first sign of insects. Application should be repeated as necessary to maintain control. NOTE: The California Department of Agriculture has set a 5-day reentry for Diazinon on citrus, grapes, peaches and nutmeats. Five days should elapse between the time of application and the time a worker may enter the field to engage in any activity requiring substantial contact with treated foliage. When a mixture of two or more organic phosphorus pesticides is applied, the interval shall be prolonged by adding to the longest applicable interval an additional 50% of that interval.

INSTRUCTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

VEGETABLE PLANT SUBJECTS

Citrus: Citruses should follow application of LACCO Diazinon AG 500 should always be applied to vegetable crops in a spray in a minimum of 2 gal. of water per acre for aerial application and a minimum of 10 gal. of water per acre for ground applications.

BEANS: Pinto Beans, Snap Beans, Lima Beans, Black Bean Aphids, Mexican Bean Beetle, Leaf-miners. Apply 1/2 pt. per acre. Cucumber Beetles. Apply 1/2 pt. in 100 gal. of water per acre. Do not apply within 7 days of bean harvest. Bean foliage or bean hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment of foliage to be cut for hay.

LIMA BEANS: Deteriorate Leafminers, Leafhoppers, Mexican Aphid. Apply 1 pt. per acre. Do not apply within 7 days of harvest. Bean foliage or bean hay may be fed to dairy cattle and sheep. Wait 4 days following treatment of foliage to be cut for hay.

BROCCOLI, CABBAGE, CABBAGEFLOWER: Aphids, Diamondback Moths, Imported Cabbageworms. Apply 1/2 pt. per acre. Applications at 7-10 day intervals will usually be required. Do not apply to broccoli or cauliflower within 5 days, or to cabbage within 7 days of harvest.

BROCCOLI SPINNY: Aphids, Diamond Moths, Imported Cabbageworms. Apply 1 pt. per acre. Generally applications at 7-10 day intervals will be required. Do not apply to broccoli sprouts within 7 days of harvest.

CARROTS: Aphids. Apply 1 pt. per acre. Do not apply within 10 days of harvest.

CELERY: Aphids, Flea Beetle. Apply 1 pt. in 100 gal. of water per acre. Do not apply within 10 days of harvest. Plants should be market trimmed (leaves removed) before being stored or used. Foliage should not be used for feed or food.

COLLARDS, RAIL, GREEN CHINA, TURNIP TOPS: Aphids, Imported Cabbageworm, Diamondback Moth, variegated Cabbage Moth. Apply 1/2 pt. per acre. Do not apply to collards, kale or turnip tops within 10 days, or to Brassica Chard within 12 days of harvest.

CUCUMBERS: Aphids, Serpentine Leafminers. Apply 1 pt. per acre. Do not apply within 7 days of harvest.

LETTUCE, ENDIVE: Aphids, Deteriorate Leafminers. Apply 1/2 pt. per acre. Do not apply within 10 days of harvest.

MELONS, SQUASH, CANTALOUPE, CASABA, CUCUMBER, HONEYDEW MELONS, BU LARBELLS, PEACH MELONS, AND HYDRANGEA OF THE SE. WATERMELONS AND THEIR HYBRIDS, SUMMER AND WINTER SQUASH: Aphids, Leafhoppers, Moths. Apply 1/2 pt. per acre. Serpentine Leafminers. Apply 1/2 pt. per acre. Do not apply to melons and water squashes within 7 days, or to winter squashes within 7 days of harvest.

ONIONBULB VEGETABLES: Onion Root Fly, Fleas and Sciarids. Apply 1/2 pt. in 50 Gal. of water (1/2 to 1 qt. of Diazinon) and spray on outside and inside of bulb and neck of onion. Do not spray onion bulb or neck after harvest. Do not use on onion bulb or neck after harvest. Do not spray onion bulb or neck after harvest. Do not use on onion bulb or neck after harvest. Do not spray onion bulb or neck after harvest. Do not use on onion bulb or neck after harvest.

ONIONS: Onion Root Fly. Apply 1 pt. per acre. Do not apply within 10 days of harvest.

PEAS: Aphids, Leafminers. Apply 1/2 pt. per acre. Peas may be picked immediately. The main and side stems may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment of foliage to be cut for hay.

PEPPERS: Aphids, Serpentine Leafminers. Apply 1/2 pt. per acre. Do not apply within 7 days of harvest.

POTATOES: Aphids, Flea Beetles, Colorado Potato Beetle, Deteriorate Leafminers. Apply 1/2 pt. per acre. Do not apply within 35 days of harvest.

RADISHES, PARSNIPS, TUBEROSES: Aphids, Flea Beetles, Leafhoppers. Apply 1/2 pt. per acre. Do not apply within 10 days of harvest.

SPINACH, BEETS: Aphids, Deteriorate Leafminers. Apply 1/2 pt. per acre. Do not apply within 7 days of harvest.

SQUASH: Cucumber and Winter Squash Moths and Sciarids.

