

PM 25 962-406

ACCEPTED
DEC 22 1983
Under the Federal Insecticide
Fungicide and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 962-406

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
Harmful if swallowed. Avoid contact with skin, eyes or clothing. In case of contact with eyes, immediately flush with plenty of water. Get Medical attention if irritation persists. Wash thoroughly after handling. Keep livestock off treated areas.

ENVIRONMENTAL HAZARDS
LACCOBOR-CHLORATE READY-TO-USE solutions carelessly spilled or applied to cropland or growing plants — including trees and shrubs — may kill or seriously retard growth. Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL
This product may crystallize if stored at temperatures below 0°C (32°F.). Do not contaminate water, food or feed by storage or disposal. Do not store where children or animals may gain access. Do not reuse empty container. Wrap container and put in trash collection. This is a ready-to-use general herbicide, nonselective in action, destroying broad and narrow leaf types of grasses and weeds.

ANNUAL WEEDS
LACCOBOR-CHLORATE-READY-TO-USE, no mixing required. Apply 1 gallon of solution as a contact spray thoroughly wetting all parts of the plants particularly the growing tips. Apply 1 gallon solution per 75 square feet when the plants are young and tender (1 to 4 inches high).



EPA Reg. No. 962-406AA
EPA Est. No. 11474-CA-01
LACCOBOR-CHLORATE
READY-TO-USE
GRASS AND WEED KILLER

FOR USE ON: Fence Lines, Asphaltic Driveways, Industrial Sites, Drainage Ditch Banks and Similar Non-Crop Areas.
CONTROLS: Johnson Grass, Leafy Spurge, Canadian Thistle, Bindweed, Bermuda Grass and quackgrass.

ACTIVE INGREDIENTS:	
Disodium Octaborate Tetrahydrate (Na ₂ B ₈ O ₁₃ · 4H ₂ O)	8.20%
Sodium Chlorate (NaClO ₃)	2.81%
Boron Trioxide (B ₂ O ₃) Equivalent	5.5%
Boron (B) Equivalent	1.7%
INERT INGREDIENTS:	88.99%
Total 100.00%	

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

LOT NUMBER
NET CONTENTS
1 GALLON

LOS ANGELES CHEMICAL COMPANY • 4545 ARDINE STREET, SOUTH GATE, CALIFORNIA 90280

PERENNIAL WEEDS
LACCOBOR-CHLORATE may be applied in late summer, fall, or early spring stages or in the fall months and spring applications. Apply ready-to-use at the rate of 1 gal Johnson grass, leafy spurge, Canadian Thistle, Grass and quackgrass.
FOR ASPHALTIC DRIVEWAYS, apply at 1 square foot, make sure to thoroughly wet weeds have emerged.

NOTICE
This material is non-selective in action and vegetation and may render the entire treated area unproductive for one or more years. Care should be taken to confine the use of this material to the area intended to be treated and to avoid its use on crops and other desirable plants which are areas which may be underlain by roots of trees. Because of the many varying conditions of vegetation, the manufacturer warns that conditions may impair or vary the results of product in any event, complete elimination is not guaranteed. It is advisable to consult your Agricultural Experiment Station weed control problems. Our recommendations for use of this product are believed to be reliable. The use of this product being beyond the scope of no guarantee, expressed or implied, is made or the results to be obtained if not used in accordance with established safe practice. The buyer must include injury or damage, resulting from combination with other materials.