

# WHITMOYER

# SAN-O-FEC®

## Disinfectant, deodorant and sanitizer

### ACTIVE INGREDIENTS

Methyldodecylbenzyl trimethyl ammonium chloride	16%
Methyldodecylxylylene bis (trimethyl ammonium chloride)	4%

### INERT INGREDIENT

Water	80.0%
-------	-------

### Phenol Coefficient

(10 min. standard A.O.A.C. method 20 C)

<i>S. typhosa</i>	50
<i>S. aureus</i>	80

EPA Reg. No. 984-19AA

**DANGER:** Keep out of reach of children. See First Aid statement and other precautions on back panel.

NET CONTENTS 1 GALLON (U.S.)



For measuring purposes, the following table may be used:  
1 pint = 16 ounces  
1 quart = 32 ounces  
1 standard measuring cup = 8 ounces  
1 gallon = 128 ounces  
1 standard measuring cup = 8 ounces  
1 quart = 32 ounces  
1 gallon = 128 ounces  
1 standard measuring cup = 8 ounces  
1 quart = 32 ounces  
1 gallon = 128 ounces

**For general disinfection and sanitation**—Mix 1 ounce SAN O FEC in 4 gallons of water for usual sanitizing applications. For disinfecting previously soiled porous surfaces, mix 1 ounce SAN O FEC in 4 gallons of water. For disinfecting previously cleaned porous surfaces, mix 1 ounce SAN O FEC in 4 gallons of water.

**In the poultry industry**—For dipping hatching eggs, rinsing egg trays, feeders, brooders, batteries, egg crates, coops, spraying or washing walls and floors. Mix 1 ounce SAN O FEC in 8 gallons of water to get sanitizing solution. Use double strength (1 ounce in 4 gallons of water) if water present in previous flocks. Hatchability is not affected when eggs are washed with SAN O FEC. Do not use for dipping of eggs for human consumption.

**Application Procedures** for proper use of SAN O FEC.

**In the swine industry**—For disinfecting farrowing pens, rearing pens, teat waterers, mix 1 ounce SAN O FEC in 4 gallons of water. See **Application P** for proper use of SAN O FEC.

**For pets**—For disinfecting and deodorizing cages, kennels, feeding pans, equipment, mix 1 ounce SAN O FEC in 4 gallons of water.

### APPLICATION PROCEDURES

For proper use of SAN O FEC in poultry and livestock premises, do the following:

1. Remove all animals, poultry and feed from premises, carts, boats, and other equipment.
2. Remove all litter and manure from floors, walls and surfaces of buildings, chutes and other facilities and fixtures, occupied or traversed by animals and poultry.
3. Empty all troughs, racks and other feeding and watering equipment.
4. Saturate all surfaces with the SAN O FEC solution.
5. Soak ropes, mats, ropes and other types of equipment, including restraining equipment, foot tanks, and all other equipment that may be used.

## DIRECTIONS FOR USE

For measuring purposes, the following table may be used:

1 table spoonful	1 ounce	8 ounce	1 standard measuring cup
------------------	---------	---------	--------------------------

Clean all surfaces and thoroughly rinse with soap or detergent before applying SAN-O-FEC solution. Do not mix with soap or synthetic detergents.

**For general disinfection and sanitation**—Mix 1 ounce SAN-O-FEC in 8 gallons of water for general sanitizing applications. For disinfecting previously cleaned non-porous surfaces, mix 1 ounce SAN-O-FEC in 4 gallons of water. For disinfecting previously cleaned porous surfaces mix 1 ounce SAN-O-FEC in 2 gallons of water.

**In the poultry industry**—For dipping hatching eggs, rinsing egg trays, fountain feeders, brooders, batteries, egg crates, coops, spraying or washing incubators, walls and floors. Mix 1 ounce SAN-O-FEC in 8 gallons of water to prepare the sanitizing solution. Use double strength (1 ounce in 4 gallons of water) if infection was present in previous flocks. Hatchability is not affected when eggs are treated with SAN-O-FEC. Do not use for dipping of eggs for human consumption. See **Application Procedures** for proper use of SAN-O-FEC.

**In the swine industry**—For disinfecting farrowing pens, rearing pens, feeders and waterers, mix 1 ounce SAN-O-FEC in 4 gallons of water. See **Application Procedures** for proper use of SAN-O-FEC.

**For pets**—For disinfecting and deodorizing cages, kennels, feeding pans and other equipment, mix 1 ounce SAN-O-FEC in 4 gallons of water.

### APPLICATION PROCEDURES

For proper use of SAN-O-FEC in poultry and livestock premises, observe the following:

1. Remove all animals, poultry and feed from premises, cars, boats, truck and other equipment.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals and poultry.
3. Empty all troughs, racks and other feeding and watering equipment.
4. Saturate all surfaces with the SAN-O-FEC solution.
5. Immerse all baskets, trays and other types of equipment used in handling and restraining animals. Treat forks, shovels and scrapers the same way.

6. Ventilate buildings, coops and other closed spaces. Do not house poultry or livestock or employ equipment until treatment has been absorbed, set or dried.

7. All treated food racks, hoppers, troughs, water troughs, feeders, fountain and waterers must be thoroughly rinsed with potable water prior to reuse.

**In packing plants and storage plants**—For sanitizing counters in packing and storage plants, mix 1 ounce SAN-O-FEC in 8 gallons of water and apply. Flush and rinse with clear water. On surfaces difficult to clean, increase strength of sanitizing solution to 1 ounce SAN-O-FEC in 2 gallons of water. Flush surfaces with a potable water rinse before reuse. For reman walls and floors, mix 1 ounce SAN-O-FEC in 2 gallons of water.

### NOT FOR USE OR STORAGE IN OR AROUND THE HOME ENVIRONMENT STORE ABOVE FREEZING TEMPERATURES

**DANGER:** Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

**FIRST AID**—In case of contact, immediately flush eye or skin with plenty of water for at least 15 minutes. For eye, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, promptly drink a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water before disposal.

LOT NO. D-131

Manufactured by

CODE NO. 4-63

**WHITMOYER**  
LABORATORIES, INC.  
MILWAUKEE, WISCONSIN  
A DIVISION OF ROHM AND HAAS COMPANY

