

WHITMOYER

SAN-O-FEC[®]-50

Disinfectant, deodorant and sanitizer

ACTIVE INGREDIENTS

Methyldodecylbenzyl trimethyl ammonium chloride 40%
Methyldodecylxylene bis (trimethyl ammonium chloride) 10%

INERT INGREDIENT

Water 50%

Phenol Coefficient

(10 min. Standard A.O.A.C. method 20°C)
S. typhosa 125
S. aureus 200

EPA Reg. No. 984-33A

DANGER: Keep out of reach of children. See First Aid statement and other precautions on back panel.

NET CONTENTS 1 GALLON (U.S.)



For general disinfection and sanitizing of water, toilet bowls, bathing tubs, showers, and kitchen and bathroom fixtures, use 1/2 ounce per gallon of water. For previously treated porous water.

In the Poultry Industry—For feeders, drinkers, batteries, pans and floors. Mix 1 ounce per gallon of water. If feces was present in previous flock, use 2 ounces per gallon with SAN-O-FEC-50. Do not use on birds.

Application Procedures for Poultry—

In the Swine Industry—For waterers. Mix 1 ounce per gallon of water.

For Pets—For disinfecting a pet's environment.

For proper use of SAN-O-FEC-50, follow these directions:

1. Remove all animals from the area to be treated.
2. Remove all animals from the area to be treated.
3. Remove all animals from the area to be treated.
4. Remove all animals from the area to be treated.
5. Remove all animals from the area to be treated.
6. Remove all animals from the area to be treated.
7. Remove all animals from the area to be treated.
8. Remove all animals from the area to be treated.
9. Remove all animals from the area to be treated.
10. Remove all animals from the area to be treated.

For more information on the following information, see the following references: *Journal of the National Sanitation Foundation*, *Journal of the National Poultry Improvement Association*, and *Measuring Up*.

Wash all surfaces thoroughly with a heavy soap or detergent before applying SAN-O-FEC-50 solution. Do not mix with soap or synthetic detergents.

For general disinfection and sanitation—Mix 1 ounce SAN-O-FEC-50 in 20 gallons of water for disinfecting applications. For disinfecting previously cleaned non-porous surfaces, mix 1 ounce SAN-O-FEC-50 in 10 gallons of water. For disinfecting previously cleaned porous surfaces, mix 1 ounce SAN-O-FEC-50 in 5 gallons of water.

In the Poultry Industry—For dipping hatching eggs, rinsing egg trays, fountains, feeders, brooders, battery egg crates, coops, spraying or washing incubators, walls and floors. Mix 1 ounce SAN-O-FEC-50 in 20 gallons of water to prepare the sanitizing solution. Use double strength (1 ounce in 10 gallons of water) if infection was present in previous flock. Hatchability is not affected when eggs are treated with SAN-O-FEC-50. Do not use for dipping of eggs for human consumption. **See Application Procedures for proper use of SAN-O-FEC-50.**

In the Swine Industry—For disinfecting farrowing pens, rearing pens, feeders, and waterers, mix 1 ounce SAN-O-FEC-50 in 10 gallons of water. **See Application Procedures for proper use of SAN-O-FEC-50.**

For Pets—For disinfecting and deodorizing cages, kennels, feeding pans, etc., mix 1 ounce SAN-O-FEC-50 in 20 gallons of water.

APPLICATION PROCEDURES

For proper use of SAN-O-FEC-50 in poultry and livestock premises, observe the following:

1. Remove all animals, poultry, and feed from premises, wash troughs, tracks, and other equipment.
2. Remove all litter and manure from floors, walls and surfaces of battery pens, stalls, hatcher and other facilities and fixtures occupied or traversed by animals or poultry.
3. Empty all troughs, tanks, and other feeding and watering equipment.
4. Saturate all surfaces with the SAN-O-FEC-50 solution.
5. To remove water, soap, and other types of equipment used in feed preparation, thoroughly wash troughs, tanks, walls and troughers, the same way.

INSTRUCTIONS FOR USE

cover hatching buildings, cages and other structures used to raise poultry chicks. For emptying equipment, until treated with solution, do not use for other

use. All treated feed racks, managers, troughs, waterers, feeders, fountains, and waterers must be thoroughly scrubbed with detergent and rinsed with potable water prior to reuse.

In packing plants and storage plants—For sanitizing containers in packing and storage plants, mix 1 ounce of SAN-O-FEC-50 in 20 gallons of water and apply. Flush and rinse with clear water. On surfaces that will be lean in reuse strength of sanitizing solution to 1 ounce SAN-O-FEC-50 in 5 gallons of water. Flush surfaces with a potable water rinse before reuse. For mangle walls and floors, mix 1 ounce SAN-O-FEC-50 in 5 gallons of water.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME ENVIRONMENT STORE ABOVE FREEZING TEMPERATURES

DANGER: Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of feet.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash or laundered clothing before reuse.

If swallowed, promptly drink a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

Rinse empty container thoroughly with water before disposal.

E.O.I. NO. 1-125

QDE NO. 4965

Whitmoyer
WHITMOYER
LABORATORIES, INC.
A Division of
ROHM AND HAAS COMPANY



1967