

A CRESYLIC ACID PRODUCT FOR USE IN SWINE, BEEF CATTLE AND HORSE BUILDINGS


ACTIVE INGREDIENT:

Cresylic Acid 100%

EPA Reg. No. 984-37AA

DANGER: Keep out of reach of children. See first aid and other precautions on side panel.

NET
CONTENTS
5 LIQUID
GALLONS (U.S.)



Rinse sprayer equipment with soap and water immediately after using HYDROL CONCENTRATE solution. The oil may soften the hoses and other rubber parts if this is not done. It is preferable to use a sprayer equipped with synthetic rubber or similar material which will not be affected by oil.

DANGER — Corrosive: Causes eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. May be fatal if swallowed or absorbed through skin. May produce severe burns. Keep away from children, domestic animals and foodstuffs. Avoid breathing the spray mist. Use with adequate ventilation. Avoid spilling or spraying HYDROL CONCENTRATE or any of its solution on animals, poultry or eggs. Wash hands thoroughly after handling. Avoid spilling or spraying on open flame.

FIRST AID — In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

Cleaning and feeding and watering equipment should be disinfected with a suitable disinfecting material.

Do not reuse empty drums or containers. Drums should be returned to reconditioner or destroyed by perforating or crushing and burying in a safe place.

Do not use around lactating dairy animals. Do not use in milking stalls, milking parlors or milk houses. Do not house livestock or place equipment until treatment has been absorbed, set or dried.

LOT NO.

CODE NO. 2536


Manufactured by

WHITMOYER


LABORATORIES, INC.

MYERSTOWN, PENNSYLVANIA 17067

SUBSIDIARY OF ROHM AND HAAS COMPANY



309147



Printed in U.S.A.

DISINFECTANT

Phenol Coefficient—6; A.O.A.C. Method

**A GENERAL DISINFECTANT FOR STABLES,
HOG PENS, POULTRY HOUSES, KENNELS
AND HOUSEHOLDS.**

ACTIVE INGREDIENTS:

Coal Tar Neutral Oil 60%
Soap 12%
Coal Tar Phenols 18%

INERT INGREDIENT:

Water 10%

DANGER: KEEP OUT OF THE REACH OF CHILDREN.

See other Precautionary statements on back panel.

EPA Reg. No. 984 - 41

NET CONTENTS: 1 LIQUID GALLON (U.S.)

MANUFACTURED FOR

BARKER, MOORE AND MEIN

Myerstown, Pa. 17067

"OVER A CENTURY OF SERVICE TO ANIMAL AGRICULTURE"

Division of Whitmoyer Laboratories, Inc.

BARKER'S

DISINFECTANT

A DISINFECTANT

FOR LIVESTOCK

PREMISES AND

HOUSEHOLD

ACCEPTED USE

3/28/73

UNDER THE FEDERAL INSECTICIDE
AND FUNGICIDE ACT
REGISTERED UNDER NO. 984-41

Comment

preparing smaller quantities
DISINFECTANT to each gallon of

Livestock Building Disinfectant
procedures. Mix 1 gallon of
of water.

1. Do not use in milking st
2. Remove all poultry and
treated. Destroy or ren
3. Remove all litter and n
barns, pens, stalls and c
by animals.
4. Empty all troughs, racks
5. Saturate all surfaces with
6. Thoroughly wet or imm
handling or restraining a
7. Ventilate buildings and c
stock or place equipment
8. All treated feed racks, n
waterers must be thorough
able water prior to reuse

Other Uses—For disinfectin
walls and woodwork of hon
filth and heavy soil deposits
BARKER'S DISINFECTANT to 2
BARKER'S DISINFECTANT

DANGER — May be fata
produce severe burns! Do
breathing spray mist. Wash
Keep away from food.

Avoid spilling or spraying c
or store near feed or foods
equipment or disposal of w
puncture and bury in a safe
areas. Do not use in milk
reuse empty container.

Lot No.

DISINFECTANT

Phenol Coefficient--6: A.O.A.C. Method

A GENERAL DISINFECTANT
FOR PENS, POULT
AND PIGS

FOR STABLE
AND KENNELS

ACTIVE INGREDIENTS

Coal Tar Neutral Oil	6.0%
Soap	1.0%
Coal Tar Phenol	1.0%

INERT INGREDIENT

Water

DANGER: KEEP OUT OF THE REACH OF CHILDREN

See other Precautionary statements on back panel.

EPA Reg. No. 105-111

NET CONTENTS: 1 LIQUID GALLON (U.S.)

MANUFACTURED FOR

BARKER, MOORE AND MEIN

Myerstown, Pa. 17067

"OVER A CENTURY OF SERVICE TO ANIMAL AGRICULTURE"

Division of Whitmoyer Laboratories, Inc.

BARKER'S

DISINFECTANT

A DISINFECTANT

FOR LIVESTOCK

PREMISES AND

HOUSEHOLD

USE

1. Add 1 ounce of BARKER'S DISINFECTANT to each gallon of

Livestock Building Disinfection procedures. Mix 1 gallon B of water.

1. Do not use on drinking water.
2. Remove all poultry and treat. Destroy or remove.
3. Remove all litter and in barns, pens, stalls and clean by animals.
4. Empty all troughs, racks.
5. Saturate all surfaces with.
6. Thoroughly wet on handling or restraining a.
7. Ventilate buildings and stock or place equipment.
8. All treated feed racks, waterers must be thoroughly clean water prior to reuse.

Other Uses For disinfecting walls and woodwork of hoards filthy and heavy soil deposits BARKER'S DISINFECTANT to 2 BARKER'S DISINFECTANT.

DANGER May be fatal produce severe burns! Do breathing spray must. Wash Keep away from food.

Avoid spilling or spraying on or store near food or foods equipment or disposal of waste and bury in a safe area. Do not use in milk rooms empty containers.

10/7/57

DISINFECTANT

A.C. Method

USE FOR STABLES,
BARNYARDS, KENNELS
AND CAGES.

- 60%
- 12%
- 18%
- 10%

KEEP AWAY FROM EACH OF CHILDREN.
See instructions on back panel.

- 41

1 GALLON (U.S.)

OR

BARKER'S DISINFECTANT

A DISINFECTANT

FOR LIVESTOCK

PREMISES AND

HOUSEHOLD

ACCEPTED USE
3/28/73
UNDER THE FEDERAL INSECTICIDE
AND RODENTICIDE ACT
EPA REGISTRATION NO. 987-41
Comment

For preparing smaller quantities of solution, add 1 cup of BARKER'S DISINFECTANT to each gallon of water.

Livestock Building Disinfectant — BARKER'S DISINFECTANT application procedures. Mix 1 gallon BARKER'S DISINFECTANT with each 29 gallons of water.

1. Do not use in milking stalls, milking parlors, milk houses or dairy barns.
2. Remove all poultry and other livestock from the building or area to be treated. Destroy or remove animal feeds.
3. Remove all litter and manure from floors, walls and other surfaces of barns, pens, stalls and chutes and any other equipment occupied or used by animals.
4. Empty all troughs, racks and other feeding and watering equipment.
5. Saturate all surfaces with the BARKER'S DISINFECTANT solution.
6. Thoroughly wet or immerse halters, ropes and other equipment used in handling or restraining animals as well as forks, shovels and scrapers.
7. Ventilate buildings and other closed spaces. Do not house poultry or livestock or place equipment until treatment has been absorbed, set or dried.
8. All treated feed racks, mangers, troughs, automatic feeders, fountains and waterers must be thoroughly scrubbed with detergent and rinsed with potable water prior to reuse.

Other Uses—For disinfecting and deodorizing sinks, garbage pails and floors, walls and woodwork of homes, public buildings and factories. Remove gross filth and heavy soil deposits and then scrub with a solution of 1 part BARKER'S DISINFECTANT to 29 parts water. Thoroughly wet all surfaces with BARKER'S DISINFECTANT.

DANGER — May be fatal if swallowed or absorbed through skin! May produce severe burns! Do not get in eyes, on skin, or on clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Keep away from food.

Avoid spilling or spraying on animals, poultry or eggs. Do not contaminate or store near feed or foodstuffs. Do not contaminate water by cleaning of equipment or disposal of wastes. Rinse disinfectant container and crush or puncture and bury in a safe place. Do not allow livestock to graze treated areas. Do not use in milk rooms, milking parlors or dairy barns. Do not reuse empty container.

Lot No.

Code No. 1652