

AFFILIATED

STERI-FAC

SANITIZER AND DISINFECTANT

A cleanser, disinfectant and deodorant for kennels, cages and runs, animal hospital floors and walls, surgical instruments and other equipment.

ACTIVE INGREDIENTS:

n-alkyl (50% C ₁₄ , 40% C ₁₂ , 10% C ₁₆)	
dimethyl benzyl ammonium chloride	5.00%
Sodium hydroxide	3.00%
Tetrasodium ethylenediamine tetraacetate	0.88%

INERT INGREDIENTS 91.12%

Detergents, Sequestering Agent and Buffers included

E.P.A. Reg. No. 984-55AA

DANGER: Keep out of reach of children. See First Aid and other precautionary statements on side panel.

NET CONTENTS
1 LIQUID QUART (U.S.)

Manufactured by

AFFILIATED
LABORATORIES DIVISION
WHITMOYER LABORATORIES, INC.
MYERSTOWN, PENNSYLVANIA 17067
A DIVISION OF ROHM AND HAAS COMPANY



ACCEPTED
Mar. 1, 1973
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 984-55

DIRECTIONS FOR USE

DILUTION RECOMMENDATIONS — Use ½ ounce (15 cc) of STERI-FAC for each gallon of water for cleaning and disinfecting non-porous surfaces such as smooth metal or plastic table and cabinet tops. For porous surfaces such as wood or concrete, which are difficult to clean or disinfect, use 1 ounce of STERI-FAC for each gallon of water. Hot or cold water may be used.

The excellent cleaning properties of STERI-FAC are achieved through the detergent and wetting agent formulation. Detergents used in STERI-FAC are biodegradable.

GENERAL CLEANING AND DISINFECTING — Clean floors, walls, runs, cages and contaminated equipment with 1 ounce of STERI-FAC in each gallon of warm water. Brush, sponge or mop until surfaces are clean. Rinse surfaces and thoroughly wet with a fresh solution of ½ ounce STERI-FAC in each gallon of water. Use 1 ounce STERI-FAC per gallon of water on wood, concrete or other porous surfaces. Heavily soiled areas should be reasonably free of gross organic soil before attempting to disinfect.

GLASSWARE, FEED AND WATER UTENSILS — Soak in solution of 1 ounce of STERI-FAC per gallon of water to loosen adhering soil. If necessary scrub or brush to remove soil. Rinse with clean water and soak for 10 minutes in a fresh solution of ½ ounce STERI-FAC per gallon of water to sanitize. Rinse in fresh water before using.

SURGICAL INSTRUMENTS — Soiled instruments should be soaked in a solution of ½ ounce STERI-FAC to ½ gallon of water. Scrub or brush to remove adhering debris. To achieve disinfection soak in

a solution of the same strength for 10 minutes. Rinse with clean water after cleaning and before autoclaving or dry heat sterilization. STERI-FAC is not recommended for cold sterilization of surgical instruments.

EXAMINATION TABLES — After examining animals, sponge examination table tops with a solution of 1 ounce STERI-FAC per gallon of water to remove animal hair and debris. To disinfect, thoroughly wet surfaces with the same strength STERI-FAC solution and allow 10 minute contact.

DANGER — Keep Out of Reach of Children. Undiluted product is corrosive and causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID — In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN — Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water before disposal.

LOT NO.

FOR VETERINARY USE ONLY

301047

Printed in U.S.A.