

# WHITMOYER



## An emulsified cresylic acid disinfectant

### ACTIVE INGREDIENTS

Cresylic acid ..... 50%

Soap ..... 38%

INERT INGREDIENTS ..... 12%

EPA Reg. No. 984-42 AA

**WARNING:** Keep out of reach of children.  
See other warning statements on back panel.

NET CONTENTS 5 LIQUID GALLONS (U.S.)

PANTEK II is a cresylic acid disinfectant which is diluted with water and is used for disinfecting poultry houses, swine farrowing and finishing houses and other livestock buildings. PANTEK II water solution is particularly effective for the control of bacterial organisms and pathogenic fungi (*T. interdigitale* by A.O.A.C. Fungicidal Test) on inanimate surfaces.

### DIRECTIONS FOR USE

Mix 1 part PANTEK II with 200 parts of water.

Do not use in dairy barns, milking stalls, milking parlors or milk houses.

#### WALLS, CEILINGS, FLOORS

**Poultry** -- Remove all poultry, moveable equipment, feed, litter and waste from the house and clean thoroughly. Empty all remaining troughs and other feeding and watering equipment before applying PANTEK II. Disinfect the cleaned house with PANTEK II using 20 to 40 gallons of solution per 1000 square feet of floor space. All corners, cracks, crevices and ledges must be thoroughly wet for best results. High pressure, high volume spray equipment is useful in getting PANTEK II to the areas to be disinfected. Clean and immerse all accessory equipment such as coops, forks, shovels and scrapers. Ventilate the building and do not place poultry or use equipment until the PANTEK II solution has been absorbed, set or dried.

**Swine and Other Livestock** -- Do not use in dairy barns, milking stalls, milking parlors or milk houses. Remove all animals, moveable equipment, feed, litter and waste from the house. Empty all troughs, racks and other feeding and watering equipment. Scrape and scrub to remove all adhering dirt and manure. Disinfect the cleaned house with PANTEK II using 20 to 40 gallons of solution per 1000 sq. ft. of floor space, preferably using high pressure, high volume application to saturate all surfaces with the disinfecting solution.

Clean and immerse all accessory equipment such as halters, ropes, forks, shovels and scrapers. Ventilate the building and do not place livestock or use equipment until the PANTEK II solution has been absorbed, set or dried.


**Equipment, Feeders, Waterers, Brooders, Hatchery Equipment** Mix 1 part PANTEK II to 200 parts of water (approximately 2 ounces PANTEK II per 3 gallons of water). Empty all troughs, racks and other feeding and watering equipment. Scrub equipment and thoroughly wet all surfaces with this solution. All treated feed racks, mangers, troughs, automatic feeders, fountains and waterers must be thoroughly scrubbed with detergent and rinsed with potable water prior to reuse.

**SHOE BATH SANITIZER** -- Add 1 ounce of PANTEK II to each gallon of water. Place the solution in shallow pans at all entrances to poultry and livestock buildings. Wash sanitizer pans and add fresh PANTEK II solution daily. Routine use of shoe baths prepared with PANTEK II will aid in reducing the spread of disease organisms between poultry houses and other livestock buildings.

**WARNING:** May be fatal if swallowed or absorbed through skin! May produce severe burns! Do not get in eyes, on skin or on clothing! Avoid breathing spray mist. Wash thoroughly after handling. Internal Antidote -- Use demulcents of egg white, milk or gruel. Seek medical help immediately.

Avoid spilling or spraying PANTEK II or any of its solution on animals, poultry or eggs. Do not use on nests or nesting material. Do not contaminate or store near feed or foodstuffs. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not allow livestock to graze on treated areas. Do not use in dairy barns, milking stalls, milking parlors or milk houses. Rinse empty container thoroughly with water before disposal.

LOT NO.



CODE NO. ~~XXX~~  
4026



Manufactured by  
**WHITMOYER**  
LABORATORIES, INC.  
MERTZTOWN, PENNSYLVANIA 19380  
A ROHM AND HAAS COMPANY



# WHITMOYER

# LARO™

**BACTERICIDE**

**Phenolic disinfectant and deodorizer**

**ACTIVE INGREDIENTS:**

Soap 10%; Isopropanol 10%; 4- & 6-chloro-2-phenylphenol 7%;  
o-phenylphenol 6%; tetrasodium ethylenediamine tetraacetate  
1.5%; p-tertiary amyphenol 1.0% ..... 35.5%

**INERT INGREDIENTS:** ..... 64.5%


Water, sodium hydroxide, potassium sulfite

EPA Reg. No. 984-57      EPA Est. 984-PA-1

**DANGER:** Keep Out of Reach of Children. See First Aid and other precautions on side panel.

**NET CONTENTS 1 GALLON (U.S.)**

**FUNGICIDE**



**DIRECTIONS FOR USE**

Laro is concentrated and must be diluted before use. Mix 1 part Laro with 200 parts water and use for all disinfecting purposes in hatcheries, poultry and livestock buildings, kennels and veterinary hospitals. Also for disinfecting equipment and vehicles and sanitizing hatching eggs.

Laro has been tested and shown to be bactericidal and fungicidal by official methods. At a use dilution of 1 part Laro to 200 parts water, the pathogenic fungi, *Trichophyton interdigitale*, *Aspergillus niger* and *Candida albicans* were killed on inanimate surfaces when subjected to the A.O.A.C. Fungicidal Test.

**Poultry and Livestock Buildings** — Use Laro for disinfecting all types of poultry houses, swine, cattle, sheep and horse buildings, stalls, all equipment and vehicles. The following procedure is recommended for disinfecting facilities for food producing animals. Do not use in milking stalls, milking parlors or milk houses.

1. Remove all animals, poultry and feed from premises, coops and crates.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities occupied or traversed by animals.
3. Empty all troughs, racks and other feeding and watering equipment.
4. Saturate all surfaces with the Laro solution.
5. Immerse all halters, ropes and other equipment used in handling and restraining animals. Treat forks, shovels and scrapers the same way.
6. Ventilate buildings, coops and other closed spaces. Do not house poultry or livestock or employ equipment until treatment has been absorbed, set or dried.
7. All treated feed racks, mangers, troughs, automatic feeders, fountains and waterers must be thoroughly scrubbed and rinsed with potable water prior to reuse.

**Veterinary Hospitals, Kennels and Animal Shelters** — Clean runs, floors, walls, cages and equipment thoroughly and disinfect with a 1 to 200 Laro solution.

**Hatcheries** — Thoroughly clean incubators, hatchers, equipment, floors and walls and apply Laro to completely wet all surfaces. Also disinfect setter and hatcher trays before reusing.

Sanitizi  
of Laro

Shoe Ba  
at all e  
solution  
spread

**DANGER**  
irritation  
rubber  
through

**FIRST A**  
at least  
clothing

If swallo  
if these  
physicia

**NOTE T**  
gastric  
convuls

Rinse e  
DO NOT

CODE 2