

WHITMOYER

SAN-O-FEC-5

Disinfectant, deodorant and sanitizer

ACTIVE INGREDIENTS
Didecyl dimethyl ammonium chloride 37.5%
INERT INGREDIENTS 62.5%

EPA Reg. No. 984-71 EPA EST. 984-PA-1

KEEP OUT OF REACH OF CHILDREN

DANGER

See first aid statement and other precautions on back panel.

NET CONTENTS 1 GALLON (U.S.)

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

DANGER: Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID — In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, promptly drink a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT
IN A MANNER INCONSISTENT WITH ITS LABELING**

PREMISE DISINFECTANT AND SANITATION OF FLOORS AND WALLS — Mix 1 ounce SAN-O-FEC-5 in 20 gallons of water for usual sanitizing applications. For disinfecting previously cleaned non-porous surfaces, mix 1 ounce SAN-O-FEC-5 in 10 gallons of water. For disinfecting previously cleaned porous surfaces mix 1 ounce SAN-O-FEC-5 in 5 gallons of water.

POULTRY INDUSTRY — For dipping hatching eggs, rinsing egg trays, fountains, feeders, brooders, batteries, egg crates, coops, spraying or washing incubators, walls and floors. Mix 1 ounce SAN-O-FEC-5 in 20 gallons of water to prepare the sanitizing solution. Use double strength (1 ounce in 10 gallons of water) if infection was present in previous flocks. Hatchability is not affected when eggs are treated with SAN-O-FEC-5. Do not use for dipping of eggs for human consumption.

SWINE INDUSTRY — For disinfecting farrowing pens, rearing pens, feeders and waterers, mix 1 ounce SAN-O-FEC-5 in 10 gallons of water. See Application Procedures for proper use of SAN-O-FEC-5.



61

BEST DOCUMENT AVAILABLE

ADVERSE STATEMENTS TO HUMANS AND DOMESTIC ANIMALS

Causes severe eye and skin damage. Do not wear goggles or face shield and rubber gloves swallowed. Avoid contamination of food. Immediately flush eyes or skin with plenty of water and call a physician. Remove and wash contaminated

quantity of milk, egg whites, gelatin solution or large quantities of water. Avoid alcohol. Call a

musical damage may contraindicate the use of circulatory shock, respiratory depression and

CHEMICAL HAZARDS

Non-flammable.

INSTRUCTIONS FOR USE

GENERAL LAW TO USE THIS PRODUCT

CONSISTENT WITH ITS LABELING

SANITIZATION OF FLOORS AND WALLS — Mix 1 ounce of water for usual sanitizing applications. For non-porous surfaces, mix 1 ounce SAN-O-FEC-5 in 10 gallons of water. For previously cleaned porous surfaces mix 1 ounce in 5 gallons of water.

For hatching eggs, rinsing egg trays, fountains, troughs, coops, spraying or washing incubators, mix 1 ounce SAN-O-FEC-5 in 20 gallons of water to prepare the solution (1 ounce in 10 gallons of water) if infection susceptibility is not affected when eggs are treated. For dipping of eggs for human consumption.

For farrowing pens, rearing pens, feeders and troughs, mix 1 ounce SAN-O-FEC-5 in 10 gallons of water. See Application Procedures for proper use of SAN-O-FEC-5.

PETS — For disinfecting and deodorizing cages, kennels, feeding pans and other equipment, mix 1 ounce SAN-O-FEC-5 in 10 gallons of water. See Application Procedures for proper use of SAN-O-FEC-5.

FOOD PACKING AND STORAGE PLANTS — For sanitizing counters in food packing and storage plants, mix 1 ounce of SAN-O-FEC-5 in 20 gallons of water (150 ppm active) and apply. Flush and rinse with clear water. On surfaces difficult to clean, increase strength of sanitizing solution to 1 ounce SAN-O-FEC-5 in 5 gallons of water. Flush surfaces with a potable water rinse before reuse. For rinsing walls and floors, mix 1 ounce SAN-O-FEC-5 in 5 gallons of water. For sanitizing there should be at least one minute contact time.

DAIRY COWS — The udders, flanks and teats of dairy cows can be sanitized by washing with a solution of 1/2 oz. SAN-O-FEC-5 in 10 gallons of warm water. No potable water rinse is required. Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt or soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

MILK CONTAINERS AND UTENSILS — At 1/2 oz./10 gallons, SAN-O-FEC-5 fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinances 1978. Recommendations of the U.S. Public Health Services in waters up to 1000 ppm of hardness calculated as CaCO₃ when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against Escherichia coli and Staphylococcus aureus.

APPLICATION PROCEDURES

Clean all surfaces and thoroughly rinse away soap or detergent before applying SAN-O-FEC-5 solution. Do not mix with soap or synthetic detergents. Apply by cloth, sponge, mop or mechanical spray. Prepare fresh solution daily or when use solution becomes visibly dirty.

For proper use of SAN-O-FEC-5 in poultry and livestock premises, observe the following:

1. Remove all animals, poultry, feed and other equipment from premises.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals and poultry.
3. Empty all troughs, racks and other feeding and watering equipment.
4. Saturate all surfaces with SAN-O-FEC-5 solution for a period of ten minutes.
5. Immerse all halters, ropes and other types of equipment used in handling and restraining animals. Treat forks, shovels and scrapers the same way.

6. Ventilate buildings, coops and other closed spaces. Do not house poultry or livestock or employ equipment until treatment has been absorbed, set or dried.
7. Thoroughly scrub treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and thoroughly rinse with potable water prior to reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL

Pesticide or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL

Triple rinse (or equivalent) and dispose in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

GENERAL

Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.


LOT NO

Manufactured by
WHITMOYER
LABORATORIES, INC.
MYERSTOWN, PENNSYLVANIA 17067

CODE 4966-011-1001
E80

Printed in USA

62



BEST DOCUMENT AVAILABLE