

WHITMOYER

SAN-O-FEC[®] 2

Disinfectant, deodorant and sanitizer

ACTIVE INGREDIENTS
Didecyl dimethyl ammonium chloride 15.0%
INERT INGREDIENTS 85.0%

EPA Reg. No. 984-72 EPA EST 984-PA-1

KEEP OUT OF REACH OF CHILDREN

63

DANGER

See first aid statement and other precautions on back panel.

NET CONTENTS 1 GALLON (U.S.)

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

DANGER: Corrosive. Causes severe eye and skin contact. Irritant. Causes severe eye and skin contact if get in eyes, on skin or on clothing. Wear goggles or face shield and when handling. Harmful or fatal if swallowed. Avoid contamination of food and feed.
FIRST AID — In case of contact, immediately flush eyes or skin with water for at least 15 minutes. For eyes, call a physician. Remove and wash clothing before reuse.

If swallowed, promptly drink a large quantity of milk, egg whites, gelatin or water if these are not available, drink large quantities of water. Avoid physical exertion.

NOTE TO PHYSICIAN: Probably mucosal damage may contraindicate gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.



PHYSICAL OR CHEMICAL HAZARD

Do not use or store near heat or open flame.

**DIRECTIONS FOR USE
GENERAL CLASSIFICATION**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

PREMISE DISINFECTION AND SANITATION OF FLOORS AND WALLS — Mix 1 ounce SAN-O-FEC-2 in 8 gallons of water for usual sanitizing and disinfecting previously cleaned non-porous surfaces; mix 1 ounce in 4 gallons of water. For disinfecting previously cleaned porous surfaces, mix 1 ounce SAN-O-FEC-2 in 2 gallons of water.

POULTRY INDUSTRY — For dipping hatching eggs, rinsing egg feeders, brooders, batteries, egg crates, coops, spraying or washing walls and floors. Mix 1 ounce SAN-O-FEC-2 in 8 gallons of water for sanitizing solution. Use double strength (1 ounce in 4 gallons of water) for sanitizing previous flocks. Hatchability is not affected when used with SAN-O-FEC-2. Do not use for dipping of eggs for human consumption.

SWINE INDUSTRY — For disinfecting farrowing pens, rearing piglets, mix 1 ounce SAN-O-FEC-2 in 4 gallons of water. Procedures for proper use of SAN-O-FEC-2.

ACCEPTED
NOV 19 1980
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 984-72

CAUTIONARY STATEMENTS REGARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive. Causes severe eye and skin damage. Do not breathe fumes. Wear goggles or face shield and rubber gloves. Fatal if swallowed. Avoid contamination of food.

In case of contact, immediately flush eyes or skin with plenty of water. In case of ingestion, call a physician. Remove and wash contaminated clothing.

Avoid contact with large quantity of milk, egg whites, gelatin solution or other large quantities of water. Avoid alcohol. Call a physician.

Severe mucosal damage may contraindicate the use of this product in case of circulatory shock, respiratory depression and other conditions.

FLAMMABLE OR CHEMICAL HAZARDS

Flammable or open flame.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

FEDERAL LAW TO USE THIS PRODUCT

CONSISTENT WITH ITS LABELING

DISINFECTION OF FLOORS AND WALLS — Mix 1 ounce of SAN-O-FEC-2 in 4 gallons of water for usual sanitizing applications. For non-porous surfaces, mix 1 ounce SAN-O-FEC-2 in 2 gallons of water. For sanitizing previously cleaned porous surfaces mix 1 ounce of SAN-O-FEC-2 in 4 gallons of water.

For dipping hatching eggs, rinsing egg trays, fountains, troughs, coops, spraying or washing incubators, mix 1 ounce SAN-O-FEC-2 in 8 gallons of water to prepare the solution. For dipping of eggs for human consumption, mix 1 ounce in 4 gallons of water if infection is suspected. Hatchability is not affected when eggs are treated with this solution.

For dipping farrowing pens, rearing pens, feeders and troughs, mix 1 ounce SAN-O-FEC-2 in 4 gallons of water. See Application Procedures for proper use of SAN-O-FEC-2.

PETS — For disinfecting and deodorizing cages, kennels, feeding pans and other equipment, mix 1 ounce SAN-O-FEC-2 in 4 gallons of water. See Application Procedures for proper use of SAN-O-FEC-2.

FOOD PACKING AND STORAGE PLANTS — For sanitizing counters in food packing and storage plants, mix 1 ounce of SAN-O-FEC-2 in 8 gallons of water (150 ppm active) and apply. Flush and rinse with clear water. On surfaces difficult to clean, increase strength of sanitizing solution to 1 ounce SAN-O-FEC-2 in 2 gallons of water. Flush surfaces with a potable water rinse before reuse. For rinsing walls and floors, mix 1 ounce SAN-O-FEC-2 in 2 gallons of water. For sanitizing there should be at least one minute contact time.

DAIRY COWS — The udders, flanks and teats of dairy cows can be sanitized by washing with a solution of 1/2 oz. SAN-O-FEC-2 in 4 gallons of warm water. No potable water rinse is required. Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

MILK CONTAINERS AND UTENSILS — At 1/2 oz./4 gallons, SAN-O-FEC-2 fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinance 1978 Recommendations of the U.S. Public Health Services in waters up to 1000 ppm of hardness calculated as CaCO₃ when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against Escherichia coli and Staphylococcus aureus.

APPLICATION PROCEDURES

Clean all surfaces and thoroughly rinse away soap or detergent before applying SAN-O-FEC-2 solution. Do not mix with soap or synthetic detergents. Apply by cloth, sponge, mop or mechanical spray. Prepare fresh solution daily or when use solution becomes visibly dirty.

For proper use of SAN-O-FEC-2 in poultry and livestock premises, observe the following:

1. Remove all animals, poultry, feed and other equipment from premises.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals and poultry.
3. Empty all troughs, racks and other feeding and watering equipment.
4. Saturate all surfaces with SAN-O-FEC-2 solution for a period of ten minutes.
5. Immerse all halters, ropes and other types of equipment used in handling and restraining animals. Treat forks, shovels and scrapers the same way.

6. Ventilate buildings, coops and other closed spaces. Do not house poultry or livestock or employ equipment until treatment has been absorbed, set or dried.
7. Thoroughly scrub treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and thoroughly rinse with potable water prior to reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL

Pesticide or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL

Triple rinse (or equivalent) and dispose in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

GENERAL

Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

LOT NO.

Manufactured by
WHITMOYER
LABORATORIES, INC.
MYERTOWN, PENNSYLVANIA 17067

CODE 4-7-011-1001
FR1

Printed in USA

BEST DOCUMENT AVAILABLE