

Form 100-100

ACCEPTED
APR 19 1957
1957-37

AGRICULTURE
UNITED STATES DEPARTMENT OF AGRICULTURE

UNITED STATES ANIMAL AND PLANT HEALTH QUARANTINE
DISEASES OF ANIMALS, AND PLANTS, AND PRODUCTS THEREOF
APPLICATION

ACTIVE INGREDIENTS

1. Dichlorodiphenyl dimethyl	
phosphate*	10.00
2. Diethyl p-aminobenzoate*	10.00
3. Hydroxyquinoline*	10.00
	<u>30.00</u>

Equivalent to 100% (w/w) insecticide
Total weight _____

Directions for use: ...

Preparation: ...

Other information: ...

Signature
Date

Printed name
Address

USE INSTRUCTIONS

Use in conventional hand, or automatic spraying equipment, for control of insects on animals.

For the control of certain insects on animals, apply in accordance with best commercial recommendations. Do not apply directly to animal unless advised by label. Do not use on animals with known hypersensitivity to organic phosphates.

USE AND DIRECTIONS FOR APPLICATION

Section 1: Use on Cattle (When used on cattle, use the following directions.)

Apply 1 to 2 fluid ounces of 4-4 solution per animal on a 100 lb. animal with hand or automatic sprayer. Spray to thoroughly cover the entire animal, including the legs, with the solution. Do not spray in excess of 2 fluid ounces per 100 lbs. of animal weight. Do not use on animals with known hypersensitivity to organic phosphates.

Section 2: Use on Horses (When used on horses, use the following directions.)

Apply 1 to 2 fluid ounces of 4-4 solution per animal on a 1000 lb. animal with hand or automatic sprayer. Spray to thoroughly cover the entire animal, including the legs, with the solution. Do not spray in excess of 2 fluid ounces per 100 lbs. of animal weight. Do not use on animals with known hypersensitivity to organic phosphates.

USE AND DIRECTIONS FOR APPLICATION

Apply 1 to 2 fluid ounces of 4-4 solution per animal on a 1000 lb. animal with hand or automatic sprayer. Spray to thoroughly cover the entire animal, including the legs, with the solution. Do not spray in excess of 2 fluid ounces per 100 lbs. of animal weight. Do not use on animals with known hypersensitivity to organic phosphates.

Apply 1 to 2 fluid ounces of 4-4 solution per animal on a 1000 lb. animal with hand or automatic sprayer. Spray to thoroughly cover the entire animal, including the legs, with the solution. Do not spray in excess of 2 fluid ounces per 100 lbs. of animal weight. Do not use on animals with known hypersensitivity to organic phosphates.