RATOREX®

Pir. 14

PRICE

ACCEPTED

SEP 1 / 1979

Under the Federal Insection of Fungicide, and Rodenticide Let. as amended, for the posticide registered under 99.

with PROLIN Kills Rats and Mice

with PROE

KILLS NORWAY RATS, ROOF RATS & HOUSE MICE

ACTIVE INGREDIENTS:

Warfarin 3-(alpha-acetonylbenzyl) -4-hydroxycoumarin _____ N'-(2-quinoxalinyl) sulfanilamide

(Sulfaquinoxaline) 0.025% INERT INGREDIENTS _____

EPA Reg. No. 995-41 EPA Est. No. 995-MN-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR REQUIRED PRECAUTIONS

NET WEIGHT 41/2 POUNDS

ANOTHER QUALITY PRODUCT BY

WINONA, MINNESOTA 55987

PRECAUTIONARY STATEMENTS:

HAZARD TO HUMANS & DOMESTIC ANIMALS:

CAUTION: Keep away from humans, domestic animals and pets. If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K1 combined with blood transfusions are indicated, as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

ENVIRONMENTAL HAZARD:

Keep Out of Lakes, Streams or Ponds.

GUARANTEE ...

RATOREX is guaranteed to give complete satisfacion when used as directed if RATOREX fails to perform as guaranteed, mail the empty carron for replacement to the ment of the satisfactors.

Language . The wife of the state of the same of the same .

with PROLIN

BEST AVAILABLE COPY



PELLETED RAT AND MICE BAIT



RATOREX

with PROLIN

RATS COST YOU MONEY-KILL 'EM!

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS — For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar manmade structures.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS — After removing as much food, water, and harborage as possible, determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow.

APPLICATION DIRECTIONS


RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 1/4 to 1/2 ounce (1-2 level tablespoons) of bait at 8 to 1-2 foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days.


MATS & MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. Repeat treatment when intestation-recurs. Where continuous source of infestation is present, establish permanent bait stations and replanish bait as needed.

DISPOSAL

DO NOT REUSE EMPTY CONTAINER.
WRAP CONTAINER AND PUT IN TRASH COLLECTION.









Kills Rats and Mice

WIEW with PROLIN

BEST AVAILABLE COPY