

pm14

995-41

189

ACCEPTED

OCT 10 1990

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 995-41

RATOREX®

with **PROLIN**

Kills Rats and Mice

PRECAUTIONARY STATEMENTS:
HAZARD TO HUMANS & DOMESTIC ANIMALS
CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K, intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD: Keep Out of Lakes, Streams or Ponds.

RATOREX®

with **PROLIN**

KILLS NORWAY RATS, ROOF RATS & HOUSE MICE



GARANTEE

RATOREX is guaranteed to give complete satisfaction when used as directed. If RATOREX fails to perform as guaranteed, mail the empty carton for replacement.

ACTIVE INGREDIENTS:	
Warfarin 3-(alpha-acetonylbenzyl)	0.025%
4-hydroxycoumarin	99.975%
INERT INGREDIENT*	
*N-(2-quinoxalanyl) sulfanilamide (Sulfaquinoxaline)	0.025%

EPA Reg. No. 995-41
EPA Est. No. 995-MN-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR REQUIRED PRECAUTIONS

NET WEIGHT
1 POUND



MANUFACTURED BY
The MACKWIN Company
WINONA, MINNESOTA 55987

RATOREX®

with **PROLIN**

PELLETED RAT AND MICE BAIT

RATORREX[®]
with **PROLIN**
Kills Rats and Mice



Place this box in locations not accessible to children, pets and domestic animals. Do not use this box as a bait dispenser in areas accessible to children or non target animals.

RATORREX[®]

with **PROLIN**

RATS COST YOU MONEY—KILL 'EM!

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS — For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS — After removing as much food as possible, determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow.

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply ¼ to ½ ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS AND MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL

STORAGE: Store only in original container in a dry place inaccessible to children and pets.

DISPOSAL: Do not reuse empty container. Securely wrap in newspaper and discard in trash.

TAKE TIME



OBSERVE LABEL
DIRECTIONS

RATORIEX®

with **PROLIN**

RATS COST YOU MONEY—KILL 'EM!

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS — For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS — After removing as much food as possible, determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow.

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

Apply ¼ to ½ ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS AND MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

RATORIEX

KILLS NORWAY RATS, ROOF RATS & HOUSE MICE

ACTIVE INGREDIENTS:



Warfarin 3-(alpha-acetylbenzyl)	
-4-hydroxycoumarin.....	0.025%
INERT INGREDIENT*.....	99.975%
*N'-(2-quinoxaliny) sulfanilamide (Sulfaquinoxaline).....	0.025%

EPA Reg. No. 995-41
EPA Est. No. 995-MN-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR REQUIRED PRECAUTIONS



1700
PA



ACCEPTED
OCT 10 1990
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 995-41

STORAGE AND DISPOSAL

STORAGE: Store only in original container in a dry place inaccessible to children and pets.

DISPOSAL: Do not reuse empty container. Securely wrap in newspaper and discard in trash.

TAKE TIME



**OBSERVE LABEL
DIRECTIONS**