

PM 14

995-41

144

[top flap]

RATOREX

Kills Rats and Mice

[front label

RATOREX

KILLS NORWAY RATS,
ROOF RATS & HOUSE MICE

ACCEPTED
OCT 14 1992
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 995-41

ACTIVE INGREDIENTS:

Warfarin [3-(alpha-acetonybenzyl)-
4-hydroxycoumarin]..... 0.025Z

INERT INGREDIENTS.. 99.975Z

TOTAL 100.000Z

EPA Reg. No. 995-41
EPA Est. 995-MN-1

KEEP OUT OF REACH OF CHILDREN [12 pt]

CAUTION [18 pt]

SEE RIGHT PANEL FOR ADDITIONAL PRECAUTIONARY
STATEMENTS [12 pt]

NET WEIGHT
1 POUND

MANUFACTURED BY

THE MACKWIN COMPANY
WINONA, MINNESOTA 55987

[bottom flap]

RATOREX

PELLETED RAT AND MOUSE BAIT

249

[left side panel]

**RATOREX
KILLS RATS AND MICE**

GUARANTEE

RATOREX is guaranteed to give complete satisfaction when used as directed. If RATOREX fails to perform as guaranteed, mail proof of purchase for replacement. Do not mail empty carton.

TAKE TIME
OBSERVE LABEL
DIRECTIONS

[right side panel]

PRECAUTIONARY STATEMENTS:

**HAZARD TO HUMANS &
DOMESTIC ANIMALS**

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K1, intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times

ENVIRONMENTAL HAZARD:

This product is toxic to mammals and birds. Do not apply this product to water, swamps, bogs, or potholes. Keep out of lakes, streams, or ponds.

[back panel]

**RATOREX
RATS COST YOU MONEY - KILL 'EM!**

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife or in tamper-resistant bait stations. Bait stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. Units must be secured or immobilized to prevent bait from being shaken from the stations. Stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of container, and unused, spoiled, or unconsumed bait as specified on this label.

USE RESTRICTIONS - For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures. Do not place bait in areas where there is a possibility of contaminating food or food surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS - After removing as much alternate food as possible, determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow.

4-7-1

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 1/4 to 1/2 ounces (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Large placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS AND MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL

STORAGE: Store only in original container in a dry place inaccessible to children and pets.

DISPOSAL: Do not reuse empty container. Securely wrap in newspaper and discard in trash.