

SUPERIOR CHEMICAL PRODUCTS, INC.
3942 Frankford Ave.

BRAND NAME SUPRO

S U P R O
(Total Release Aerosol Can)

99-67

Kills Roaches, Fleas, Ticks, Silverfish, Flies, Mosquitoes, Gnats.
IN HOMES AND BUILDINGS (Inches, Industrial Plant, Warehouse, etc.)

DIRECTIONS: To be used by licensed Pest Control Operators Only. Use ^{five} one 6 oz. cans for each unobstructed closed area of 10,000 cu. ft. but not less than 7,000 cu. ft. (One 6 oz. can per 3,000 cu. ft.) During fogging, and to release contents, use a mask or respirator of a type passed by the U.S. Dep't of Agric for DDVP insecticide protection. Open cabinets and doors to areas to be treated. Shut off air conditioning and fans before using. All occupants must be removed from house before treatment is started, and should not reoccupy building for at least two hours. Open doors and windows and ventilate well before occupying. Unwrapped or unpackage food should be removed or placed in refrigerator before treatment. Food handling equipment and dishes should be washed after application. Do not contaminate water, feed or foodstuffs. Remove all pets. Cover or remove fish tanks. Cover house plants within 8 ft. of point of release, with newspaper to prevent burning.

Place aerosol on table in center of area to be treated, or near point of greatest infestation. Place newspaper under canister. When using more than one canister, place to give best distribution of vapors where free flow of mist is not assured.

Snap clip over lip of container, which will release entire contents automatically, and leave building at once.

DANGER: KEEP OUT OF REACH OF CHILDREN
SEE ANTIDOTE STATEMENT AND OTHER CAUTIONS ON BACK PANEL.

ACTIVE INGREDIENTS

2,2-dichlorovinyl dimethyl phosphate	6.5 %
Petroleum distillates	4.0 %
Xylol <i>Xylene</i>	4.0 %
Ronnel (O,O-dimethyl O(2,4,5-trichlorophenyl) phosphorothioate	1.0 %
-Related compounds	.5 %
INERT INGREDIENTS	85.0 %

Net contents 6 ounces

SUPERIOR CHEMICAL PRODUCTS, INC.
3942 Frankford Ave. Philadelphia Penna. 19124

Camerman's Note
Poor Copy

100

ACCEPTED
July 13, 1968
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PEST CONTROL
REGISTERED UNDER NO. 998-69
TO ATTACHED COMMENTS.

(Back panel)

S U P R O

WARNING: SUPRO insecticide is poisonous if swallowed, inhaled or absorbed through skin. Do not get in skin or eyes. Wear natural rubber gloves, protective clothing and goggles, when handling. Do not breathe fumes. Wear a protective mask or respirator of type approved by USDA for Vapona. Keep all unprotected personnel out of operational area. In case of contact, wash immediately with soap and water. Wash face, hands and arms thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and water before reuse. Contents are under pressure. High temperature may burst container. Do not expose to direct sunlight, radiators, stove, hot water or other sources of heat.

ANTIDOTE AND FIRST AID TREATMENT

Atropine is the emergency antidote for Vapona insecticide. Consult your physician about an emergency supply of 1/100 grain atropine tablets for emergency use. CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.

INTERNAL: If swallowed, induce vomiting immediately, by either introducing a finger into the throat or giving warm salt water to swallow. (1 tb per glass). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If spilled on skin, remove patient from area where insecticide, remove contaminated clothing and wash skin with soap and running water. If material gets into eyes, wash immediately with running water for at least 10 minutes.

IF WARNING SYMPTOMS APPEAR, administer 2 1/100 gr. atropine tablets immediately. NEVER GIVE TABLETS UNLESS SYMPTOMS APPEAR. (See warning symptoms below) Keep patient prone and quiet. Use artificial respiration immediately if patient is not breathing. Take patient immediately to nearest physician

NOTE TO PHYSICIAN: WARNING SYMPTOMS include weakness, headache, tightness in chest, blurred vision, non-reactive pinpoint pupils, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there any signs of this, atropine sulfate should be injected at 10 minute intervals in doses of 1 to 2 milligrams until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

PRECAUTIONS IN USING: SUPRO is poisonous, thereby precautions must be observed when handling. (See antidote and warnings. Store in a safe place. Children or animals must not be able to reach or handle. Do not contaminate feed, water or feedstuffs.

DISPOSAL OF CONTAINER: Use care in handling empty can. Completely empty used can by removing to open air away from children or pets, and press release button until completely empty. Puncture can to prevent reuse. Bury empty can 18" deep in isolated area away from water supplies.

USDA Reg. # 998-

COPY