


SUPERIOR TEX 4

FOR CONTROL OF MOSQUITO LARVAE, MOSQUITOES, HOUSE FLIES
CONTAINS BAYTEX

General Information

ACTIVE INGREDIENTS:
Fenthion O, O-Dimethyl O-
(3-methyl-4-methylthio)
phenyl phosphorothioate 45%
Aromatic Petroleum Distillate 47%
INERT INGREDIENTS: 8%
Total 100%

Baytex is a reg. TM of the Parent
Company of Farbentfabriken Bayer
GmbH, Leverkusen. U.S. Patent No.
3,042,703



Tex 4 emulsifiable insecticide is an
organic insecticide which gives ef-
fective control of mosquitoes and
flies, as well as ornamental pests
ants, aphids, mites, leaf-
hoppers, armyworms, juniper scale
and bagworms. Do not apply Tex 4
to hawthorn, American linden, sugar
maple, or the rose variety, Delightful,
because of possible plant injury.

Tex 4 is formulated for use only by
Pest Control Operators, Commercial
Nurserymen, Public Health Officials
and Mosquito Abatement Districts.

FOR CONTROL OF INSECTS ON ORNAMENTALS, NURSERY PLANTS, FOREST AND SHADE TREES:

TREES AND SHRUBS	Arborvitae	Dogwood	Oak	Tulip tree
	Ash	Euonymus	Spruce	Viburnum
	Aspen	Juniper	Sweet gum	Yew
	Birch	Maple		
FLOWERING PLANTS	Begonias	Lilies	Phlox	Snapdragons
	Chrysan- themums	Merigolds	Petunias	Verbena
	Geraniums	Peonies	Roses	Zinnias

Directions for Use

HOW TO MIX SPRAY:
Add 2 teaspoons TEX 4 to one gal-
lon of water (2 pints per 100 gallons
or 0.123% concentration)

Precautionary Note:
Do not apply residual sprays (1.5%
to 2.25%) to ornamentals as these
higher concentrations will cause
plant injury.

How to Apply:
Spray when air is still to avoid drift.
Do not treat plants in direct sunlight
during the heat of day or when the
temperature is in excess of 90
degrees F. Apply thorough coverage
to upper and lower surfaces of
leaves. Avoid wetting plants exces-
sively-only a fine mist is required.
Repeat as necessary.

MOSQUITOES, HOUSE FLIES:
Residual Spray: For long residual
control of mosquitoes and residual
fly control, apply 4 ounces in 1 gallon
of water and apply it to 500 square
feet or to run-off in localized areas on
patios or other outdoor structural
surfaces, in buildings, including dairy,
poultry and beef barns. Do not apply
directly to animals. Residual control
lasts 7 to 8 weeks for the lower rate
and even longer at the higher rate.
Repeat as necessary. Do not apply as
space spray.

SPRAYING: For mosquito larvae
control, applications should be made
to moist areas, standing water pools,
marshes, swamps and tidal areas.
Also spray around and in catch
basins where mosquitoes breed. Do
not apply to ponds, lakes, and
streams. Allow at least three weeks
between applications. For large-area
mosquito control, applications should
be made to wet grassy, bushy or
woody areas. For localized adult
mosquito control, applications should
be made to surface of buildings,
screen doors, window frames and
around outdoor light fixtures. For fly
control, spray around windows and
on other surfaces frequented by flies
on or in buildings, except where
food or feed products are present.
Use only in intermittent flooded
areas, standing water, temporary rain
pools, and log ponds.

WARNING: KEEP OUT OF REACH OF CHILDREN

RESTRICTIONS:

Do not use, pour, spill, or store
near heat or open flame.

1. Do not apply oil mixtures to forage
crops or rice.
2. Do not apply when weather
conditions favor drift of spray.
3. Do not use inside homes or in
buildings where food is processed or
stored.
4. Do not apply oil spray directly to
flowers, or other ornamental plants
which may be damaged by the oil.
Fog applications will not cause plant
injury.

ENVIRONMENTAL CAUTIONS:

This product is toxic to fish, birds
and other wildlife. Do not spray
where runoff is likely to occur. Do
not apply when weather conditions
favor drift from areas treated. Shrimp
and crab may be killed at application
rates recommended on this label. Do
not apply where these are important
resources. Do not contaminate water
by cleaning of equipment or disposal
of wastes. Apply this product only as
specified on this label.

This product is highly toxic to bees
exposed to direct treatment or resi-
dues on crops. Protective informa-
tion may be obtained from your
Cooperate Agricultural Extension
service.

**WASTE AND CONTAINER DIS-
POSAL:** Do not re-use container.
Destroy it by perforating or crushing.
Bury or discard in a safe place. Do
not contaminate water by cleaning of
equipment or disposal of wastes.

WARNING: May be fatal if swal-
lowed, inhaled, or absorbed through
the skin. Rapidly absorbed through
the skin. Do not get in eyes, on skin,
or on clothing. Do not breathe spray
mist. Wash thoroughly with soap and
warm water after handling. Wash
clothing with soap and hot water
before reuse. Avoid contamination of
feed or food. Keep out of reach of
children.

TO PHYSICIAN:

Antidote- Administer atropine sulfate
in large therapeutic doses. Repeat as
necessary to the point of tolerance 2
PAM is also antidotal and may be
administered in conjunction with
atropine.

EPA REG. NO. 998-117

ESTABLISHMENT NO. 998-PA-1

NET CONTENTS 1 GALLON

SUPERIOR CHEMICAL PRODUCTS INC.

3942 FRANKFORD AVENUE

PHILADELPHIA, PA. 19124