

EPA No. 1001-11

EPA Est. No. 1001-NJ-1

SPOTRETE F

FLOWABLE THIRAM TURF FUNGICIDE AND ANIMAL REPELLENT

Active ingredient	
Thiram (Tetramethylthiuram disulfide)*	..... 42%
Inert ingredients	..... 58%

\*Equivalent to 4 lbs/US gallon or 480 g/l

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF On SKIN: Wash with plenty of soap and water. Get medical attention.

IF In EYES: Flush eyes with plenty of water. Call a physician if irritation persists.


SEE RIGHT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

W. A. CLEARY CORPORATION

1049 Somerset St.

Somerset, N.J. 08873

**ACCEPTED**  
 MAY 14 1985  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act,  
 as amended for the pesticide  
 registered under  
 EPA Reg. No. 1001-11



## PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Wash thoroughly with soap and water after handling. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Harmful if inhaled. Avoid breathing spray mist. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling. Do not take alcohol immediately before or after exposure to this product. The consumption of alcohol by the applicator can increase the toxicity of the product. This product can cause skin itching, rashes and other dermal reactions to the user. Do not graze treated areas or feed clippings to livestock

## ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly to water or wetlands. This product is toxic to fish. Do not apply where conditions favor drift from areas treated. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

## PHYSICAL AND CHEMICAL HAZARDS

Keep away from heat and open flame

## STORAGE AND DISPOSAL

**STORAGE:** Do not contaminate water, food, or feed by storage or disposal. Store in original containers only in a secured area inaccessible to children and unauthorized persons. Absorb spills onto clay, sand or vermiculite and hold for proper disposal. Keep from freezing.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## LIMITED WARRANTY AND DISCLAIMER

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of fitness or of merchantability, and no agent of seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

GENERAL INFORMATION

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SPOTRETE-F prevents the infestation of common turf diseases such as Dollar Spot and Brown Patch if sprayed at regular weekly interval throughout the summer season.

SPOTRETE-F will control Snow Mold when 2 applications are made during Fall through Spring season.

SPOTRETE-F will control Dollar Spot and Brown Patch when applied at a higher dosage to turf already infested with these diseases.

SPOTRETE-F, when used according to label directions, protects nursery stock shrubs and ornamentals from rabbit, deer, and rodent depra-dation. One application will protect treated parts from 1 to 3 months, depending upon weather conditions. Spotrete-F is also useful as a repellent in deterring deer migration onto airport runways as a result of local feeding and browsing. The addition of a latex based sticker, such as CLEARSPRAY, prolongs the residual effectiveness of the treatment. Repellency is the result of rendering the treated plant parts distasteful.

DIRECTIONS FOR USE  
TURF APPLICATIONS

PREVENTATIVE DOSAGE: For Dollar Spot or Brown Patch apply 3<sup>3</sup>/<sub>4</sub> fluid ounces of Spotrete-F per 1000 square feet (5 quarts per acre) in 5 to 10 gallons of water. Spray at seven day intervals throughout the warm season. For snow mold apply 12 fluid ounces of Spotrete-F per 1000 square feet (4 gallons per acre) in 5 to 10 gallons of water. Spray in the autumn prior to the first snowfall. Repeat application in early spring.

CURATIVE LOSAGE: For Dollar Spot and Brown Patch apply 7<sup>1</sup>/<sub>2</sub> fluid ounces per 1000 square feet (2<sup>1</sup>/<sub>2</sub> gallon per acre) in 5 to 10 gallons of water. Apply when disease first appears and repeat application at 7 to 10 day intervals, as necessary.

RABBIT AND RODENT REPELLENCY

SPRAY APPLICATION, GROWING SEASON

Add 1 quart Spotrete-F to 7 quarts of water and apply as a full coverage spray to all portions of the plant to be protected. Spray to runoff. For greater retention and residual efficacy add 1 pint of latex based sticker to the spray solution. Do not use wetting agents or surfactants as foliar damage may result. For best results spray when temperatures are greater than 40° F and relative humidity is less than 70 percent. Coverage depends on plant density.

SPRAY APPLICATION, TREE TRUNKS AND DORMANT PLANTS

Add 1 quart of Spotrete-F to 3 quarts of water and apply as a spray to all portions of the plant to be protected. Spray to runoff. For greater retention and residual efficacy add 1 pint of latex based sticker to the spray solution. Coverage depends on stem density.

BRUSH ON APPLICATION

Apply Spotrete-F undiluted with a paint brush to stem parts and trunks to be protected. For greater retention and residual efficacy blend 4 fluid ounces of latex based sticker into 1 gallon Spotrete-F prior to brushing.

DEER REPELLENCY

SPOTRETE-F is useful as a repellent for deterring deer from migrating onto airport runways. It works by discouraging deer from localized feeding and browsing in areas adjacent to the airport. The product is most effective when used in combination with mechanical deterrents such as ditches, fences, etc. Prepare and spray Spotrete-F exactly as for rabbit and rodent repellency (above), depending upon season and type of application desired. Spray to a height of 5 feet (1.5 meters).

Printed in USA

Made in USA 12/84

