

4/24/72

METHAR 30
DISODIUM METHANEARSONATE
For post-emergent weed control
in cotton
EPA REG. NO. 1001-36

W. A. CLEAR CORPORATION
NEW BRUNSWICK, NEW JERSEY

CAUTION

Harmful if taken internally. Avoid contact with skin. Avoid breathing dust or spray mist. Wash thoroughly after using. Store in a safe place away from feed and food products. Keep children and domestic animals off treated areas until this product has been washed into the soil. DO NOT CONTAMINATE WATERS USED FOR DOMESTIC CONSUMPTION, OR WATER USED FOR IRRIGATION PURPOSES.

ANTIDOTE: IF TAKEN INTERNALLY, INDUCE VOMITING AND CALL PHYSICIAN AT ONCE.

Use of this product in a manner, or at any time other than in accordance with the directions may produce plant injury, excessive residues or other undesirable results.

NOTICE

No warranty, expressed or implied, is made concerning this material except that it conforms to the chemical description on the label. The buyer assumes all risk in using this material. The seller shall not be responsible for any personal injury or property damage resulting from the handling, storage or use of this material.

MIXING INSTRUCTIONS

Fill the spray equipment reservoir about half full with water and add the required amount of Methar 30 with agitation. Finish filling the reservoir with water and apply. Continue to agitate thoroughly during application. If the spray equipment is shut off when it still contains some Methar 30 solution, agitate thoroughly before applying again. Do not store spray solution in reservoir or other tank for a prolonged period and clean application equipment thoroughly after use by flushing with water.

UNDER THE FEDERAL INSECTICIDE AND RODENTICIDE ACT OF 1947
OR ECONOMIC POISON ACT
UNDER NO. 1001-36

METHAR 30
DISODIUM METHANE ARSONATE
LIQUID DSMA

APPLICATION METHODS

Methar 30 spray solutions should be applied with a low volume sprayer having satisfactory pumping and bypass action. ADJUST NOZZLES TO DIRECT THE SPRAY IN A MANNER TO ALLOW MAXIMUM COVERAGE OF WEEDS. KEEP SPRAY OFF COTTON FOLIAGE.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT

DIRECTIONS FOR USE

COTTON: Methar 30 is useful as a directed spray application to weeds in cotton for post-emergent control of weeds and grasses, including Johnsongrass, nutsedge, watergrass, goosegrass, dallisgrass, and cocklebur.

Mix at a rate of 1 1/2 gallons of Methar 30 (equivalent to 4 1/2 pounds DSMA Hexahydrate) plus 1.5 to 2 quarts of a suitable surfactant in 50 gallons of water. Apply at a broadcast rate of 40 gallons of this solution per acre. For band application, apply at a rate of one gallon of this solution for each one inch of band width to be treated on cotton grown in 40 inch rows. A second application may be required during humid growing conditions. Time the second application, if necessary, about 1 to 3 weeks after the first application.

APPLY ONLY WHEN COTTON IS 3 INCHES HIGH TO FIRST BLOOM. DO NOT FEED TREATED FOLIAGE TO LIVESTOCK, OR GRAZE TREATED AREAS.

Slight burning and a reddish discoloration of the cotton leaf may occasionally occur following the recommended treatment, but the cotton plant will develop normally.

NON-CROP: Methar 30 is useful for controlling Johnsongrass, nutsedge, watergrass, goosegrass and other weeds on drainage ditch banks, rights of way, storage yards and similar non-crop areas.

Mix Methar 30 at a rate of 1 1/2 gallons of Methar 30 (equivalent to 4 1/2 pounds DSMA Hexahydrate) plus 1.5 to 2 quarts of a suitable surfactant in 50 gallons of water. Spray the unwanted vegetation thoroughly to the point of run-off. Any spray equipment that gives good coverage may be used. If regrowth occurs, retreat as required.

4/24/72

METHAR 30
DISODIUM METHANE ARSONATE
FOR POST-EMERGENT WEED CONTROL
ON COTTON
EPA REG. NO. 1001-36

CAUTION

Harmful if taken internally. Avoid contact with skin. Avoid breathing dust or spray mist. Wash thoroughly after using. Store in a safe place away from feed and food products. Keep children and domestic animals off treated areas until this product has been washed into the soil. **DO NOT CONTAMINATE WATERS USED FOR DOMESTIC CONSUMPTION, OR WATER USED FOR IRRIGATION PURPOSES.**

ANTIDOTE: IF TAKEN INTERNALLY, INDUCE VOMITING AND CALL PHYSICIAN AT ONCE

Use of this product in a manner, or at any time other than in accordance with the directions may produce plant injury, excessive residues or other undesirable results.

No warranty, expressed or implied, is made concerning this material except that it conforms to the chemical description on the label. The buyer assumes all risk in using this material. The seller shall not be responsible for any personal injury or property damage resulting from the handling, storage or use of this material.

Fill the spray equipment reservoir about half full with water and add the required amount of Methar 30 with agitation. Finish filling the reservoir with water and apply. Continue to agitate thoroughly during application. If the spray equipment is shut off when it still contains some Methar 30 solution, agitate thoroughly before applying again. Do not store spray solution in reservoir or other tank for a prolonged period and clean application equipment thoroughly after use by flushing with water.

ACCEPTED
JUN 15 1972
UNDER THE FEDERAL INSECTICIDE ACT AND THE FEDERAL FUNGICIDE ACT, THIS PRODUCT IS LISTED AS A WEED KILLER FOR ECONOMIC PURPOSES, REGISTERED UNDER NO. 1001-36

METHAR 30
DISODIUM METHANE ARSONATE

Registration No. 1001-36

FOR POST-EMERGENT WEED CONTROL

Active ingredient: Disodium methanearsonate anhydrous	18.9 %
Inert ingredients	81.1 %
TOTAL	100 %
Net weight, all in water soluble form	7.65%
Equivalent to 3 pounds DISODIUM METHANEARSONATE MONOHYDRATE per gallon.	

CAUTION
KEEP OUT OF THE REACH OF CHILDREN
(See safety sheet for additional cautions)

NET CONTENT GALLONS
W. A. CLIBBY CORPORATION
NEW BRUNSWICK, N. J.

Methar 30 spray solutions should be applied with a low volume sprayer having satisfactory pumping and bypass action. **ADJUST NOZZLES TO DIRECT THE SPRAY IN A MANNER TO ALLOW MAXIMUM COVERAGE OF WEEDS. KEEP SPRAY OFF COTTON FOLIAGE.**
READ ENTIRE LABEL BEFORE USING THIS PRODUCT

COTTON

Apply to weeds in cotton fields after the cotton has reached the 4th true leaf stage. Apply to weeds in cotton fields after the cotton has reached the 4th true leaf stage. Apply to weeds in cotton fields after the cotton has reached the 4th true leaf stage.

NON-CROP

Apply to weeds in non-crop areas after the cotton has reached the 4th true leaf stage. Apply to weeds in non-crop areas after the cotton has reached the 4th true leaf stage. Apply to weeds in non-crop areas after the cotton has reached the 4th true leaf stage.