

MCP-2,4-D

POSTEMERGENCE BROADLEAF WEED CONTROL

ACTIVE INGREDIENTS:

2-(2-Methyl-4-chlorophenoxy)propionic acid, diethanolamine salt*...	31.5%
2,4-Dichlorophenoxyacetic acid, diethanolamine salt**.....	14.8%

INERT INGREDIENTS:	53.7%
--------------------------	-------

*2-(2-Methyl-4-chlorophenoxy)propionic acid equivalent 2 lb./gal. or 240 g/l or 21.2% by wt.

**2,4-Dichlorophenoxyacetic acid equivalent 0.95 lb./gal. or 114 g/l or 10.1% by wt. (isomer specific

by AOAC Method No. 6.D01-5)

KEEP OUT OF REACH OF CHILDREN

CAUTION


REGISTERED
 1001-47

SEE RIGHT PANEL FOR ADDITIONAL PRECAUTIONS

E.P.A. REG. NO. 1001-47

E.P.A. EST. NO. 1001-NJ-1

MANUFACTURED BY:



W-CLEARY CHEMICAL CORPORATION

1049 Somerset St. Somerset NJ 08873 (201) 247-8000

NET CONTENTS: 2-1/2 GALLONS

RIGHT PANEL

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed or absorbed through the skin. May cause eye and skin irritation. Avoid breathing spray mist. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Keep container closed. In case of contact with eyes, flush eyes immediately with a copious amount of water for at least 15 minutes and seek medical attention. If swallowed have the patient drink several glasses of water and induce vomiting. Seek medical attention. In case of contact with the skin wash the exposed area with soap and water. If irritation persists, seek medical attention.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or contaminate water by cleaning of equipment or disposal of waste. Do not contaminate irrigation ditches or water used for domestic purposes. Avoid spray contact or drift onto susceptible plants or any off-target plants. Do not spray when the wind is blowing sufficiently to cause drift. Use coarse sprays to minimize drift. Maintain spray boom pressure as low as possible to minimize drift. Never spray when the wind or gusts exceed 8 mph. Flush spray tank and lines in a suitable non-crop area after use. Do not use the same spray equipment to spray ornamentals or any other valuable or desirable plants.

TURF USE PRECAUTIONS

Do not use on new grass seedings until they have developed to the stage of first mowing. Do not use grass clippings from first mowing after treatment as a mulch for flowers or ornamental shrubs. Do not apply during a period of drought without watering thoroughly. Do not apply when temperatures exceed 90°F., as grass may be discolored or injured.

STORAGE AND DISPOSAL

- STORAGE:** Do not contaminate water, food or feed by storage or disposal. Store in original containers only in a secured area inaccessible to children and unauthorized persons.
- PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
- CONTAINER DISPOSAL:** Do not reuse empty container. Triple rinse (or equivalent) and offer for recycling or reconditioning or dispose of in a sanitary landfill, or by other approved Federal, state or local regulations.

**READ ENTIRE LABEL BEFORE USING THIS PRODUCT
GENERAL INFORMATION**

MCPP-2,4-D is a postemergence herbicide specifically formulated for use on turfgrass for selective control of many broadleaf weeds commonly found in lawns. MCPP-2,4-D is for use on bluegrass, fescue, Bermudagrass and bentgrass on fairways, greens, tees and aprons. Not for use on home lawns.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

APPLICATION RATE: For established turf, use 1½ fluid ounces in 1 to 2 gallons of water per 1000 square feet or 2 quarts per acre in 30-40 gallons of water. For St. Augustine grass or Centipede grass only, apply ¾ fluid ounce in 1 to 2 gallons of water per 1000 square feet or 1 quart per acre in 30-40 gallons of water.

MIXING: Use clean spray equipment. Partially fill the tank with clean water, add herbicide, mix thoroughly, and add the balance of water.

TIMING: Apply anytime weeds are young and actively growing. It is important that application be made when there is plenty of soil moisture, following a good rain or thorough watering. Do not water for at least 24 hours after application. If rain occurs within this period, effectiveness may be reduced, but wait at least a week to determine whether another application is necessary. Full effects of application may not be visible for about three weeks.

MOWING: Since action is through the leaves, avoid cutting or mowing turf for 1 to 2 days before and after spraying.

EQUIPMENT: Use equipment which applies product uniformly. A boom type sprayer is recommended.

The following is a partial list of weeds controlled by MCPP-2,4-D

Black medic	Dock	Mallow	Purslane
Common chickweed	English daisy	Mouse-ear chickweed	Ragweed
Clover	knotweed	Pigweed	Wood sorrel
Dandelion	Lambsquarters	Plantain	Yarrow

LIMITED WARRANTY AND DISCLAIMER

W. A. Cleary warrants that this material conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions and under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to W. A. Cleary. In no event shall W. A. Cleary's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.