

NET WEIGHT: 8 OZ.

FOR PREVENTION AND CONTROL OF
RED THREAD HELMINTHOSPORIUM
COPPER SPOT BROWN PATCH DOLLAR
SPOT FUSARIUM ROSEUM STRIPE SMUT

ACTIVE INGREDIENT:

Diethyl 4, 4'-o-phenylenebis
[3- thioallophanate] 50%

INERT INGREDIENTS: 50%

EPA Reg. No. 1001-50

CAUTION:

KEEP OUT OF THE REACH OF CHILDREN - MAY BE HARMFUL
IF SWALLOWED, AVOID CONTACT WITH EYES, SKIN, OR
CLOTHING. AVOID BREATHING DUST OR SPRAY MIST. IN
CASE OF CONTACT WASH OFF SKIN OR EYES IMMEDIATELY
WITH PLENTY OF WATER.

FOR EYES CALL A PHYSICIAN

SMALL AREA CONVERSION TABLE

When applying 3336 to smaller areas, the fungicide is frequently measured with tablespoons. Use the following conversion tables:

Level Tablespoons	Equivalent in ounces
4 tbs.	1/2 oz.
8 tbs.	1 oz.
16 tbs.	2 oz.

**DOLLAR SPOT
STRIPE SMUT**

**FUSARIUM ROSEUM
BROWN PATCH**

HELMINTHOSPORIUM

CAUTION:

IF SWALLOWED INDUCE VOMITING WITH A MIXTURE OF
SALT, SOAP IN WARM WATER. CALL A PHYSICIAN IMMEDI-
ATELY. DO NOT RE-USE THIS CONTAINER. BURY WHEN
EMPTY. DO NOT APPLY TO CROPS USED FOR FOOD OR LIVE-
STOCK. DO NOT GRAZE ANIMALS ON TREATED TURF. DO
NOT FEED CLIPPINGS TO LIVESTOCK OR POULTRY.

W-A-CLEARY

CORPORATION
Somerset, N. J. 08873

NET WEIGHT: 8 OZ.

FOR PREVENTION AND CONTROL OF
RED THREAD HELMINTHOSPORIUM
COPPER SPOT BROWN PATCH DOLLAR
SPOT FUSARIUM ROSEUM STRIPE SMUT

ACTIVE INGREDIENT:

Diethyl 4, 4'-o-phenylenebis
[3- thioallophanate] 50%

INERT INGREDIENTS: 50%

EPA Reg. No. 1001-50

KEEP OUT OF THE REACH OF CHILDREN - MAY BE HARMFUL
IF SWALLOWED, AVOID CONTACT WITH EYES, SKIN, OR
CLOTHING. AVOID BREATHING DUST OR SPRAY MIST. IN
CASE OF CONTACT WASH OFF SKIN OR EYES IMMEDIATELY
WITH PLENTY OF WATER.

FOR EYES CALL A PHYSICIAN

When applying 3336 to smaller areas, the fungicide is frequently
measured with tablespoons. Use the following conversion tables:


Level Tablespoons	Equivalent in ounces
4 tbs.	1/2 oz.
8 tbs.	1 oz.
16 tbs.	2 oz.

**DOLLAR SPOT
STRIPE SMUT**

**FUSARIUM ROSEUM
BROWN PATCH
HELMINTHOSPORIUM**

IF SWALLOWED INDUCE VOMITING WITH A MIXTURE OF
SALT, SOAP IN WARM WATER. CALL A PHYSICIAN IMMEDI-
ATELY. DO NOT RE-USE THIS CONTAINER. BURY WHEN
EMPTY. DO NOT APPLY TO CROPS USED FOR FOOD OR LIVE-
STOCK. DO NOT GRAZE ANIMALS ON TREATED TURF. DO
NOT FEED CLIPPINGS TO LIVESTOCK OR POULTRY.

W-A-CLEARY
CORPORATION
Somerset, N. J. 08873



3336 is a wettable powder for use on golf course greens, tees, and fairways of Bentgrasses, Bluegrasses, Bermuda grasses, Fescues, Ryegrasses, St. Augustine grasses or their mixtures. 3336 is not phytotoxic to any of the above mentioned grasses when used in accordance with the label. 3336 is to be used for the prevention and control of the below mentioned diseases.

Apply with hand sprayer, power sprayer, or a hose-on proportioner. Add the recommended amount of 3336 to a partially filled spray tank under agitation. Complete filling the tank; continuous agitation is necessary to insure even suspension of the wettable powder. Use spray mixture the same day it is prepared. Spray uniformly over the area to be treated. Apply after mowing or avoid mowing for twelve hours after application.

Apply recommended amounts in sufficient water to obtain thorough coverage, usually five gallons per 1,000 sq. ft. of turf area.

DOLLAR SPOT (*Sclerotinia homoeocarpa*), COPPER SPOT (*Gloeocercospora sorghi*), LARGE BROWN PATCH (*Rhizoctonia solani*), RED THREAD (*Corticium Fuciforme*), LEAF SPOT (*Helminthosporium spp.*) - Apply 3336 at 2 oz. per 1,000 sq. ft. beginning when the disease first appears and continue at 7-10 day intervals through the season or as needed.

FUSARIUM BLIGHT (*Fusarium roseum*) and STRIPE SMUT (*Ustilago striiformis*) - Apply 3336 at 4-8 oz. per 1,000 sq. ft. Apply applications at 7-10 day intervals beginning applications when the disease first appears. ³³

**IRRIGATE IMMEDIATELY AFTER APPLICATION TO DRENCH
3336 INTO THE ROOT ZONE.**

Stripe Smut usually occurs in the spring and the fall. Apply one treatment of 3336 at those times to prevent this disease.