

FRONT PANEL

135

3336 TM
TURF FUNGICIDE

For prevention and control of Dollar Spot, Copper Spot, Brown Patch, Red Thread, Fusarium Blight, Leaf Spot, and Stripe Smut on fine turf applications and for disease control on roses, flowers, ornamentals and shade trees.

Active Ingredient

Diethyl 4,4'-o-phenylenebis - (3-thioallophanate).....	50%
Inert Ingredients	50%
	100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautions

NET WT. 2 Pounds (907 grams)

EPA Reg. No. 1001-50
EPA Est. No. 36209-NJ-1

ACCEPTED
JUN 26 1984
 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 1001-50

W. A. CLEARY CORPORATION
SOMERSET, N.J. 08873

BEST DISEASE AVAILABLE

BACK PANEL

It is a violation of Federal Law to use this product in a manner inconsistent with its labelling.

3336 is a wettable powder for use on golf course greens, tees, and fairways of Bentgrasses, Bluegrasses, Bermuda grasses, Fescues, Ryegrasses, St. Augustine grasses or their mixtures. 3336 is not phytotoxic to any of the above mentioned grasses when used in accordance with the label.

DIRECTIONS FOR USE

Apply recommended amounts in sufficient water to obtain thorough coverage, usually five gallons per 1,000 sq.ft. of turf area:

DOLLAR SPOT (*Sclerotinia homoeocarpa*), COPPER SPOT (*Gloeocerospora sorghi*), LARGE BROWN PATCH (*Rhizoctonia solani*), RED THREAD (*Corticium fuciforme*) - Apply 3336 at 2 oz. per 1,000 sq.ft. beginning when the disease first appears and continue at 7-10 day intervals through the season, or as needed.

STRIP SMUT (*Ustilago striiformis*) - FUSARIUM BLIGHT (*Fusarium roseum*) - and LEAF SPOT (*Helminthosporium*) - Apply 3336 at 4-8 oz. per 1,000 sq.ft. Apply two applications at 7-10 day intervals, beginning applications when the disease first appears. IRRIGATE IMMEDIATELY AFTER APPLICATION.. TO DRENCH 3336 INTO THE ROOT ZONE. TO PREVENT STRIPE SMUT APPLY 3336 IN THE SPRING AND FALL.

DOLLAR SPOT CONTROL IN FAIRWAYS

Dollar spot is easily controlled with Cleary's 3336. For the prevention and control of this disease on fairways, apply one ounce per 1,000 sq.ft. at two to four week intervals.

ROSES, FLOWERS, ORNAMENTALS and SHADE TREES

Begin applications when disease first appears, or during suspected periods of disease incidence. Use Super Wet (3-4 oz./100 gal.), or equivalent wetting agent at recommended rate, for plants which have leaves that are hard to wet. Use at the recommended rates for the prevention or control of the following diseases:

FOLIAR SPRAY FOR FIELD & GREENHOUSE

<u>DISEASES CONTROLLED</u>	<u>LBS/100 GAL.</u>	<u>REMARKS</u>
Stem Blight of Juniper (<i>Phomopsis</i> spp.)	1 1/2 lb.	Repeat every 7-10 days as needed during disease periods
Rust of Crabapple (<i>Gymnosporangium</i> spp.)	1 lb.	Repeat every 10-14 days as needed during disease periods
Ovulinia Blight of Azalea & Rhododendron	1/2 lb.	Apply as flowers open. Repeat every 4-6 days as needed during disease periods
Anthracoze, Shade Trees & Woody Ornamentals	3/4 lb.	Apply at bud break. Repeat every 7-10 days as needed during disease period. Do not exceed a total of 4 applications per year.
Ascochyta Leaf Blight (<i>Ascochyta</i> spp.) Black Spot of Rose (<i>Diplocarpon</i> Roshe) Botrytis Flower Blight (<i>Botrytis</i> spp.) Scab of Pyracantha & Flowering Crab (<i>Fusicladium</i> spp. & <i>Venturia</i> spp.) Leaf Blight (<i>Curvularia</i> spp.) Corynespora Leafspot of Ligustrum (<i>Corynespora</i> spp.) Iris Leafspot (<i>Didymellina</i> , spp.)	3/4 lb.	Repeat every 7-10 days as needed during disease period
<u>DISEASES CONTROLLED</u>	<u>DRENCH TREATMENT LBS/100 GAL.</u>	<u>REMARKS</u>

BEST DOCUMENT AVAILABLE

STRIP SMUT (*Ustilago striiformis*) - FUSARIUM BLIGHT (*Fusarium roseum*)- and LEAF SPOT (*Helminthosporium*) - Apply 3336 at 4-8 oz. per 1,000 sq. ft. Apply two applications at 7-10 day intervals, beginning applications when the disease first appears. IRRIGATE IMMEDIATELY AFTER APPLICATION. TO DRENCH 3336 INTO THE ROOT ZONE. TO PREVENT STRIPE SMUT APPLY 3336 IN THE SPRING AND FALL.

DOLLAR SPOT CONTROL IN FAIRWAYS

Dollar spot is easily controlled with Cleary's 3336. For the prevention and control of this disease on fairways, apply one ounce per 1,000 sq. ft. at two to four week intervals.

ROSES, FLOWERS, ORNAMENTALS and SHADE TREES

Begin applications when disease first appears, or during suspected periods of disease incidence. Use Super Wet (3-4 oz./100 gal.), or equivalent wetting agent at recommended rate, for plants which have leaves that are hard to wet. Use at the recommended rates for the prevention or control of the following diseases:

FOLIAR SPRAY FOR FIELD & GREENHOUSE

DISEASES CONTROLLED	LBS/100 GAL.	REMARKS
Stem Blight of Juniper (<i>Phomopsis</i> spp.)	1½ lb.	Repeat every 7-10 days as needed during disease periods
Rust of Crabapple (<i>Gymnosporangium</i> spp.)	1 lb.	Repeat every 10-14 days as needed during disease periods
Ovulinia Blight of Azalea & Rhododendron	½ lb.	Apply as flowers open. Repeat every 4-6 days as needed during disease periods
Anthracoze, Shade Trees & Woody Ornamentals	¾ lb.	Apply at bud break. Repeat every 7-10 days as needed during disease period. Do not exceed a total of 4 applications per year.
Ascochyta Leaf Blight (<i>Ascochyta</i> spp.) Black Spot of Rose (<i>Diplocarpon</i> Roshe) Botrytis Flower Blight (<i>Botrytis</i> spp.) Scab of Pyracantha & Flowering Crab (<i>Fusicladium</i> spp. & <i>Venturia</i> spp.) Leaf Blight (<i>Curvularia</i> spp.) Corynespora Leafspot of Ligustrum (<i>Corynespora</i> spp.) Iris Leafspot (<i>Didymellina</i> , spp.)	¾ lb.	Repeat every 7-10 days as needed during disease period

BEST DOCUMENT AVAILABLE

DISEASES CONTROLLED	DRENCH TREATMENT LBS/100 GAL.	REMARKS
Stem, Crown & Root Rot of Herbaceous Annual, Perennials & Bedding Plants (<i>Botrytis</i> spp., <i>Fusarium</i> spp., <i>Rhizoctonia</i> spp. & <i>Sclerotinia</i> spp.)	¾ lb.	Apply at rate of 2-3 pints/sq. ft. Apply after transplant to propagation bed or container
Cylindrocladium Rot of Azalea, Conifers, Poinsettias & Rhododendrons (<i>Cylindrocladium</i> spp.)		Repeat every 14-21 days as needed during disease period.

DISEASES CONTROLLED	BULB TREATMENT LBS/100 GAL.	REMARKS
Bulb Rot of Daffodil, Easter Lily, Gladiolus, Iris & Tulip (<i>Fusarium</i> spp. & <i>Penicillium</i> spp.)	¾-1 lb.	Soak immediately after digging for 30 minutes in dip (80-85° Dry well before storing. Heat treat before treatment if bulb are used for forcing.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. May cause irritation of eyes, nose, throat and skin. Avoid breathing dust. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse. In case of contact with eyes, flush with plenty of water for 15 minutes. Get medical attention if irritation persists. In case of contact with skin, wash with soap and water.

Do not graze animals on treated turf. Do not feed clippings to livestock or poultry.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply, allow to drift or drain or flush equipment onto or near croplands. Application equipment should be thoroughly cleaned after use.

STORAGE AND DISPOSAL

STORAGE Do not contaminate water, food, or feed by storage or disposal. Store in a cool, dry area, in original unopened container. Store in a secured area unavailable to unauthorized persons. : : :
PESTICIDE DISPOSAL Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. : : :
CONTAINER DISPOSAL Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. : : :

LIMITED WARRANTY AND DISCLAIMER

Seller warrants that the composition of this product conforms to the ingredient statement and is suited to the purposes described when used according to directions. The use of this product being beyond the control of the seller, no guarantee, expressed or implied, is made to the effects or results to be obtained if not used in accordance with directions or established safe practices. Buyer accepts all risks of use, storage or handling of this material not in strict accordance with directions given herewith.