

1001-51

ACCEPTED

Under the Federal Insecticide, Fungicide, and Rodenticide Act, this product is registered under the Fungicide, Fertilizer, and Plant Growth Regulator Act, Pesticide Registration Act, 1001-51

BROMOSAN

SYSTEMIC TURF FUNGICIDE

(WITH "CLEARY'S 3336")

FOR CONTROL OF DOLLAR SPOT, COPPER SPOT, ANTHRACNOSE, HELMINTHOSPORIUM, RUST, MELD AND SNOW MOLD

ORNAMENTAL TURF USE ONLY

ACTIVE INGREDIENT		
Triphenyltin hydroxide (as triphenyltin phosphate)	16.67%	
Triphenyltin disulfide	50.00%	
INERT INGREDIENTS		33.33%
EPA REG NO. 51		EPA EST NO. 36209-NJ-01

KEEP OUT OF REACH OF CHILDREN

CAUTION: May be harmful if swallowed. Avoid contact with eyes, skin or clothing. Avoid breathing dust or spray mist. In case of contact wash off skin or eyes immediately with plenty of water. For eyes call a physician. See other cautions on rear panel.

WT 18 LBS
6-3 LB Bags Per Case

W.A. CLEARY CORPORATION

1045 SOMERSET STREET SOMERSET, N.J. 08873

CAUTION :

Do not take alcohol before or after exposure to this product

Contact may cause allergic reactions

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of a label. Apply this product only as directed on this label.

Do not treat areas or feed clippings to livestock. Do not reuse container. Destroy when empty.

GENERAL INFORMATION

"Bromosan" is a specifically formulated wettable powder for use in fine Turfgrass areas, such as Golf Greens and Fairways for Turf Diseases. "Bromosan" is not phytotoxic, nor will it discolor THE FINEST BENI-GRASSES, BLUE GRASSES, FESCUE OR THEIR MIXTURES.

DIRECTIONS FOR USE AS A SPRAY

DOLLAR SPOT CONTROL: Mix 6 ounces BROMOSAN in 5 to 10 gallons water and apply to 2,000 sq. ft. Applications should start early spring as a preventive and repeat 7 to 10 days later as directed on the label.

OTHER DISEASES

During hot humid weather it may be necessary to increase the rate to 8 oz. per 1500 sq. ft. to control other diseases such as Copper Spot, Brown Patch.

For the control of Helminthosporium (Leaf Spot) use the rate 6 oz. per 1500 sq. ft.

SNOW MOLD CONTROL (Fusarium Nivale & Typhula sp.) For the Fall application, before first snow cover, mix 9-12 ounces BROMOSAN in 5-10 gallons water and apply to 1,000 sq. ft. For a mid-winter application, or during spring thaw, mix 4.5 ounces BROMOSAN in 5-10 gallons water and apply to 1,000 sq. ft. **DO NOT APPLY TO SNOW COVERED TURF.**

1001-51