

# Spotrete

## THIRAM TURF FUNGICIDE

EPA REG NO 1001-60  
EPA EST NO 36209-NJ-01

For the control of Dollar Spot, Brown Patch  
and Snow Mold

Active Ingredient  
Thiram (Tetramethylthiuram Disulfide) 75%  
Inert Ingredients 25%

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

May be harmful if swallowed. Avoid contact with eyes, skin or clothing. Contact may cause allergic reactions. Avoid breathing dust or spray mist. Do not take alcohol before or after exposure to this product.


#### PRACTICAL TREATMENT

If swallowed induce vomiting. Call a physician immediately. In case of contact wash off skin or eyes immediately with plenty of water. For eyes call a physician.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of any body of water. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

NET WT 3 LBS



#### INSTRUCTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**Note to User:** Do not graze treated areas or feed clippings to livestock.

SPOTRETE prevents the infestation of common turf diseases such as Dollar Spot and Brown Patch if sprayed at regular weekly intervals throughout the summer season.

SPOTRETE will control Snow Mold when 2 applications are made during Fall through Spring seasons.

SPOTRETE will control Dollar Spot and Brown Patch when applied at a higher dosage to turf already infested with these diseases.

#### Preventive Dosage:

A. For Dollar Spot and Brown Patch: 2 1/2 ozs per thousand square feet at seven day intervals throughout the warm season.

B. For Snow Mold: 8 ozs per thousand square feet late in the fall prior to snowfall. A Spring application of 3 ozs per thousand square feet during Spring thaw may be necessary.

#### Curative Dosage:

A. For Dollar Spot and Large Brown Patch: 5 ozs per thousand square feet. Begin when disease first appears. Repeat at 7 to 10 day intervals as necessary.

#### Amount of Water:

For uniform coverage use 5 to 10 gals per thousand square feet.

#### STORAGE AND DISPOSAL

**Prohibitions:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**Pesticide Disposal:** Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

**Pesticide Disposal:** Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

**Container Disposal:** Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

**General:** Consult federal, state, or local disposal authorities for approved alternative procedures such as limited open burning.

W-A-CLEAR-Y

SOMERSET, NEW JERSEY 08873

42

BEST DOCUMENT AVAILABLE