

MAR 7 1994

Mr. Richard F. Fletcher
W.A. Cleary Chemical Corp.
1049 Somerset St.
Somerset, NJ 08875-0010

Dear Mr. Fletcher:

Subject: 3336 WP
EPA Reg. No. 1001-63
Your Submission of March 3, 1994

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable provided that you:

1. Submit and/or cite all data required for registration or reregistration of your product under FIFRA sec. 3(c)(5) and sec. 4 when the Agency requires all registrants of similar products to submit such data.

2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Note that the worker protection standard labeling is being reviewed by a separate group and acceptability of that labeling will be determined by them. The acceptance of this label does not imply acceptance of the worker protection labeling. You may use this labeling immediately if you have elected the self-verification method of compliance but you must make appropriate revisions if required by the Agency Worker Protection Standard Review Group.

Sincerely yours,

15
Sidney C. Jackson
Acting Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (7505C)

Encl.
7505C:C.Grable:cg:3/7/94

BEST AVAILABLE COPY

2 of 9

SPECIMEN LABEL

VERSION 4

3336 WP

IN WATER SOLUBLE BAGS

TURF AND ORNAMENTAL SYSTEMIC FUNGICIDE

For the prevention and control of fine turf diseases and diseases of Annual and Perennial Flowers, plus Deciduous and Evergreen Trees and Shrubs.

ACTIVE INGREDIENTS

Dimethyl 4,4-o-phenylenebis-(3,thioallophanate).....50%

INERT INGREDIENTS.....50%

TOTAL.....100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water, induce vomiting by touching the back of the throat, contact a physician or poison control center.

IF INHALED: Remove to fresh air.

IF ON SKIN: Wash exposed areas with soap and water, seek medical attention as needed.

IF IN EYES: Flush eyes with water, contact a physician for irritation as needed.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

SOLD BY:

CLEARY LOGO

SOMERSET, N.J. 08875

ACCEPTED
with COMMENTS
in EPA Letter Dated

MAR 07 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

1001-63

EPA Registration No. 1001-63

EPA Est. No. 33967-NJ1

NET CONTENTS 2 LBS. (907.2 grams)

BEST AVAILABLE COPY

3019

PRECAUTIONARY STATEMENTS-HAZARDS TO HUMANS AND DOMESTIC ANIMALS

PRECAUTIONARY STATEMENTS: Harmful if swallowed, inhaled or absorbed through the skin. May cause irritation of eyes, nose, throat, and skin. Avoid breathing dust and contact with eyes and skin. Do not graze animals on treated turf. Do not feed clippings to livestock or poultry.

PERSONAL PROTECTIVE EQUIPMENT

WPS USES (Nursery and Greenhouse Uses): Applicators and other handlers must wear long-sleeved shirt, long pants, and shoes with socks.

NON-WPS USES (Turf and Ornamental Uses): Applicators and other handlers should wear long-sleeved shirt, long pants, and shoes with socks.

USER SAFETY RECOMMENDATIONS

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, where surface water is present, or to intertidal areas below the mean high water mark. Application equipment should be thoroughly cleaned after use. Do not contaminate water when disposing of equipment washwater. Do not apply, allow to drift, or drain, or flush equipment onto non-target areas.

STORAGE AND DISPOSAL

STORAGE:Do not contaminate water, food, or feed by storage or disposal. Store in a cool, dry area in original unopened container. Store in a secured area unavailable to unauthorized persons.

PESTICIDE DISPOSAL:Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:When all packets are used, dispose of empty foil package in a sanitary landfill or by incineration or as allowed by State and local authorities by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

AGRICULTURAL USE REQUIREMENTS (Greenhouse and Nursery Uses Only)

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific statements and exceptions pertaining to the statements on this label about personal protective equipment(PPE) and restricted-entry interval. The requirements in this paragraph only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Personal Protective Equipment (PPE) required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is; coveralls, waterproof gloves, and shoes with socks.

GENERAL INFORMATION

Apply 3336WP with ground or aerial equipment, using sufficient volume of spray to provide thorough coverage. Add required amount of 3336WP to a partially filled tank, agitate by mechanical or hydraulic means and then add remaining required amount of water. Continuous agitation is required to keep the material in suspension. Do not tank mix with copper-containing materials or with highly alkaline pesticides such as Bordeaux mixture of lime sulfur. No claim of compatibility with other pesticides is implied. Use the higher rates under conditions of severe disease pressure. Also see local State Extension Services recommendations for application schedules. **IMPORTANT**, if after using 3336 WP as recommended a treatment is not effective, a tolerant strain of fungi may be present. Consult your local Cleary representative or your State Agricultural Extension Service for advice on the prompt use of some other suitable fungicide. **NOTICE:** Read and understand the entire label.

TURF

BEST AVAILABLE COPY

For use on all fine turf applications such as commercial and home lawns, athletic fields, cemeteries, parks, golf course greens, tees, and fairways of Bentgrasses, Bluegrasses, Bermudagrasses, Fescues, Ryegrasses, St. Augustine grasses or their mixtures. 3336 WP is not phytotoxic to any of the above mentioned grasses when used

409

in accordance with the label. 33 WP is to be used for the prevention and control of the below mentioned diseases (provided both curative and protective action).

Use spray mixture the same day it is prepared. Spray uniformly over the area to be treated with a properly calibrated power sprayer. Apply after mowing or avoid mowing twelve hours after application.

Apply recommended amounts in sufficient water to obtain through coverage, (three to five gallons per 1,000 square feet suggested). When treating golf greens, always treat aprons and approaches. Use the highest recommended rate under conditions of severe disease pressure.

DISEASE(S) CONTROLLED	RATE	REMARKS
Anthracnose (Colletotricum graminicola)	2 oz.	Apply when disease first appears. Continue at 10-14 day intervals as needed.
Dollar Spot (Mollorodiscus spp., Lanzia spp.)		Apply when disease first appears. Continue at 7-10 day intervals as needed.
Copper Spot (Gloeocerospora sorghi)		
Large Brown Patch (Rhizoctonia solani)		
Red Thread (Corticium fuciforme)		
Pink Snow Mold (Fusarium nivale)		Apply first application Mid-Late November and continue at 10-14 day intervals throughout the season. May be used in combination with Cleary's Spotrete and PCNB products at reduced rates.
Summer Patch (Magnaportha poae)	4-8 oz.	For prevention, apply 3 applications beginning early May using 21 day intervals. Water material into root zone thoroughly soon after application. For suppression, apply two applications at 10-14 day intervals beginning when disease first appears.
Fusarium Blight (Fusarium roseum and triticum)		Apply two applications at 10-14 day intervals beginning when disease first appears.
Necrotic Ring Spot (Leptosphaeria korrae)		
Stripe Smut (Ustilago striiformis)		For prevention apply 3336 WP in the spring and fall. Apply two applications at 7-10 day intervals when disease first appears. Water material into root zone soon after application.
Leaf Spot (Dreschlera spp.)		Apply two applications at 7-10 day intervals when disease first appears. Water material into root zone soon after application. May be used in combination with Cleary's Protect T/O at lower rate.

such as
HORTICULTURAL USES (FIELD, NURSERY AND GREENHOUSE) *and landscape*
ANNUAL AND PERENNIAL FLOWERS, ^fDECIDUOUS AND EVERGREEN TREES AND SHRUBS.
GROUND COVERS

Begin applications when disease first appears, or during suspected periods of disease incidence. Apply additional applications every 7-10 days or as otherwise instructed for the prevention or control of the listed diseases. Use of a wetting agent such as Cleary's Super Wet is recommended for plants that have leaves that are difficult to wet properly. Use of a spreader-sticker such as Cleary's Clearspray is recommended to enhance product performance in wet weather conditions. Cleary's 3336 WP may be used to control listed diseases on "backyard" fruit trees such as almond, apple, apricot, cherry, nectarine, peach, pecan, plum, and prune trees. Do not apply within 1 day (24 hours) of harvest.

NON-COMMERCIAL

DISEASE(S) CONTROLLED - FOLIAR	RATE/100 GAL	REMARKS
Ovulinia Blight of Azalea and Rhododendron (Ovulinia spp.)	8 oz.	Apply as flowers open. Repeat every 7-10 days as needed during disease period.
Powdery Mildew (Oidium spp.)		Apply mid-summer or at first sign of disease.
Summer Patch (Magnaportha poae)	12-16 oz.	
Anthracnose		Apply as buds break. Repeat every 7-10 days as needed during disease period. Do not exceed four applications per season.

5029

Ascochyta Leaf Blight	12-16 oz.	Repeat every 7-10 days as needed during disease period.
Black Spot of Rose (Diplocarpon rosae)		Apply mid-summer or at first sign of disease.
Botrytis Flower Blight		
Corynespora Leafspot		
Iris Leaf Spot (Didymellina spp.)		
Curvularia Leaf Blight		
Rust of Crabapple (Gymnosporangium spp.)		Repeat every 10-14 days as needed during disease periods.
Leaf Scabs (Fusicledium and Venturia spp.)		Apply mid- ^{SPRING} summer or at first sign of disease. (Crabapple, Pyracantha, Hawthorn, Pear, Mountain Ash)
Tip Blight of Pine (Sphaeropsis sapinea, Diplodia pinea)		Apply along with a compatible spreader/sticker at recommended rates to entire tree, thorough coverage is essential for optimal disease control. Begin application in the spring when new growth starts. Make a second application just before needles emerge from the sheath and a third application 10 days later.
Twig Blight of Juniper (Phomopsis spp., Kabatina spp.)	24 oz.	Repeat every 7-10 days as needed during disease period.
DISEASE(S) CONTROLLED - DRENCH	RATE/100 GAL.	REMARKS
Stem Rots, Crown Rots, and Root Rots (Botrytis spp., Fusarium spp., Gleocladium spp., Myrothecium, Ramularia spp., Rhizoctonia spp., and Sclerotinia spp.)	12-16 oz.	Apply after transplanting to propagation bed, container, or pot at a rate to thoroughly soak the growing media (usually 1-3 pints per square foot depending on the media type and depth). Repeat every 21-28 days to provide adequate crop protection.
Black Root Rot (Thielaviopsis spp.)		
Cylindrocladium Rot		
DISEASE(S) CONTROLLED - BULB TREATMENT	RATE/100 GAL.	REMARKS
Bulb, Corm, and Rhizome Rots (Fusarium spp. and Penicillium spp.)	12-16 oz.	Soak immediately after digging for 30 minutes in dip (80-85 degrees F). Dry well before storing.

LIMITED WARRANTY AND DISCLAIMER
W.A. CLEARY CHEMICAL CORPORATION warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. W.A. CLEARY CHEMICAL CORPORATION MAKES NO OTHER EXPRESSED OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESSED OR IMPLIED WARRANTY. IN NO CASE SHALL W.A. CLEARY OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT INCLUDING, BUT NOT LIMITED TO LOSS OF PROFITS, BUSINESS REPUTATION OR CUSTOMERS, LABOR COSTS OR OTHER EXPENSES INCURRED IN PLANTING OR HARVESTING. W.A. CLEARY and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of W.A. CLEARY.

BEST AVAILABLE COPY

W.A. CLEARY CHEMICAL CORPORATION
Corporate Headquarters Sales and Distribution Center
1049 Route 27, P.O.Box 10 178 Ridge Road, Suite A
Somerset, New Jersey 08875-0010 Dayton, N.J. 08810
(800) 238-7813 (908) 329-8399

3336 WP

IN WATER SOLUBLE BAGS



For the prevention and control of turf diseases and the diseases of Annual and Perennial flowers, ground covers, plus Deciduous and Evergreen trees and shrubs.

ACTIVE INGREDIENTS

Dimethyl 4, 4-o-phenylenebis-(3,thioallophanate)	50%
Inert Ingredients.....	50%
TOTAL	100%



CAUTION

STATEMENT OF PRACTICAL TREATMENT

- **IF SWALLOWED:** Drink 1 or 2 glasses of water, induce vomiting by touching the back of the throat, contact a physician or poison control center.
- **IF INHALED:** Remove to fresh air.
- **IF ON SKIN:** Wash exposed areas with soap and water, seek medical attention as needed.
- **IF IN EYES:** Flush eyes with water, contact a physician for irritation as needed.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

BEST AVAILABLE COPY

SOLD BY:



W.A. Cleary Chemical

C O R P O R A T I O N

Somerset, NJ 08875

EPA Reg. No. 1001-63 EPA Est. No. 33967-NJ1

NET CONTENTS 2 LBS.

7049

PRECAUTIONARY STATEMENTS-HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed, inhaled or absorbed through the skin. May cause irritation of eyes, nose, throat, and skin. Avoid breathing dust and contact with eyes and skin. Do not graze animals on treated turf. Do not feed clippings to livestock or poultry.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

WP USES (Nursery and Greenhouse Uses): Applicators and other handlers must wear long-sleeved shirt, long pants, and shoes with socks.

NON-WPS USES (Turf and Ornamental Uses): Applicator and other handlers should wear long-sleeved shirt, long pants, and shoes with socks.

USER SAFETY RECOMMENDATIONS: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, where surface water is present, or to intertidal areas below the mean high water mark. Application equipment should be thoroughly cleaned after use. Do not contaminate water when disposing of equipment washwater. Do not apply, allow to drift, or drain, or flush equipment onto non-target areas.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal. Store in a cool, dry area in original unopened container. Store in a secured area unavailable to unauthorized persons.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: When all packets are used, dispose of empty foil package in a sanitary landfill or by incineration or as allowed by State and local authorities by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific statements and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Personal Protective Equipment (PPE) required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, and shoes with socks.

GENERAL INFORMATION

Apply 3336 WP with ground or aerial equipment, using sufficient volume of spray to provide thorough coverage. Add required amount of 3336 WP to a partially filled tank, agitate by mechanical or hydraulic means and then add remaining required amount of water. Continuous agitation is required to keep the material in suspension. Do not tank mix with copper-containing materials or with highly alkaline pesticides such as Bordeaux mixture of lime sulfur. No claim of compatibility with other pesticides is implied. Use the higher rates under conditions of severe disease pressure. Also see local State Extension Services recommendations for application schedules. **IMPORTANT**, if after using 3336 WP as recommended a treatment is not effective, a tolerant strain of fungi may be present. Consult your local Cleary representative or your State Agricultural Extension Service for advice on the prompt use of some other suitable fungicide. **NOTICE: Read and understand the entire label.**

BEST AVAILABLE COPY

For use on all fine turf applications such as commercial lawns, athletic fields, home lawns, cemeteries, golf course greens, tees, and fairways of Bentgrasses, Bluegrasses, Bermudagrasses, Fescues, Ryegrasses, St. Augustine grasses or their mixtures. 3336 WP is not phytotoxic to any of the above mentioned grasses when used in accordance with the label. 3336 WP is to be used for the prevention and control of the below mentioned diseases (provided both curative and protective action).

For best results use spray mixture the same day it is prepared. Spray uniformly over the area to be treated with a properly calibrated sprayer. Apply after mowing or avoid mowing twelve hours after application.

Apply recommended amounts in sufficient water to obtain through coverage, (3-5 gallons per 1,000 square feet suggested). Use the highest recommended rate under conditions of severe disease pressure.

DOLLAR SPOT (*Sclerotinia Homoeocarpa*), **COPPERSPOT** (*Gloeocerospora Sorghi*), **LARGE BROWN PATCH** (*Rhizoctonia Solani*), **RED THREAD** (*Corticium Fuciforme*), and **FUSARIUM PATCH** (*Fusarium Nivale*) - Apply 3336 WP at 8 oz. (1 packet) per 4,000 sq. ft. beginning when the disease first appears and continue at 7-10 day intervals through the season or as needed.

FUSARIUM BLIGHT (*Fusarium Roseum*), **SUMMER PATCH** (*Magnaporthe Poae*), **NECROTIC RING SPOT** (*Leptosphaeria Korrea*) - Apply 3336 WP at 8-16 oz. (1-2 packets) per 2,000 sq.ft.. Water into the root zone by irrigating with one inch of water immediately after application. Make two applications at 10-14 day intervals beginning applications when the disease first appears.

ANTHRACNOSE (*Colletotricum Graminicola*) - Apply 8 oz.(1 packet) per 8,000 sq.ft. beginning when the disease first appears and continue at 10-14 day intervals as needed. As a curative, use 8 oz. (1 packet) per 4,000 sq.ft..

STRIPE SMUT (*Ustilago Striformis*) and **LEAF SPOT** (*Helminthosporium*) - Apply 3336 WP at 8-16 oz.(1-2 packets) per 2,000 sq.ft.. Apply two applications at 7-10 day intervals, beginning applications when the disease first appears. Irrigate immediately after application to drench 3336 WP into the root zone. To prevent Stripe Smut apply 3336 WP in the spring and fall.

DOLLAR SPOT (control in fairways) - Dollar spot is easily controlled with Cleary's 3336 WP. For the prevention and control of this disease on fairways, apply 8 oz. (1 packet) per 8,000 sq.ft. at 2-4 week intervals.

ORNAMENTALS, FIELD AND GREENHOUSE

ROSES, FLOWERS, ORNAMENTALS, and SHADE TREES - Begin applications when disease first appears, or during suspected periods of disease incidence. Use Super Wet 3-4 oz. per 100 gallons, or equivalent wetting agent at recommended rates for plants which have leaves that are hard to wet. Use at the recommended rates for the prevention or control of the following diseases. Cleary's 3336 WP may be used to control listed diseases on ornamental almond, apple, apricot, cherry, nectarine, peach, pecan, plum, and prune trees. Apply at early bloom. Make a second application at full bloom if conditions favor disease development. Do not apply within 1 day (24 hours) of harvest.

SPRAYED PLANT TYPES AND DISEASES CONTROLLED

DISEASE(S) CONTROLLED	OZ./100 GALLONS	REMARKS
Stem Blight of Juniper (<i>Phomopsis</i> spp.)	24 oz.	Repeat every 7-10 days as needed during disease periods.
Rust of Crabapple (<i>Gymnosporangium</i> spp.)	16 oz.	Repeat every 10-14 days as needed during disease periods.
Tip Blight	16 oz.	Apply along with a compatible spreader/sticker at recommended rates to entire tree. Thorough coverage is essential for optimum disease control. Begin application in the spring when new growth starts. Make a second application just before needles emerge from the sheath, and a third application ten days later.
Ovulinia Blight of Azalea and Rhododendron	8 oz.	Apply as flowers open. Repeat every 4-8 days as needed during disease periods.
Anthracoese, Shade trees and Woody Ornamentals	12 oz.	Apply as buds break. Repeat every 7-10 days as needed during disease period. Do not exceed a total of 4 applications per year.

BEST AVAILABLE COPY

Ascochyta Leaf Blight (Ascochyta spp.)	12 oz.	Repeat every 7-10 days as needed during disease period.
Black Spot of Rose (Diplocarpon Roshe)	12 oz.	
Botrytis Flower Blight (Botrytis spp.)	12 oz.	Apply mid-summer or at first sign of disease (Viburnum, Azalea, Catalpa, Crabapple, Hydrangea, Lilac, Rhododendron, and other woody plants) or at first sign of disease (Zinnia and flowering plants).
Scab of Pyracantha and Flowering Crabapple (Fusicadium spp. and Venturia spp.)	12 oz.	
Leaf Blight (Curvularia spp.)		
Corynespora Leafspot of Ligustrum (Corynespora spp.)		
Iris Leafspot (Didymellina spp.)		
Powdery Mildew (Erysiphe spp.)		

DRENCH TREATMENTS

DISEASE(S) CONTROLLED	OZ./100 GALLONS	REMARKS
Stem, Crown, and Root Rot of Herbaceous Annuals, Perennials, and Bedding Plants. (Botrytis spp., Fusarium spp., Rhizoctonia spp., and Sclerotinia spp.)	12 oz.	Apply at a rate of 2-3 pints per sq.ft.. Apply after transplant to propagation bed or container Repeat every 14-21 days as needed during disease periods.
Cylindrocladium Rot of Azalea, Conifers, Poinsettias, and Rhododendron. (Cylindrocladium spp.)		
Thielaveopsis Rot of Woody Ornamentals (such as Azalea, Rhododendron, Conifer and Poinsettias).		

BULB TREATMENTS

DISEASE(S) CONTROLLED	OZ./100 GALLONS	REMARKS
Bulb Rot of Daffodil, Easter Lily Gladiolus, Iris, and Tulip (Fusarium spp. and Penicillin spp.)	12-16 oz.	Soak immediately after digging for 30 minutes in dip (80-85°). Dry well before storing. Heat treat before treatment if bulbs are used for forcing.

Note: 3336 WP does not control *Pythium* spp. or *Phytophthora* spp.

BEST AVAILABLE COPY



COMPANY
NUMBER 34688

PRODUCT
SERIAL NO. 69

LABEL
APPROVAL
DATE 3-10-93

PM 22 34688-69

ACCEPTED

MAR 10 1993

By the Federal Insecticide, Fungicide, and Rodenticide Act, approved for the product **AKZO** No. 34688-69

SINESTO B

ACTIVE INGREDIENT

Alkyl* trimethylammonium chloride.....14.1%
(*Alkyl as in fatty acids of coconut oil)

INERT INGREDIENTS.....85.9%

TOTAL 100%

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES
Immediately flush eyes with plenty of water for at least 15 minutes. Then get immediate medical attention.

IF ON SKIN
Immediately wash exposed skin with soap and water. Get prompt medical attention if irritation develops. Remove contaminated clothing, including shoes, and thoroughly clean and dry before reuse.

IF SWALLOWED
Do not induce vomiting. Drink plenty of water. Get immediate medical attention.

IF INHALED
Move to fresh air. Get immediate medical attention.

NOTE TO PHYSICIAN
Possible mucosal damage may contraindicate the use of gastric lavage.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

MANUFACTURED BY
Akzo Chemicals Inc.
300 South Riverside Plaza
Chicago, Illinois 60606

EPA REGISTRATION NUMBER 34688-69
EPA ESTABLISHMENT NUMBER 34688-1L-1

NET VOLUME
55 gals. (208.2 liters)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
Danger. Corrosive. Causes severe eye and skin damage. Harmful or fatal if swallowed.

Do not get in eyes, on skin, or clothing. Avoid breathing vapor or mist. Wear chemical goggles, face shield, rubber gloves and impervious clothing when handling. Do not contaminate food. Do not take internally. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or public waters unless this product is specifically identified and addressed in an NPDES permit. Do not discharge effluent containing this product to sewer systems without previously notifying the sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA. Do not contaminate water by cleaning of equipment or disposal of waste. Apply this pesticide only as specified on this label.

STORAGE AND DISPOSAL

PROHIBITIONS
Observe handling instructions given to prevent eye damage and skin burns. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Plastic Containers: Triple rinse (or Equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL

Consult federal, state or local disposal authorities for approved alternative procedures.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with labeling.

Sinesto B is a concentrated ammonium salts in water solution used in lumber mills as an anti-sap stain on freshly cut lumber.

Sinesto B will prevent mildew stains (blue stain) and transportation of lumber. It is not intended or expected to be used on the use site. It is suitable for use with water and subsequent application on freshly cut lumber. Apply dilute solution by spraying. The wood should be treated within 24 hours after cutting.

Wear rubber gloves or eye protection when handling solution on freshly cut lumber.

Sinesto B should be diluted with 10 and 30 parts of water, depending on the severity of the stain. Vigorous stirring to mix. A board test is recommended to determine the optimum dilution rate. The dilution rate should be at least 30 seconds.

Treated wood should be kept dry after treatment and must be protected from inclement weather. Dip should be roofed, paved (or drained) to prevent dilution. Apply anti-sap stain solution. Follow to the application of the solution properly maintained.

Treated sawdust and other debris should be disposed of properly.

FOR EMERGENCY INFORMATION

(312) 900-1111