

1001-63

11/15/2013

1/21



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

NOV 15 2013

Carrie M. Tackema  
Cleary Chemical, LLC.  
4020 Aerial Center Parkway, Suite 101  
Morrisville, NC 27560

Subject: 3336 WP Turf and Ornamental Systemic Fungicide  
EPA Reg. No. 1001-63  
Notification dated October 17, 2013  
Decision Number 484522

Dear Ms. Tackema:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated October 17, 2013. The Registration Division (RD) has conducted a review of this request for applicability under PRN 98-10 and finds that your application falls within the scope of PRN 98-10. The inclusion of the marketing tag line on your product labeling is acceptable and the label will be placed in the file for our records.

If you have any questions, please call Marcel Howard at 703-305-6784 or email at [howard.marcel@epa.gov](mailto:howard.marcel@epa.gov).

Sincerely,

A handwritten signature in cursive script that reads "Shaja Joyner".

Shaja Joyner  
Product Manger (20)  
Fungicide Branch  
Registration Division (7504P)

2/21

EXPEDITE

	United States <b>Environmental Protection Agency</b> Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> <b>Other:</b>	OPP Identifier Number
---	---	--	-----------------------

**Application for Pesticide - Section I**

1. Company/Product Number <b>1001-63</b>	2. EPA Product Manager <b>Hope Johnson</b> <i>Shaja Jayar</i>	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) <b>3336 WP</b>	PM# <b>Acting 21 20</b>	
5. Name and Address of Applicant (Include ZIP Code) <b>Cleary Chemicals, LLC.</b> <b>4020 Aerial Center Parkway, Suite 101</b> <b>Morrisville, NC 27560</b> <u>Please send all correspondence to "contact point" listed below</u>  <input checked="" type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(l), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

**Section - II**

<input type="checkbox"/> Amendment – Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below

**Explanation:** Use additional page(s) if necessary. (For Section I and Section II.)


**NOTIFICATION PER PRN 98-10**

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

**CONTACT:** Carrie M. Tackema (919) 379-2528 or [carrie.tackema@us.nufarm.com](mailto:carrie.tackema@us.nufarm.com)  
 4020 Aerial Center Parkway, Suite 101  
 Morrisville, NC 27560

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input checked="" type="checkbox"/> Other foil/mylar	
*Certification must be submitted		If "Yes" Unit Packaging wgt.    No. per container	If "Yes" Package wgt.    No. per container		
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 4x8 oz.		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On labeling accompanying product	
6. Manner in Which Label is Affixed to Product				<input type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Other foil/mylar <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled	

**Section - IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application)		
Name <b>Carrie M. Tackema</b>	Title <b>Regulatory Manager</b>	Telephone No. (Include Area Code) <b>(919) 373-2528</b>
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received <b>(Stamped)</b>
2. Signature 	3. Title <b>Regulatory Manager</b>	
4. Typed Name <b>Carrie M. Tackema</b>	4. Date <b>October 17, 2013</b>	



Cleary Chemical LLC  
11901 S. Austin Avenue  
Alsip, IL 60803  
Phone: 708-377-1330 Fax: 708-377-1333

4020 Aerial Center Parkway, Suite 101  
Morrisville, NC 27560  
Phone: 919-379-2510 Fax: 919-467-5923  
www.us.nufarm.com

3/21

10/28

Via Courier Delivery

October 17, 2013

Ms. Hope Johnson, PM#21  
USEPA (7504P)  
Document Processing Desk (NOTIF)  
Room S4900, One Potomac Yard  
2777 S. Crystal Drive  
Arlington, VA 22202

Shaja Joyner, PM 20

**Subject: Notification per PRN 98-10  
EPA Reg. No(s). 1001-63; 1001-69; 1001-70; 1001-87**

Dear Ms. Johnson:

On behalf of Cleary Chemical, LLC, enclosed please find a Notification for each of the above-referenced products. The purpose of these Notifications is to include the marketing tag "Cleary|An Exclusive Line From Nufarm."

Also enclosed in support of this Notification are the following:

- Application for Pesticide Registration - NOTIFICATION;
- 1 copy of each of the above-referenced, proposed product labeling (in PDF format on CD-rom).

Should you have any questions or require additional information, please contact me at (919) 379-2528 or by email at [carrie.tackema@us.nufarm.com](mailto:carrie.tackema@us.nufarm.com).

Sincerely,

Carrie M. Tackema  
Regulatory Manager

Enclosure(s)

4/21

Master Label

# 3336 WP

## TURF AND ORNAMENTAL SYSTEMIC FUNGICIDE IN WATER SOLUBLE BAGS

Commercial Product, General Use: For Turf and Ornamental  
Applications Homeowner Product, General Use: Ornamental  
Applications Only

(Not subject to Worker Protection Standard)

For the prevention and control of turf diseases and the diseases of annual and  
perennial flowers, bedding plants, foliage plants, ground covers, plus deciduous and  
evergreen trees and shrubs.

ACTIVE  
INGREDIENT:

Thiophanate-methyl (dimethyl 4,4'-o-phenylenebis[3-thioallophanate])	....	50.00%
OTHER INGREDIENTS	.....	50.00%
TOTAL	.....	100.00%

**NET CONTENTS:** 2 Pounds (0.9 Kg)  
(Containing 4 – 8 ounce WATER SOLUBLE BAGS)

EPA Reg. No. 1001-63  
EPA Est. No. \_\_\_\_\_

**MANUFACTURED FOR**  
Cleary Chemicals, LLC  
11901 S. Austin Ave  
Alsip, IL 60803

**NOTIFICATION**

**NOV 15 2013**

5/21

Keep Out Of Reach Of Children

**CAUTION**

*[Optional referral statement for use with booklet label:  
See booklet label for additional precautionary statements and directions for use.]*

**FIRST AID**

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or when going for treatment.

For 24-Hour Emergency Assistance Call Chemtrec 1-800-424-9300

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Causes moderate eye irritation. Avoid contact eyes or clothing.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Natural rubber, polyvinyl chloride and polypropylene are chemical-resistant for this product. If you want more options, follow the instructions on an EPA chemical-resistance category selection chart.

Mixers, loaders, applicators and other handlers supporting dip treatment must wear: Coveralls over long sleeved shirt and long pants, Chemical-resistant gloves, Chemical-resistant footwear plus socks, A chemical-resistant apron. All other mixers, loaders, applicators and handlers must wear: Long sleeved shirt and long pants, Shoes plus socks, and Chemical-resistant gloves for all mixers and loaders and for applicators using handheld equipment.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

6/21

**USER SAFETY RECOMMENDATIONS:** User should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. User should remove PPE immediately after handling this product. As soon as possible wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. Do not apply, allow to drift, or drain or flush equipment onto non-target areas.

**DIRECTIONS FOR USE**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling, and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted- entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exemption: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls over long sleeved shirt and long pants, Chemical-resistant gloves made of any waterproof material, Chemical-resistant footwear plus socks, Chemical-resistant headgear for overhead exposure.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Turf and Landscape Uses:** Do not enter or allow others to enter the treated area until sprays have dried.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in the original container in a dry, temperature controlled area. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, contain/re-capture spillage and dispose of in accordance with the Pesticide Disposal Instructions listed below.

**Pesticide Disposal:** Wastes resulting from the use of this product are toxic and may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Securely wrap original container in several layers of newspaper and discard in trash. Offer for recycling, if available. When all water soluble bags are used, dispose of empty foil package in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Home Use Disposal:** If empty: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

**COMMERCIAL TURF AND ORNAMENTAL**

**GENERAL INFORMATION**

3336 WP is a broad spectrum fungicide exhibiting preventative, curative and systemic properties. It is useful on a wide variety of turf and ornamental disease problems. Apply 3336 WP with ground or overhead equipment, using sufficient volume of spray to provide thorough coverage. Do not apply with fixed wing or rotary aircraft. Use the higher rates under conditions of severe disease pressure. Also, see local State Extension Service recommendations for application schedules. This product is not for homeowner use and can only be used by individuals/firms licensed or registered by the state to apply ornamental or turf pest control products. Chemigation instructions follow "Directions For Use". Read and follow these instructions carefully for this method of application.

**Resistance Management:** To avoid the development of tolerant strains of fungi, 3336 WP should be used with fungicides of different modes of action. Cleary Chemical does not recommend the use of products containing thiabendazole in combination or rotation with 3336 WP. These utilize similar chemistry and mode of action and can contribute to development of disease tolerance. If, after using 3336 WP as recommended, the treatment is not effective, a tolerant strain of fungi may be present. Consult your local Cleary Chemical representative, your State Agricultural Experiment Station, or your State Cooperative Extension Service for proper disease identification and advice on the prompt use of some other suitable fungicide or disease control strategy. As long as recommended precautions are followed, 3336 WP can remain useful for disease control.

**Mixing Instructions:** 3336 WP consists of a protective outer resealable package containing water soluble bags. Do not allow the bags to become wet prior to adding to the tank. Do not handle water soluble bags with wet hands or wet gloves. Depending on the degree of agitation and water temperature, the water soluble bag should be completely dissolved within five minutes. Avoid adding bags near recirculation outlets as they might block outlets before completely dissolving. High pH environments cause a shortened tank life for suspended product. The buffering of tank water to pH 6-7 prior to the addition of 3336 WP is recommended. Add required amount of 3336 WP to a partially filled tank (1/2 total volume) and agitate by mechanical or hydraulic means while adding the remaining required amount of water. Once the water soluble bags have completely dissolved, add tank mix product if used (see below). Continuous agitation is recommended to keep the material in proper suspension. For best results, use spray mixture the same day it is prepared.

**Tank Mixing Instructions:** 3336 WP is compatible with most commonly used pesticides. If tank mixing with other materials, add products in the following order: water soluble bags, wettable powders, dry flowables, liquid flowables, emulsifiable concentrates, and soluble materials such as fertilizers. No claim of compatibility with other products is implied. **Do not tank mix with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur.** Consult the intended tank mix partner product label for appropriate application rates and use instructions. Follow the label directions for the most restrictive of label precautions and limitations. This product cannot be mixed with any product containing a label prohibition against such mixing.

Read and observe the most restrictive precautionary statements and other information appearing on product labels used in mixtures. 3336 WP may be applied in conjunction with chemically neutral liquid fertilizers. Application in conjunction with highly alkaline fertilizers, such as aqueous ammonia, may cause a degradation of the pesticide, resulting in reduced performance and should be avoided. Tank mix products containing boron or releasable free chlorine will affect the solubility of water soluble (PVA) film. Thoroughly rinse the spray tank of any boron containing spray solution prior to adding any water soluble bags. Even minimal carryover concentrations of boron in the spray tank may cause the water soluble bag material dissolved in water to precipitate and form insoluble residue inside the spray tank.

## CHEMIGATION

### Generic Requirements

1. Apply this product only through the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, solid set, or hand move; flood (basin); or drip trickle irrigation systems. Do not apply this product through any other type of irrigation system.
2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
3. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
5. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

9/21

### Specific Requirements

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back towards the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being drawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

### Non-Specific Requirements

1. Remove scale, pesticide residue, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.
2. Prepare a suspension of product in the mix tank or stock bucket. Fill the tank with 1/2 or 3/4 of the desired amount of water. Start agitation and add the required amount of product to the solution along with the remaining volume of water.
3. Maintain a gentle agitation in the mix tank during application to assure a uniform suspension. Follow mixing instructions and tank mixing instructions previously indicated.
4. Start system and then uniformly inject the suspension of 3336 WP into the irrigation line so as to deliver the desired rate per acre. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation system.
5. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time.
6. The suspension of 3336 WP should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing.

### Sprinkler (Overhead) Chemigation

Observe all instructions in the Generic, Specific and Non-Specific requirements sections above and the following additional requirements:

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Set sprinkler system to deliver 1/10 to 1/4 inches of water per acre. Volumes of water higher than this may reduce efficacy. Application of more than recommended quantities of irrigation water per acre may result in decreased product performance. Where sprinkler distribution patterns do not overlap sufficiently, unacceptable disease control may result. When system connections or fittings leak, when nozzles do not provide uniform distribution or when

10/21

lines containing the product cannot be flushed and must be dismantled and drained in a center pivot system, block the nozzle set nearest the well pivot injection unit to prevent spray being applied to this area. Allow sufficient time for pesticides to be flushed through all lines and all nozzles before turning off irrigation water.

**Flood (Basin) Chemigation**

1. Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and down stream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.

2. Systems utilizing a pressurized water and pesticide injection system must meet the following requirements: Observe all instructions in the Generic, Specific and Non-Specific requirements sections above and items 1 and 2 of the sprinkler irrigation requirements.

**Drip (Trickle) Chemigation**

Observe all instructions in the Generic, Specific and Non-Specific requirements sections above.

**COMMERCIAL TURF APPLICATIONS**

3336 WP may be used on all fine turf applications such as Commercial, Residential and Public (such as home lawns, parks, athletic fields, schools, and day care centers), and Golf Courses (greens, tees, fairways and aprons) of cool and warm season grasses such as Bentgrass, Bluegrass, Bermudagrass, Fescue, Ryegrass, St. Augustinegrass, Zoysiagrass, or their mixtures. 3336 WP is not phytotoxic to any of the above mentioned grasses when used in accordance with the label. 3336 WP is to be used for the prevention and control of the diseases mentioned below. It has both preventative and curative activity. Do not graze animals on treated turf. Do not feed clippings to livestock or poultry. Not for use on turf being grown for sale or other commercial use as sod.

**Application Instructions:** Apply material with properly calibrated hand held, mechanical or motorized spray equipment or by chemigation through appropriate sprinkler irrigation systems. Spray uniformly over the area to be treated. Apply recommended amounts in sufficient water to obtain thorough coverage of treatment area (2-4 gal per 1,000 sq ft is suggested). When treating golf greens, always treat aprons. Use the highest recommended rate under conditions of severe disease pressure. For best results, apply after mowing or avoid mowing twelve hours after application. For root pathogens, lightly water the treatment area to move the fungicide into active root zone with one to two tenths inch of water. Excessive irrigation may move application below active root zone and reduce application effectiveness. Green design and drainage will influence irrigation practices. When tank mixing with contact action fungicides for foliar diseases, applications should be allowed to dry on leaf surfaces. Normal watering may proceed after sprays have dried.

**Table 1: Maximum Individual Application Rates and Minimum Re-Treatment Intervals**

Use Site	Maximum Application Rate of 3336 WP	Minimum Re-treatment Interval	Comments
Residential or Public Areas	5.44 lbs/Acre (2 oz /1,000 sq.ft.)	14 days	
Golf Course Tees, Greens, Aprons	16.32 lbs/Acre (6 oz /1,000 sq.ft.)	14 days	
Golf Course Fairways -except Florida	10.88 lbs/Acre (4 oz /1,000 sq.ft.)	14 days	Excludes Florida

11/21

Golf Course Fairways- Florida Only	5.44 lbs/Acre (2 oz /1,000 sq.ft.)	14 days	Florida Only Only One Application Per Year
--	---------------------------------------	---------	--

**Table 2: Maximum Seasonal Application Rates**

Do not exceed the following amounts of product per Acre per season

Use Site	Maximum Pounds 3336 WP per Acre per Season	Ounces 3336 WP per 1,000 sq.ft.	Comments
Residential or Public Areas	21.76 lbs.	8oz.	4 Applications per year
Golf Course Tees, Greens, Aprons	43.52 lbs.	16 oz.	
Golf Course Fairways -except Florida	10.88 lbs.	4 oz.	Excludes Florida
Golf Course Fairways- Florida Only	5.44 lbs.	2 oz.	Florida Only

**Table 3: Turf Disease Control**

Observe the maximum individual application rates and maximum seasonal application limits in Table 1 and Table 2.

Disease(s) Controlled	Rate of 3336 WP oz/1,000 sq ft*	Remarks
<b>Anthracnose, basal</b> <i>Colletotrichum graminicola</i> <b>Anthracnose, foliar</b> <i>Colletotrichum graminicola</i>	4-6 2-4	For prevention in historic areas of disease pressure, apply twice at 14 day intervals when soil temperature reaches 60°F. For curative control, apply when disease first appears and continue at 14 day intervals. Rotations and/or tank mix combinations with chlorothalonil or triadimefon can be utilized.
<b>Bermudagrass Decline</b> <i>Gaeumannomyces graminis</i> var. <i>Graminis</i> <b>Take-All-Patch</b> <i>Gaeumannomyces graminis</i> var. <i>avenae</i>	4-6	Apply in mid-July or when disease symptoms first appear and repeat at 14 day intervals for suppression. Use higher rates under most severe disease expression. Follow proper agronomic recommendations to maintain plant vigor.
<b>Cool Season Brown Patch</b> <i>Rhizoctonia cerealis</i> <b>Necrotic Ring Spot</b> <i>Leptosphaeria korrea</i> <b>Spring Dead Spot</b> <i>Leptosphaeria korrea</i>	4-6	For prevention, apply in Fall before turf has stopped all growth activity. Apply second application in early Spring when soil temperatures reach 55-60°F or when disease first appears. For curative action, apply when disease first appears in early Spring and continue at 14 day intervals.
<b>Coprinus Snow Mold</b> <i>Coprinus psychromorbidus</i>	4-6	Apply 2 treatments at 21 day intervals in late Fall to early Winter, with the last application made just prior to first permanent snow cover. Rotations and/or tank mix combinations with PCNB can be utilized.
<b>Dollar spot</b> <i>Moellerodiscus, Lanzia,</i> <i>Sclerotinia</i> <b>Large Brown Patch</b> <i>Rhizoctonia solani</i> <b>Ascochyta Leaf Blight</b> <i>Ascochyta</i> <b>Copper Spot</b> <i>Gloeocercospora sorghi</i> <b>Fusarium Patch</b> <i>Fusarium nivale</i>	2-4	Apply when disease first appears and continue at 14 day intervals. Rotations and/or tank mix combinations with chlorothalonil, iprodione, or mancozeb (Protect™) can be utilized.

12/21

<b>Red Thread</b> <i>Laetisaria fuciformis</i>		
<b>Zoysia Patch</b> <i>Rhizoctonia solani</i>		
<b>Fusarium Blight</b> <i>Fusarium roseum, F. triticum</i>	4-6	Apply when disease first appears at 14 day intervals.
<b>Gray Leaf Spot (Blast)</b> <i>Pyricularia grisea</i>	4-6	Apply preventative application before expected period of disease development. Continue applications at 14 day intervals.
<b>Leaf Spot</b> <i>Drechslera</i> <b>Leaf, crown, and root diseases</b> <i>Bipolaris, Curvularia, Exserohilum</i>	4-6	Apply when disease first appears and make applications at 14 day intervals. Rotations and/or tank mix combinations with chlorothalonil, iprodione, or mancozeb (Protect™) are recommended.
<b>Pink Snow Mold</b> <i>Microdochium nivale</i>	2-4	Apply in late Fall to early Winter before turf has stopped all growth activity. A second application may be used in combination with chlorothalonil, PCNB, or thiram (Spotrete™) at recommended rates before snow cover or during Spring thaw.
<b>Rusts</b> <i>Puccinia, Uromyces</i>	4-6	Make 2 applications at 14 day intervals when disease first appears. Rotation and/or tank mix combinations with chlorothalonil or mancozeb (Protect) are recommended.
<b>Stripe Smut</b> <i>Ustilago striiformis</i>	4-6	Apply at 14 day intervals when disease first appears. For prevention, apply in spring and fall.
<b>Summer Patch</b> <i>Magnaporthe poae</i>	4-6	For prevention, apply 3 applications starting late April or early May using 21 day intervals. Rotations and/or tank mix combinations may be used as part of the three application program. For suppression, apply at 14 day intervals when disease first appears.
<b>Bentgrass Dead Spot</b> <i>Ophiosphaerella agrostis</i>	4-6	For prevention, apply in early June or based upon local Extension Service recommendations. Apply at 14 day intervals. Rotations and/or tank mix combinations may be used for season long disease prevention.

\*For residential or public areas do not apply more than 2 oz per 1,000 sq. ft. per application.

## COMMERCIAL HORTICULTURAL APPLICATIONS

**Nursery, Greenhouse, Shade or Lath House, Landscape & Interiorscape  
Annual and Perennial Flowers, Bedding Plants, Foliage Plants, Ground Covers,  
Vegetable Transplants, plus Deciduous and Evergreen Trees and Shrubs**

Do not apply to home orchards/fruit trees after fruit set.

Do not use fruit, nuts or sap from treated trees as food or feed.

3336 WP is a broad spectrum systemic fungicide which controls a variety of foliar, stem, and root diseases on a wide range of commercially important plants. 3336 WP is also effective as a pre-plant dip on cuttings and bulbs. For soil drench applications, best crop protection is achieved with preventative treatments repeated every 21-28 days. For foliar applications, begin treatments when disease first appears, or during suspected periods of disease incidence. Apply additional applications every 7-14 days or as otherwise instructed for the prevention or control of the listed diseases. Use of a wetting agent is recommended for plants that have leaves that are difficult to wet properly. Use of a spreader-sticker is recommended to enhance product performance in wet weather conditions or during periods of overhead irrigation. 3336 WP may be applied as a ground application using hand

13/21

held, mechanical or motorized spray equipment, or as a chemigation spray or through an applicable sprinkler irrigation system; or as an overhead application where applicable. See specific instructions below. For foliar applications other than cut flowers do not exceed three (3) pounds active ingredient per acre per single application. For cut flowers do not exceed one-half (0.5) pounds active ingredient per single application, Do not exceed thirty six (36) pounds active ingredient per acre per crop season from all thiophanate-methyl containing products. For soil drench applications, do not exceed 300 pounds active ingredient per acre per crop season from all thiophanate-methyl containing products.

**Note:** The "Directions For Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results.

This product is not recommended for use on Swedish Ivy (*Plectranthus australis*), Boston Fern (*Nephrolepis exhalta*), and Easter Cactus (*Hatiora gaertneri*).

**Application Instructions:** Apply material with properly calibrated, hand held, mechanical or motorized spray equipment or by chemigation thorough appropriate sprinkler irrigation, flood, or drip systems. Begin applications when disease first appears and repeat at 7-14 day intervals or as needed during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix 8-24 oz of 3336 WP per 100 gal water and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Spray volume may range up to 400 gallons of finished spray per acre depending upon plant species and plant growth stage. For applications through irrigation systems, refer to use rates indicated in the foliar application chart.

**Special Instructions For Proportional Injectors (e.g. Dosatron, Dosmatic, Anderson, and similar equipment)**

Determine the treatment rate for crop and pathogen from the foliar application table below. Determine the injection ratio for the individual system to be used for application.. For systems using a 1:100 ratio, measure and add the exact amount of recommended material per 100 gallons to each gallon of water in a stock bucket or tank. For systems using a 1:200 ratio, multiply the recommended amount per 100 gallons by 2. For systems using a 1:50 ratio, divide the recommended amount per 100 gallons added by 2. For systems using a 1:16 ratio, divide the recommended amount per 100 gallons by 6. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time. An injection ratio of 1:100 is recommended for most greenhouse and nursery systems.

Do not apply with fixed wing or rotary aircraft.

**FOLIAR APPLICATION:**

Diseases Controlled	Rate of 3336 WP	Remarks
<b>Anthraco</b> <b>nose</b>	12-16 oz/100 gallon	Apply as buds break or at first sign of disease. Repeat at 7-14 day intervals as needed during disease period.
<b>Black Spot of Rose</b> <i>Diplocarpon rosae</i>	12-16 oz/100 gallon	Apply late Spring or at first sign of disease. Repeat every 7-14 days as needed during disease period.

14/21

<b>Fusicladium and Venturia Leaf Scabs</b> on: Crabapple, Hawthorn, Pear, Mountain Ash, Pyracantha, etc.	12-16 oz/100 gallon	Apply as buds break. Repeat every 7-14 days as needed during disease period. Effective control requires coverage during leaf expansion. Rotations and/or tank mix combinations with mancozeb (Protect), chlorothalonil or propiconazole can be utilized. Do not use fruit from treated crabapple or pear trees for food purposes.
<b>Leaf Spots and Blights</b> caused by: <i>Ascochyta, Blumeriella, Botrytis, Cercospora, Coccomyces, Corynespora, Curvularia, Didymellina, Entomosporium, Fabraea, Fusarium, Ramularia, Rhizoctonia, Marssonina, Mycosphaerella, Myrothecium, Phoma, Physalospora, Schizothyrum, Septoria, Sphaceloma</i>	12-16 oz/100 gallon	Apply when disease symptoms first appear. Repeat every 7-14 days as needed during disease period. Rotations and/or tank mix combinations with mancozeb (Protect) or chlorothalonil can be utilized.
<b>Ovulinia Blight</b>	12-16 oz/100 gallon	Apply as flowers open. Repeat every 7-14 days as needed during disease period.
<b>Powdery Mildews</b> <i>Erysiphe, Microsphaera, Phyllactinia, Podosphaera, Oidium, Sphaerotheca</i>	12-24 oz/g100 gallon	Apply when disease first appears and repeat every 7-14 days during disease period. Rotations and/or tank mix combinations with mancozeb (Protect™) or triadimefon can be utilized.
<b>Rust Diseases</b> caused by: <i>Puccinia, Gymnosporagnium, Uromyces</i>	12-16 oz/100 gallon	Apply late Spring or when symptoms first appear. Repeat every 7-14 days as needed during disease period. Rotations and/or tank mix combinations with mancozeb (Protect) or chlorothalonil are recommended.
<b>Tip Blight of Pine</b> <i>Sphaeropsis sapinea, Diplodia pinea</i>	16-24 oz/100 gallon	Begin application in spring when new growth starts. Make a second application just before needles emerge from the sheath and a third application 14 days later. Thorough coverage is essential for optimal disease control.
<b>Twig Blights, Cankers, and Diebacks</b> <i>Diaporthe, Kabatina, Phoma, Phomopsis</i>	16-24 oz/100 gallon	Apply when symptoms first appear. Repeat every 7-14 days as needed during disease period.

**VEGETABLE TRANSPLANTS:  
Not intended for field vegetable production**

Vegetable Transplants	Diseases Controlled	Rate of 3336 WP	Remarks
<b>Beans, dry and succulent</b> including: Lima bean, Snap bean, Kidney bean, Mung bean, Navy bean, Pinto bean, Wax bean, Broad bean, Fava bean, Asparagus bean, Blackeyed pea, Cowpea, Sweet lupine, White lupine, White sweet lupine, Grain lupine, Chick pea, Garbanzo bean	<b>Anthracnose</b> <i>Colletotrichum</i>	2.1-2.8 lbs/Acre <sup>1</sup>	For one application: Apply when 100% of plants have at least one open bloom or when conditions are favorable for disease development.  or  For multiple applications: Make the first application when 10% to 30% of plants have at least one open bloom and follow with sequential applications on a 4 to 7 day interval. Apply prior to the development of disease for best results.  Do not apply more than 5.6 lbs. Of product (2.8 lbs a.i.)/A/season.
	<b>Gray Mold</b> <i>Botrytis</i>	or	
	<b>White Mold</b> <i>Sclerotinia</i>	1.4-2.1 lbs/Acre <sup>2</sup>	

15/21

			<p>Pre-harvest interval: <b>California only</b>, 14 days for succulent beans, 28 days for dry beans and lima beans.</p> <p>Pre-harvest interval: <b>all other States</b>, 14 days for succulent beans and lima beans, 28 days for dry beans.</p>
<b>CUCURBITS</b> (Cantaloupe, Casaba, Cucumbers, Melons, Pumpkins, Summer and Winter Squash, and Watermelons)	General Information		<p>Do not apply more than 4.2 lbs of product (2.1 lbs a.i.)/A/season from any combination of application timings.</p> <p>Follow resistance management guidelines under Directions for Use.</p>
	<b>Anthracnose*</b> <i>Colletotrichum</i> <b>Gummy Stem Blight*</b> <i>Didymella</i> <b>PowderyMildew*</b> <i>Erysiphe,</i> <i>Sphaerotheca,</i> <i>Podosphaera</i> <b>Target Spot*</b> <i>Corynespora</i>	0.7 lbs/Acre <sup>3</sup>	<p>Begin applications when plants begin to run or when disease first appears, and repeat at 7 to 14 day intervals.</p> <p>For Target Spot use at 7 day intervals.</p>
	<b>Belly Rots*</b> <i>Rhizoctonia,</i> <i>Fusarium</i>	0.7 lbs/Acre <sup>3</sup>	Apply in sufficient volume to allow runoff to the soil. Will not control <i>Pythium</i> or <i>Phytophthora</i> .

\*Not for this use in California.

1. Apply, for example, in 50-200 gallons of water per acre. In volumes of water below 50 gallons, use a minimum of 2.1 pounds per acre. If more than 200 gallons of water per acre are required for good plant coverage, apply a maximum rate of 2.8 pounds per acre. For example, if 200 gallons of water are required, use 1.4 pounds per 100 gallons.
2. Apply, for example, in 50-200 gallons of water per acre. In volumes of water below 50 gallons, use a minimum of 1.4 pounds per acre. If more than 200 gallons of water per acre are required for good plant coverage, apply a maximum rate of 2.1 pounds per acre. For example, if 200 gallons of water are required, use 1.05 pounds per 100 gallons.
3. Apply, for example, in 50-200 gallons of water per acre. In volumes of water below 50 gallons, use a minimum of 0.7 pounds per acre. If more than 200 gallons of water per acre are required for good plant coverage, apply a maximum rate of 0.7 pounds per acre. For example, if 200 gallons of water are required, use 0.35 pounds per 100 gallons.

**SOIL DRENCH APPLICATION**

Diseases Controlled	Rate of 3336 WP	Remarks
<b>Stem, Crown, and Root Rots</b> caused by: <i>Botrytis, Cyllindrocladium, Fusarium,</i> <i>Gliocladium, Myrothecium, Penicillium,</i> <i>Ramularia, Rhizoctonia, Sclerotinia</i>	8-16 oz/100 gal	<p>Apply as a drench or directed spray using hand held, mechanical, or motorized spray equipment, or as a chemigation drench or directed spray using applicable sprinkler irrigation systems, after seeding or sticking of cuttings (8 Oz) or after transplanting (12-16 oz) to propagation beds, containers, pots, trays, or nursery or landscape beds at a rate to thoroughly soak the growing media through the root zone. A general guide is 0.25-3 pints pf finished mixture per sq ft depending on the media type and depth (about 4 oz per 4 inch pot or 8 oz per 6 inch pot). Repeat every 21-</p>

16/21

		28 days for adequate crop protection. Note: 3336 WP does not control <i>Pythium</i> or <i>Phytophthora</i> . Tank mix combinations with metalaxyl, mefenoxam, etridiazole, fosetyl-A, mono and dipotassium salts of phosphorous acid or propamocarb are required for the control of <i>Pythium</i> and <i>Phytophthora</i> .
--	--	--

**PLANT DIP APPLICATION**

Diseases Controlled	Rate of 3336 WP	Remarks
<b>Plant or Cutting Diseases</b> caused by: <i>Botrytis, Cylindrocladium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia, Thielaviopsis</i>	16-24 oz/100 gal	Immerse plants or cuttings for 10-15 min. Remove and allow to drain. <b>Note:</b> Follow accepted hygiene practices to minimize the introduction and spread of water borne bacterial and water mold fungal diseases.
<b>Bulb, Corm, and Rhizome Rots</b> caused by: <i>Botrytis, Cylindrocladium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia, Thielaviopsis</i>	16-24 oz/100 gal	Soak cleaned bulgs for 15-30 min in warm solution (80-85°F). For storage disease prevention, treat bulbs preferably within 48 hours after digging. After treatment, dry well before storing. If bulbs are for forcing, treat bulbs that have been heat-cured. <b>Note:</b> Follow accepted hygiene practices to minimize the introduction and spread of water born bacterial and water mold fungal diseases.

**HOMEOWNER ORNAMENTAL**

**HOMEOWNER DIRECTIONS FOR USE**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or residential grounds.

For use on ornamentals only.

**HOMEOWNER GENERAL INFORMATION**

3336 WP is a broad-spectrum fungicide exhibiting preventative, curative and systemic properties. It is useful on a wide variety of ornamental disease problems. Apply 3336 WP with ground or overhead equipment, using sufficient volume of spray to provide thorough coverage..Use the higher rates under conditions of severe disease pressure. Also, see local State Extension Service recommendations for application schedules.

Do not allow people or pets to enter treated area until sprays have dried. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

**Resistance Management:** To avoid the development of tolerant strains of fungi, 3336 WP should be used with fungicides of different modes of action. Cleary Chemical does not recommend the use of products containing thiabendazole in combination or rotation with 3336 WP. These utilize similar chemistry and mode of action and can contribute to development of disease tolerance. If, after using 3336 WP as recommended, the treatment is not effective, a tolerant strain of fungi may be present. Consult your local Cleary Chemical representative, your State Agricultural Experiment Station, or your State Cooperative Extension Service for proper disease identification and advice on the prompt use of some other suitable fungicide or disease control strategy. As long as recommended precautions are

77/21

followed, 3336 WP can remain useful for disease control.

**Mixing Instructions:** 3336 WP consists of a loose powder inside a protective outer resealable package. Add required amount of 3336 WP to a partially filled tank (1/2 total volume) and agitate by mechanical means while adding the remaining required amount of water. Once the product has completely dissolved, add tank mix product if used (see below). **Do not tank mix 3336WP with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur.** High pH environments cause a shortened tank life for product. Buffer the tank water to pH 6-7 prior to the addition of 3336 WP. Cleary's NutriGrow Magnum may be used for pH buffering by adding one teaspoon per gallon. Continuous agitation is recommended to keep the material in proper suspension. For best results, use spray mixture the same day it is prepared.

**Tank Mixing Instructions:** 3336 WP is compatible with most commonly used pesticides. If tank mixing with other materials, add products in the following order: water soluble bags, wettable powders, dry flowables, liquid flowables, emulsifiable concentrates, and soluble materials such as fertilizers. No claim of compatibility with other products is implied. **Do not tank mix with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur (see Mixing Instructions).** Consult the intended tank mix partner product label for appropriate application rates and use instructions. Follow the label directions for the most restrictive of label precautions and limitations. This product cannot be mixed with any product containing a label prohibition against such mixing. Read and observe the most restrictive precautionary statements and other information appearing on product labels used in mixtures. 3336 WP may be applied in conjunction with chemically neutral liquid fertilizers. Application in conjunction with highly alkaline fertilizers, such as aqueous ammonia, may cause a degradation of the pesticide, resulting in reduced performance and should be avoided. Tank mix products containing boron or releasable free chlorine will affect the solubility of water soluble (PVA) film. Thoroughly rinse the spray tank of any boron containing spray solution prior to adding any water soluble bags. Even minimal carryover concentrations of boron in the spray tank may cause the water soluble bag material dissolved in water to precipitate and form insoluble residue inside the spray tank.

## HOMEOWNER HORTICULTURAL APPLICATIONS

**Annual and Perennial Flowers, Bedding Plants, Foliage Plants, Ground Covers, plus Deciduous and Evergreen Trees and Shrubs**

**Not intended for homeowner use on vegetables. Do not apply to home orchards/fruit trees.**

**Do not use fruit, nuts or sap from treated trees as food or feed.**

3336 WP is a broad spectrum systemic fungicide which controls a variety of foliar, stem, and root diseases on a wide range of commercially important plants. 3336 WP is also effective as a pre-plant dip on cuttings and bulbs. For soil drench applications, best crop protection is achieved with preventative treatments repeated every 21-28 days. For foliar applications, begin treatments when disease first appears, or during suspected periods of disease incidence. Apply additional applications every 7-14 days or as otherwise instructed for the prevention or control of the listed diseases. Use of a wetting agent is recommended for plants that have leaves that are difficult to wet properly. Use of a spreader-sticker is recommended to enhance product performance in wet weather conditions or during periods of overhead irrigation. 3336 WP may be applied as a ground application using hand held, mechanical or motorized spray equipment where applicable. See specific instructions below. For foliar and drench applications, do not exceed 1.8 pounds active ingredient per

18/21

acre (3.6 lbs of 3336 WP product) in a single application.

**Note:** The "Directions For Use" of this product reflects the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on Swedish Ivy (*Plectranthus australis*), Boston Fern (*Nephrolepis exalta*), and Easter Cactus (*Hatiora gaertneri*).

**Application Instructions**

Apply material with properly calibrated hand held, mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 7-14 day intervals or as needed during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications mix 1-4 teaspoons of 3336 WP per one (1) gal water and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Continuous agitation is required to keep product in suspension. The maximum application rate is 1.8 lbs ai per acre per application.

**FOLIAR APPLICATION**

Diseases Controlled	Rate of 3336 WP per gallon	Remarks
<b>Anthracnose</b>	2 ¼ - 3 teaspoons	Apply as buds break or at first sign of disease. Repeat at 7-14 day intervals as needed during disease period.
<b>Black Spot of Rose</b> <i>Diplocarpon rosae</i>	2 ¼ - 3 teaspoons	Apply early summer or at first sign of disease. Repeat every 7-14 days as needed during disease period.
<b>Brown Rot and Blight</b> <i>Monilinia, Sclerotinia, Whetzelinia</i>	2 ¼ - 3 teaspoons	Apply late Spring or at first sign of disease. Repeat every 7-14 days during disease period.
<b>Fusicladium and Venturia Leaf Scabs</b> on: Crabapple, Hawthorn, Pear, Mountain Ash, Pyracantha, etc.	2 ¼ - 3 teaspoons	Apply as buds break. Repeat every 7-14 days as needed during disease period. Effective control requires coverage during leaf expansion. Rotations and/or tank mix combinations with mancozeb (Protect), chlorothalonil or propiconazole can be utilized. Do not use fruit from treated crabapple or pear trees for food purposes.
<b>Leaf Spots and Blights</b> caused by: <i>Ascochyta, Blumeriella, Botrytis, Cercospora, Coccomyces, Corynespora, Curvularia, Didymellina, Entomosporium, Fabraea, Fusarium, Ramularia, Rhizoctonia, Marssonina, Mycosphaerella, Myrothecium, Phoma, Physalospora, Schizothyrium, Septoria, Sphaceloma</i>	2 ¼ - 3 teaspoons	Apply when disease symptoms first appear. Repeat every 7-14 days as needed during disease period. Rotations and/or tank mix combinations with mancozeb (Protect) or chlorothalonil can be utilized.
<b>Ovulinia Blight</b>	1 ½ - 3 teaspoons	Apply as flowers open. Repeat every 7-14 days as needed during disease period.
<b>Powdery Mildews</b> <i>Erysiphe, Microsphaera, Phyllactinia, Podosphaera, Oidium, Sphaerotheca</i>	2 ¼ - 4 ½ teaspoons	Apply when disease first appears and repeat every 7-14 days during disease period. Rotations and/or tank mix combinations with mancozeb (Protect™) or triadimefon can be utilized.
<b>Rust Diseases</b> caused by: <i>Puccinia, Gymnosporangium, Uromyces</i>	2 ¼ - 3 teaspoons	Apply late Spring or when symptoms first appear. Repeat every 7-14 days as needed during disease period. Rotations and/or tank mix combinations with mancozeb (Protect) or chlorothalonil are recommended.
<b>Tip Blight of Pine</b> <i>Sphaeropsis sapinea, Diplodia pinea</i>	3 - 4 ½ teaspoons	Begin application in spring when new growth starts. Make a second application just before

19/21

		needles emerge from the sheath and a third application 14 days later. Thorough coverage is essential for optimal disease control.
<b>Twig Blights, Cankers, and Diebacks</b> <i>Diaporthe, Kabatina, Phoma, Phomopsis</i>	3 – 4 ½ teaspoons	Apply when symptoms first appear. Repeat every 7-14 days as needed during disease period.

**SOIL DRENCH APPLICATION**

Diseases Controlled	Rate of 3336 WP per gallon	Remarks
<b>Stem, Crown, and Root Rots</b> caused by: <i>Botrytis, Cythindrocladium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia</i>	1 ½ - 3 teaspoons	Apply as a drench or directed spray using hand held, mechanical, or motorized spray equipment, or as a chemigation drench or directed spray using applicable sprinkler irrigation systems, after seeding or sticking of cuttings (8 Oz) or after transplanting (12-16 oz) to propagation beds, containers, pots, trays, or nursery or landscape beds at a rate to thoroughly soak the growing media through the root zone. A general guide is 0.25-3 pints pf finished mixture per sq ft depending on the media type and depth (about 4 oz per 4 inch pot or 8 oz per 6 inch pot). Repeat every 21-28 days for adequate crop protection. Note: 3336 WP does not control <i>Pythium</i> or <i>Phytophthora</i> . Tank mix combinations with metalaxyl, mefenoxam, etridiazole, fosetyl-A, mono and dipotassium salts of phosphorous acid or propamocarb are required for the control of <i>Pythium</i> and <i>Phytophthora</i> .
<b>Black Root Rot</b> <i>Thielaviopsis</i>		

**PLANT DIP APPLICATION**

Diseases Controlled	Rate of 3336 WP per gallon	Remarks
<b>Plant or Cutting Diseases</b> caused by: <i>Botrytis, Cythindrocladium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia, Thielaviopsis</i>	3 – 4 ½ teaspoons	Immerse plants or cuttings for 10-15 min. Remove and allow to drain. <b>Note:</b> Follow accepted hygiene practices to minimize the introduction and spread of water borne bacterial and water mold fungal diseases.
<b>Bulb, Corm, and Rhizome Rots</b> caused by: <i>Botrytis, Cythindrocladium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia, Thielaviopsis</i>	3 -4 ½ teaspoons	Soak cleaned bulbs for 15-30 min in warm solution (80-85°F). For storage disease prevention, treat bulbs preferably within 48 hours after digging. After treatment, dry well before storing. If bulbs are for forcing, treat bulbs that have been heat-cured. <b>Note:</b> Follow accepted hygiene practices to minimize the introduction and spread of water born bacterial and water mold fungal diseases.

**WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

**LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

**If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations of liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.**

Protect and Spotrete are registered trademarks of Cleary Chemicals LLC.

All other trademarks are the property of their respective owners.



AN EXCLUSIVE LINE FROM



21/21

**LABEL HISTORY**

<b>File Name</b>	<b>Revision Date</b>
001001-00063.draft080813	RV103009
001001-00063.master090513	RV090513
001001-00063.20131017.MASTER	RV101713