1001-65 Form Approved: OMB No. 2070-0060. Approval expires 2-28-95 Please read instructions on reverse before completing form. OPP Identifier Number Registration **United States Environmental Protection Agency** Amendment 214268 Washington, DC 20460 Other Application for Pesticide - Section I 1. Company/Product Number 2. EPA Product Manager 3. Proposed Classification 1001-65 CONNIE WELCH None Restricted 4. Company/Product (Name) 2 L PROTECT T/O 5. Name and Address of Applicant (Include ZIP Code) 6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) W.A. CLEARY CHEMICAL CORP. (b)(i), my product is similar or identical in composition and labeling 178 RIDGE ROAD, SUITE A EPA Reg. No. DAVION NJ 08810-1501 Check if this is a new address **Product Name** Section - II Final printed labels in repsonse to Amendment - Explain below. Agency letter dated Resubmission in response to Agency letter dated "Me Too" Application. Notification - Explain below. Other - Explain below. Explanation: Use additional page(s) if necessary. (For section I and Section II.) NOTIFICATION PER PRN 88-6. This is to certify that this Notification meets every requirement of PRN 88-6. Section - III 1. Material This Product Will Be Packaged in: Child-Resistant Packaging Unit Packaging Water Soluble Packaging 2. Type of Container Metal Yes Yes Yes Plastic No No No Glass Paner If "Yes" No. per If "Yes" No. per \* Certification must Unit Packaging wgt. container Package wgt Other (Specify) be submitted 3. Location of Net Contents Information 4. Size(s) Retail Container 5. Location of Label Directions Container 6. Manner in Which Label is Affixed to Product Lithograph Paper glued Stenciled Other Section - IV 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) Title . Name Telephone No. (Include Area Code) 329-8399 6. Date Application Received I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete.

I acknowledge that any knowlinglly false or misleading statement may be punishable by fine or imprisonment or

Richard F. Fletcher

EPA Form 8570-1 (Rev. 3-94) Previous editions are obsolete.

both under applicable law.

White - EPA File Copy (original)

Yellow - Applicant Copy

(Stamped)

**SPECIMEN LABEL** 

# PROTECT<sup>T/O</sup>

IN WATER SOLUBLE BAGS

# TURF AND ORNAMENTAL FUNGICIDE

An 80% Coordination Product of Manganese and Zinc Ethylenebisdithiocarbamate.

## **ACTIVE INGREDIENTS**

# KEEP OUT OF REACH OF CHILDREN

# CAUTION

## STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
- IF ON SKIN: Wash with plenty of soap and water. Get medical attention.
- IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS

## PRECAUTIONARY STATEMENTS-HAZARDS TO HUMANS AND DOMESTIC ANIMALS

May cause irritation of eyes. Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling.

## • PERSONAL PROTECTIVE EQUIPMENT (PPE)

APPLICATORS AND OTHER HANDLERS MUST WEAR: Long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, protective eyewear when mixing or loading, chemical-resistant apron when cleaning equipment, mixing or loading. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.


USER SAFETY RECOMMENDATIONS: Users should wash hands before: eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## • ENVIRONMENTAL HAZARD

This product is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

#### CHEMICAL HAZARDS

Do not allow to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of the product as a fungicide and create vapors which may be flammable. Keep container closed when not in use.



EPA REG. NO. 1001-65 EPA Est. No. 33967-NJ1

### STORAGE AND DISPOSAL

Storage: Store in the original container in a dry area. If allowed to become wet the product will deteriorate and represent a fire hazard. Keep away from sources of ignition (e.g. sparks and open flame). Close bag when not in use. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling sweep up spillage and dispose of in accordance with the Pesticide Disposal Instructions listed below.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Completely empty bag into application equipment. Dispose of bag in a sanitary landfill, by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or Tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific statements and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Personal Protective Equipment (PPE) required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over long-sleeved shirt and long pants, waterproof gloves and shoes plus socks.

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. TURF AND LANDSCAPE USES: Do not enter or allow entry into

TURF AND LANDSCAPE USES: Do not enter or allow entry into treated areas until sprays have dried unless wearing the personal protective equipment listed on this label.

## GENERAL INFORMATION

PROTECTT/OTERFAND ORNAMENTAL FUNGICIDE is a wettable powder containing a coordination product of zinc ion and manganese ethylenebisdithiocarbamate, and is recommended for use as a spray for the control of many important plant diseases. When used according to directions, it provides very high fungicidal activity and can be safely used on both turf grasses and certain ornamentals. PROTECTT/O provides control of most common turfgrass diseases and it is also effective in controlling major fungus diseases of certain ornamentals.

Diseases of this grass and ornamentals can attack suddenly and unexpectedly causing severe can age and may excirresult in total loss of large areas of valuable turf grass and ornamental plants. The use of a regular protective spray program will minimize the risk of disease damage and can generally be accomplished with lower rates and less frequent fungicide applications. Once

diseases have become established, higher rates of fungicide and more frequent applications are required to bring them under control. Follow a regular protective program for maximum product performance.

## INSTRUCTIONS FOR APPLICATION

TO MIX: Slowly add PROTECT T/O into spray tank while it is being filled and the agitator is running. Add insecticides and nutritional sprays last. PROTECT T/O is compatible with most commonly used pesticides. Read and observe the precautionary statements and other information appearing on product labels used in mixtures.

FOR AERIAL APPLICATION: Use at rate indicated in sufficient water for thorough coverage or a minimum of 2 gallons per acre. Use a spreader-sticker, such as Cleary's Clearspray T/O, at a rate of 4 fl.oz. per 100 gallon spray if needed. Add product slowly to water in the spray tank with agitation or premix thoroughly in separate holding tank for concentrate or aircraft sprayers. Continuous agitation is required to keep the product in suspension.

## **TURF APPLICATIONS**

Refer to General Use Instructions except when more specific directions are given for individual crops or diseases.

GENERAL USE: To be used on golf courses, sod farms and for professional applications to industrial, municipal and residential lawns. Do not use on pasture or range grasses. Do not feed clippings to livestock. Do not use if grass is grown for seed.

Start applications when grass greens up in the spring or when disease first appears and repeat at 7-14 day intervals or until disease threat is past. When conditions are especially favorable for disease development, apply maximum fungicide use rate on a 7 day spray schedule. Apply in sufficient water for adequate coverage 2-4 gallons are recommended for mechanical applications, smaller volumes are acceptable for hand applications. Do not graze treated areas. Do not feed clippings to livestock.

Rate of PROTECT T/O Oz./1000 Sq. Ft.	Remarks
4-8	
ıt, 4	Apply on a 7 day schedule for Rhizoctonia.
8	Apply at 5 day intervals or more frequently if conditions are especially favorable for disease development.
6-8	
6	
6-8	Apply at 2 week intervals during winter.
	PROTECT T/O Oz./1000 Sq. Ft. 4-8  at, 4  8  6-8 6

## HORTICULTURAL APPLICATIONS

## FIELD, NURSERY, GREENHOUSE, and LANDSCAPE

For use on the listed Annual and Perennial Flowers, Ground Covers, plus Deciduous and Evergreen Trees and Shrubs. Refer to General Use Instructions except when more specific directions are given for individual crops or diseases. Do not use edible portions of any listed plant for food or feed purposes.

GENERAL USE: Apply the equivalent of 1.5 lbs. PROTECT T/O per 100 gallons of water (2 water soluble bags) in full coverage spray. To improve performance, add 2-4 oz. spreader-sticker, such as Cleary's Clearspray T/O, per 100 gallons of spray. Begin spraying when plants are well leafed out or at first sign of disease and apply at 7-10 day intervals throughout the season, or follow State Extension Service recommendations for disease control on the listed ornamental plants. Note: The Directions for Use of this product reflect cumulative inputs from both field use experience and product testing programs. However, it is impossible to test this product on all plant species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied. Wait 5-7 days after treatment to evaluate trial results.

## Cleary's PROTECT T/O fungicide is recommended for use on the plant species listed below for the control of the following diseases and pathogens:

Alternaria - leaf spots and flower blights

Anthracnose Blights - Colletotrichum, Discula, Elsinoe,

Gloeodes, Gloeosporium, Glomerella, Gnomonia, Guignardia

Aschochyta Blights

Blackspot of Rose - Diplocarpon

Black Leafspot of Elm

Botryosphaeria Dieback

Botrytis Blights, Gray mold

Brown Rot Blight - Monolinia, Sclerotinia, Whetzelinia

Cylindrocladium Leaf Blight

Downy Mildews - Plasmopara, Peronospora

Flower Blights - Aschochyta, Botrytis, Monolinia, Sclerotinia,

Whetzelinia

Gray mold - Botrytis

Ink Spot - Ciborinia, Sclerotinia, Whetzelinia

Kabatina Dieback

Leafspots - Alternaria, Bipolaris, Cephalosporium, Cercospora,

Cylindrosporium, Dactylaria, Didymellina, Diplocarpon, Dreschleria, Exerohilum, Helminthosporium, Entomosporium, Fabraea, Leptosphaeria, Marssonina, Mycosphaerella, Pestalotia, Phyllosticta, Rhytisma, Septoria, Sphaeropsis, Stigmina

Needlecasts of Conifers - Cyclaneusma, Ismiella, Lophodermium, Naemacyclus, Rabdicline, Rhizosphaeria, Scirrhia

Phytophthora Dieback

Ramularia Blight

Rhizoctonia Web Blight

Rusts - Gymnosporangium, Melampsora, Melampsoridium,

Phragmidium, Puccinia, Uromyces

Scab - Fusicladium, Venturia, Sphaceloma, Cladosporium

Shothole Leafspot - Blumeriella, Coccomyces

Stemphyllium Blight

Tip Blights, Phomopsis, Diaporthe

Tip Blight of Pine - Diplodia, Sphaeropsis

Volutella Blight

## Ornamental plants for which Cleary's PROTECT T/O is recommended:

African violet - Saintpaulia

Ageratum - Ageratum

Almond (ornamental) - Prunus

Alyssum - Lobularia

Andromeda - Pieris

Anthurium - Anthurium

Apple (ornamental) - Malus sp.

Arborvitae - Thuia

Ash - Fraxinus

Ash, Mountain - Sorbus

Aspen - Poplus

Aster, China - Callistephus

Azalea\*- Rhododendron

Baby's Breath - Gysophyla

Begonia - Begonia

Birch - Betula

Boxwood - Buxus

Buckeye - Aescolus

Camellia\*- Camellia

Cardinal Flower - Lobelia

Carnation - Dianthus

Catharanthus - Catharanthus

Cedar, Red - Juniperus

Celosia - Celosia

Cherry (ornamental) - Prunus

Chinese evergreen - Aglaonema

Christmas Cactus - Schlumbergera

Christmas Trees

Chrysanthemum\*- Chrysanthemum Dendranthema

Coleus - Coleus

Columbine - Aquilegia

Conifers

Common Dracaena - Cordyline

Cotoneaster - Cotoneaster

Cyclamen - Cyclamen

Dahlia - Dahlia

Crape Myrtle - Lagerstroemia

Crabapple (ornamental) - Malus

Croton - Codiaeum

\* See "Special Instructions" on the next page for additional information on use.

Daisy, Transvall - Gerbera

Daisy, Shasta - Chrysanthemum

Delphinium - Delphinium

Dieffenbachia - Dieffenbachia

Dogwood\* - Comus

Dracaena - Dracaena

Dusty Miller - Centauria Elm - Ulmus

Euonymus - Euonymus

Fatsia - Fatsia

Ferns - Nephrolepus, Polypodium,

Polystichum, Pteris

Ficus - Ficus

Fig (ornamental) - Ficus

Firs - Pseudotsuga Firethorn - Pyracantha

Fittonia - Fittonia

Fuchsia - Fuchsia

Garden Balsam - Impatiens

Geranium - Geranium, Pelargonium

Gerbera - Gerbera

Gladiolus - Gladiolus

Gold Dust Tree - Aucuba

Gomphrena - Gomphrena

Gypsophila - Gypsophila Hawthorn - Crataegus

Hibiscus - Hibiscus

Hickory - Carva

Holly - Ilex

Hollyhock - Alcea

Horsechestnut Aescolus

Hydrangea - Hydrangea Impatiens - Impatiens

Iris - Iris

Ivy, Boston - Parthenocissus

Ivy, English - Hedera

Jadeplant - Crassula

Juniper - Juniperus

Kalanchoe - Kalanchoe

Larkspur - Delphinium

Laurel, Cherry - Prunus

Lavender Cotton - Santolina

Laurel, Mountain - Kalmia

Lily - Lilium

Liriope - Liriope Lobelia - Lobelia

Loquat - Eriobotrya

Magnolia - Magnolia

Mahonia - Mahonia

Maple - Acer Myrtle - Myrtus

Nasturtium - Tropaeolum

Nephthytis - Nephthytis

Nicotiana - Nicotiana Nierembergia - Nierembergia

Oak - Ouercus

Orchids - Cymbidium, Phalaenopsis

Osmanthus - Osmanthus

Pachysandra\*- Pachysandra

Palm, Areca - Chrysalidocarpus

Palm, King - Archotophoenix

Palm, Chamaedorea - Chamaedorea

Palm, Madagascar -Chrysalidocarpus

Palm, Washingtonia

Pansy - Viola

Peach (ornamental) - Prunus

Pear (ornamental) - Pyrus

Peony\*- Paeonia

Peperomia - Peperomia

Periwinkle - Vinca

Petunía - Petunia

Philodendron - Philodendron

Phlox - Phlox

Photinia - Red Tip

Pieris - Pieris

Pilea - Pilea

Pine - Pinus Pine, Norfolk Island - Araucaria

Pittosporum - Pittosporum

Plane Tree - Platanus

Plum (ornamental) - Prunus

Poinsettia - Euphorbia

Poplar - Populus

Portulaca - Portulaca

Pothos - Syngonium

Prayer Plant - Primula Privet - Ligustrum

Protea - Protea

Pyracantha - Pyracantha

Umbrella Tree - Schefflera

Red Tip - Photinia Redwood - Sequoia

Rhododendron\*- Rhododendron

Rose - Rosa

Sage - Salvia

Sand Cherry - Prunus

Santolina - Santolina

Schefflera - Schefflera

Senecio - Senecio

Sequoia - Sequoia

Snake Plant - Sansevieria Snapdragons - Antirrhinum

Spathiphylum - Spathiphylum

Spruce - Picea

Spurge, Japanese\*- Pachysandra

Statice - Limonium Strawflower - Helichrysum

Sycamore - Platanus

Syngonium - Syngonium

Vanda - Vanda

Velvetleaf - Abuulon. Verbena - Verbena

Viburnum - Viburnum

Wanter Juglans

Zinnia - Zinnia

Willow Salix

Yucca Yucca

Zebra Plant - Appelandra



535

# Special Instructions - Follow the Directions listed below for the specified plant and pathogen.

Azaleas, Camellias, Rhododendron - Petal Blight. Spray 2-3 times a week while flowers are opening. Direct spray into flowers and spray ground under plants thoroughly.

Chrysanthemum - Botrytis Petal Spot. Apply 1-2 lbs. per 100 gallons or 1.5 tablespoons per gallon in full coverage spray. Apply twice weekly during the blooming season.

Dogwood, Flowering - Anthracnose. Apply when buds begin to open, when bracts have fallen, 4 weeks later, and again in late summer after flower buds for next season are formed.

Pachysandra - Volutella Blight. Use a drenching spray of 4 lbs. per 100 gallons of water per 10,000 sq. ft. of bed area. Start application at first sign of disease and make at least 5 applications at 10-14 day intervals.

Peony - Phytophthora Blight, Botrytis Blight. Apply in early spring and early fall, drenching soil around plants as well as foliage. Promptly destroy all infected plant parts.

## **CHEMIGATION**

# APPLICATION THROUGH IRRIGATION SYSTEMS

#### **General Information**

- 1. Apply this product only through solid set or hand move, hand held, or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system.
- 2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- 3. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- 4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide labelprescribed safety devices for public water systems are in place.
- 5. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

# SPECIFIC INFORMATION FOR IRRIGATION SYSTEMS


#### Connected to Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

#### LIMITED WARRANTY AND DISCLAIMER

W. A. CLEARY CHEMICAL CORPORATION warrants that this material conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the Directions for Use, subject to the risks referred to therein. W. A. CLEARY CHEMICAL CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL W. A. CLEARY OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT INCLUDING BUT NOT LIMITED TO, LOSS OF PROFITS, BUSINESS REPUTATION. OF CUSTOMERS, LABOR COSTS, OR OTHER EXPENSES INCURRED IN PLANTING OR HARVESTING.

W. A. CLEARY and seller offer this product and the buyer and user accept it subject to the foregoing conflictions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of W. A. CLEARY CHEMICAL CORPORATION.



- 3. The pesticide injection pipeline must contain a functional, automatic, quickclosing check valve to prevent the flow of fluid back toward the injection pump.
- 4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 6. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- 7. Do not apply when wind speed favors drift beyond the area intended for treatment.

## SPRAY PREPARATION

- Remove scale, pesticide residues, and other foreign matter from the chemical stock tank and the entire injector system. Flush with clean water.
- 2. Follow the Directions for Use to determine the amount of PROTECT T/O that must be added to the stock tank. This amount should correspond with the site, host, and pathogen for which the product is used, and the intended amount of area to be treated.
- Fill the stock tank with approximately 1/2 the desired amount of water.Slowly add the required amount of PROTECTT/O to the stock tank, then add the remaining volume of water.
- 4. Maintain a gentle continuous agitation in the stock tank during the mixing and application process to ensure a uniform suspension of material.
- 5. PROTECT T/O may be applied in conjunction with chemically neutral liquid fertilizers and pesticides. Applications in conjunction with highly alkaline fertilizers or pesticides may cause a degradation of the pesticide resulting in reduced performance and should be avoided.
- Set sprinkler system to deliver 0.1-1.25 inches of water per acre. Volumes of water higher than this may reduce efficacy.
- 7. Start sprinkler and then uniformly inject the suspension of PROTECT T/O into the water line with a positive displacement pump ahead of a right angle turn to insure adequate mixing and the desire rate of product.



# W.A. Cleary Chemical

Corporate Headquarters 1049 Route 27, P.O. Box 10 Somerset, NJ 08875-0010 908-247-8000

Chemical Sales and Distribution Center 178 Ridge Road, Suite A Dayton, NJ 08810 908-329-8399 800-524-1662

Effective Date: 9/95 Replaces: 3/95

WAC43325P10W

