

1001-69

8/12/2009

1 of 23



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

AUG 12 2009

Mr. Pedro Perdomo
Cleary Chemicals, LLC
178 Ridge Road, Suite A
Dayton, NJ 08810

Subject: Label Notification(s) for Pesticide Registration Notice 2007-4

Dear Registrant:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated June 26, 2009 for:

EPA Registration 1001-69 3336 F

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Banza Djapao of my staff at 703-305-7269.

Sincerely,

Linda Arrington *for*
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

20923

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 1001-69	2. EPA Product Manager M. Waller	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) 3336 F	PM# 21	
5. Name and Address of Applicant (Include ZIP Code) Cleary Chemicals, LLC 178 Ridge Road, Suite A Dayton, NJ 08810 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
		If "Yes" Package wgt.	No. per container	<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Pedro Perdomo	Title Director - Regulatory Affairs	Telephone No. (Include Area Code) (732) 929-8399 x 1031
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Pedro Perdomo</i>	3. Title Director - Regulatory Affairs	
4. Typed Name Pedro Perdomo	5. Date June 26, 2009	

30f23



Cleary Chemical Corporation

800.524.1662

732.329.8399

Fax: 732.274.0894

178 Ridge Road

Dayton, NJ 08810

www.clearychemical.com

DATE: June 26, 2009

Mary Waller, Product Manager/Team 21
Registration Division - Document Processing Desk
USEPA - Office of Pesticide Programs (7504P)
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Dear Ms. Waller:


Cleary Chemicals, LLC is submitting a label notification to EPA, per EPA PR Notice 2007-4, to add container reuse and residue removal statements to the product label for 3336F (EPA Reg No. 1001-69).

Enclosed please find 1) transmittal document (this letter); 2) one copy of the completed form 8570-1; 3) one copy of the master label with changes highlighted; and 4) two final (clean) copies of the master label.

If you have any questions, please do not hesitate to contact me directly at 1.800.524.1662 ext. 1031.

Thank you.

Pedro Perdomo
Pedro Perdomo
Director - Regulatory



4023

Master Label

3336 F

TURF AND ORNAMENTAL SYSTEMIC FUNGICIDE

Commercial Product, General Use: For Turf and Ornamental Applications (May be subject to Worker Protection Standard)

Home Owner Product, General Use: Ornamental Applications Only

For the prevention and control of turf diseases and the diseases of annual and perennial flowers, bedding plants, foliage plants, ground covers, plus deciduous and evergreen trees and shrubs.

ACTIVE INGREDIENT:

Thiophanate-methyl (dimethyl 4,4'-o-phenylenebis[3-thioallophanate])	41.25%
OTHER INGREDIENTS	58.75%
TOTAL	100.00%


This product contains 4.0 lb thiophanate-methyl per gallon.

**Cleary Chemical LLC
178 Ridge Road, Suite A
Dayton, NJ 08810**

EPA Reg. No. 1001-69

Keep Out Of Reach Of Children

CAUTION



Optional referral statement for use with booklet label:
See booklet label for additional precautionary statements and directions for use.

FIRST AID

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.

IF SWALLOWED: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or when going for treatment.

For 24-Hour Emergency Assistance Call Chemtrec 1-800-424-9300

Cleary Chemical Corporation

178 Ridge Road, Suite A, Dayton, New Jersey 08810-1501

(732)-329-8399 (800)-524-1662

EPA Reg. No. 1001-69

EPA Est. No. 1812-GA-3

NET CONTENTS: 1 QUART, 1 GALLON, 2.5 GALLON

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if absorbed through skin. Harmful if inhaled. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category C on an EPA chemical resistance selection chart.

Handlers mixing, loading and applying the product as a dip must wear:

- Coveralls over long-sleeved shirt and long pants

- Chemical resistant gloves
- Chemical-resistant footwear plus socks
- Chemical-resistant apron.

All other mixers and loaders and applicators must wear:

- Long sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves
- Chemical-resistant apron for mixers, loaders and other handlers exposed to concentrate.

USER SAFETY REQUIREMENTS:

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS: User should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. User should remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling, and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exemption: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposures.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets out of treated areas until sprays have dried.

COMMERCIAL TURF AND ORNAMENTAL

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in the original container in a dry, temperature controlled area. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, contain/re-capture spillage and dispose of in accordance with the Pesticide Disposal instructions listed below.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal (Nonrefillable container 5 gallons or less): Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Offer for recycling, if available.

Residue Removal: Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

GENERAL INFORMATION

3336 F is a broad spectrum fungicide exhibiting preventative, curative and systemic properties. It is useful on a wide variety of turf and ornamental disease problems. Apply 3336 F with ground or overhead equipment, using sufficient volume of spray to provide thorough coverage.

Not for homeowner use. For use only by certified applicators or those under their immediate supervision. Do not apply with fixed wing or rotary aircraft. Not for use on turf being grown for sale or other commercial use as sod. Use the higher rates under conditions of severe disease pressure. Also, see local State Extension Service recommendations for application schedules. Chemigation instructions follow "Directions For Use". Read and follow these instructions carefully for this method of application.

Resistance Management: To avoid the development of tolerant strains of fungi, 3336 F should be used with fungicides of different modes of action. Cleary Chemical does not recommend the use of products containing thiabendazole in combination or rotation with 3336 F. These utilize similar chemistry and mode of action and can contribute to development of disease tolerance. If, after using 3336 F as recommended, and the treatment is not effective, a tolerant strain of fungi may be present. Consult your local Cleary Chemical representative, your State Agricultural Experiment Station, or your State Cooperative Extension Service for proper disease identification and advice on the prompt use of some other suitable fungicide or disease control strategy. As long as recommended precautions are followed, 3336 F can remain useful for disease control.

Mixing Instructions: SHAKE WELL BEFORE USING. Some settling may occur during prolonged periods of non-use. High pH environments cause a shortened tank life for diluted product. The buffering of tank water to pH 6-7 prior to the addition of 3336 F is recommended. Add required amount of 3336 F to partially filled tank (1/2 total volume), agitate by mechanical or hydraulic means, add tank mix product if used (**Do not tank mix 3336F with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur.** For more information, see 'Tank mixing instructions below), agitate again and then add remaining required amount of water. Continuous agitation is recommended to keep the material in proper suspension. For best results, use spray mixture the same day it is prepared.

Tank Mixing Instructions: 3336 F is compatible with most commonly used pesticides. If tank mixing with other materials, add products in the following order: water soluble bags, wettable powders, dry flowables, liquid flowables, emulsifiable concentrates, and soluble materials such as fertilizers. No claim of compatibility with other products is implied. **Do not tank mix with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur.** Consult the intended tank mix partner product label for appropriate application rates and use instructions. Follow the label directions for the most restrictive of label precautions and limitations. This product cannot be mixed with any product containing a label prohibition against such mixing. Read and observe the most restrictive precautionary statements and other information appearing on product labels used in mixtures. 3336 F may be applied in conjunction with chemically neutral liquid fertilizers. Application in conjunction with highly alkaline fertilizers, such as aqueous ammonia, may cause a degradation of the pesticide, resulting in reduced performance and should be avoided.

CHEMIGATION

Generic Requirements

1. Apply this product only through the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, solid set, or hand move; flood (basin); or drip trickle irrigation systems. Do not apply this product through any other type of irrigation system.
2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
3. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
5. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Specific Requirements

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back towards the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being drawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Non-Specific Requirements

1. Remove scale, pesticide residue, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.
2. Prepare a suspension of product in the mix tank or stock bucket. Fill the tank with 1/2 or 3/4 of the desired amount of water. Start agitation and add the required amount of product to the solution along with the remaining volume of water.
3. Maintain a gentle agitation in the mix tank during application to assure a uniform suspension. Follow mixing instructions and tank mixing instructions previously indicated.
4. Start system and then uniformly inject the suspension of 3336 F into the irrigation line so as to deliver the desired rate per acre. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation system.
5. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time.
6. The suspension of 3336 F should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing.

Sprinkler (Overhead) Chemigation

Observe all instructions in the Generic, Specific and Non-Specific requirements sections above and the following additional requirements:

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Set sprinkler system to deliver 1/10 to 1/4 inches of water per acre. Volumes of water higher than this may reduce efficacy. Application of more than recommended quantities of irrigation water per acre may result in decreased product performance. Where sprinkler distribution patterns do not overlap sufficiently, unacceptable disease control may result.

When system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product cannot be flushed and must be dismantled and drained in a center pivot system, block the nozzle set nearest the well pivot injection unit to prevent spray being applied to this area. Allow sufficient time for pesticides to be flushed through all lines and all nozzles before turning off irrigation water.

Flood (Basin) Chemigation

1. Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and down stream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.
2. Systems utilizing a pressurized water and pesticide injection system must meet the following requirements: Observe all instructions in the Generic, Specific and Non-Specific requirements sections above and items 1 and 2 of the sprinkler irrigation requirements.

Drip (Trickle) Chemigation

Observe all instructions in the Generic, Specific and Non-Specific requirements sections above.

COMMERCIAL TURF APPLICATIONS

3336 F may be used on all fine turf applications such as Commercial, Residential and Public (such as home lawns, parks, athletic fields, schools, and day care centers), and Golf Courses (greens, tees, fairways, and aprons) of cool and warm season grasses such as Bentgrass, Bluegrass, Bermudagrass, Fescue, Ryegrass, St. Augustinegrass, Zoysiagrass, or their mixtures. 3336 F is not phytotoxic to any of the above mentioned grasses when used in accordance with the label. 3336 F is to be used for the prevention and control of the diseases mentioned below. It has both preventative and curative activity. Do not graze animals on treated turf. Do not feed clippings to livestock or poultry.

Application Instructions: Apply material with properly calibrated hand held, mechanical or motorized spray equipment or by chemigation through appropriate sprinkler irrigation systems. Spray uniformly over the area to be treated. Apply recommended amounts in sufficient water to obtain thorough coverage of treatment area (2-4 gal per 1,000 sq ft is suggested). When treating golf greens, always treat aprons. Use the highest recommended rate under conditions of severe disease pressure. For best results, apply after mowing or avoid mowing twelve hours after application. For root pathogens, lightly water the treatment area to move the fungicide into the active root zone with one to two tenths inch of water. Excessive irrigation may move application below active root zone and reduce application effectiveness. Green design and drainage will influence irrigation practices. When tank mixing with contact action fungicides for foliar diseases, applications should be allowed to dry on leaf surfaces. Normal watering may proceed after sprays have dried.

Table 1: Maximum Individual Application Rates and Minimum Re-Treatment Intervals

Do not exceed the amounts per acre or reduce the re-treatment interval indicated below.

Use Site	Maximum application rate of 3336 F	Minimum Re-Treatment Interval	Comments
Residential or Public Areas	0.68 Gallons/Acre (2 fl. oz / 1,000 sq.ft.)	14 days	
Golf Course Tees, Greens, Aprons	2.04 Gallons/Acre (6 fl. oz / 1,000 sq.ft.)	14 days	
Golf Course Fairways –except Florida	1.36 Gallons/Acre (4 fl. oz / 1,000 sq.ft.)	14 days	Excludes Florida
Golf Course Fairways – Florida Only	0.68 Gallons/Acre (2 fl. oz / 1,000 sq.ft.)	14 days	Florida Only.

Table 2: Maximum Yearly Application Rates

Do not exceed the following amounts of product per Acre per year

Use Site	Maximum Gallons 3336F per Acre per Season	Fluid Ounces 3336 F per 1,000 sq.ft.	Comments
Residential or Public Areas	2.72 Gallons	8 fl.oz.	4 Applications per year
Golf Course Tees, Greens, Aprons	5.44 Gallons	16 fl. oz.	
Golf Course Fairways –except Florida	1.36 Gallons	4 fl.oz.	Excludes Florida
Golf Course Fairways – Florida Only	0.68 Gallons	2 fl.oz.	Florida Only

Table 3: Turf Disease Control

Disease(s) Controlled	Rate of 3336 F fl oz/1,000 sq ft	Remarks*
Anthracnose, basal <i>Colletotrichum graminicola</i>	4-6	For prevention in historic areas of disease pressure, apply twice at 14 day intervals when soil temperature reaches 60°F. For curative control, apply when disease first appears and continue at 14 day intervals. Rotations and/or tank mix combinations with chlorothalonil or triadimefon can be utilized.
Anthracnose, foliar <i>Colletotrichum graminicola</i>	2-4	
Bermudagrass Decline <i>Gaeumannomyces graminis</i> var. <i>graminis</i> Take-All-Patch <i>Gaeumannomyces graminis</i> var. <i>avenae</i>	4-6	Apply in mid-July or when disease symptoms first appear and repeat at 14 day intervals for suppression. Use higher rates under most severe disease expression. Water treatment into active root zone. Follow proper agronomic recommendations to maintain plant vigor.
Cool Season Brown Patch <i>Rhizoctonia cerealis</i> Necrotic Ring Spot <i>Leptosphaeria korrea</i> Spring Dead Spot <i>Leptosphaeria korrea</i>	4-6	For prevention, apply in Fall before turf has stopped all growth activity. Apply second application in early Spring when soil temperatures reach 55-60°F or when disease first appears. For curative action, apply when disease first appears in early Spring and continue at 14 day intervals. Water treatment into active root zone.
Coprinus Snow Mold <i>Coprinus psychromorbidus</i>	4-6	Apply 2 treatments at 21 day intervals in late Fall to early Winter, with the last application made just prior to first permanent snow cover. Rotations and/or tank mix combinations with PCNB can be utilized.
Dollar Spot <i>Moellerodiscus, Lanzia,</i> <i>Sclerotinia</i> Large Brown Patch <i>Rhizoctonia solani</i> Ascochyta Leaf Blight <i>Ascochyta</i> Copper Spot <i>Gloeocercospora sorghi</i> Fusarium Patch <i>Fusarium nivale</i> Red Thread <i>Laetisaria fuciformis</i> Zoysia Patch <i>Rhizoctonia solani</i>	2-4	Apply when disease first appears and continue at 14 day intervals. Rotations and/or tank mix combinations with chlorothalonil, iprodione, or mancozeb (Protect™) can be utilized.
Fusarium Blight <i>Fusarium roseum, F. triticum</i>	4-6	Apply when disease first appears at 14 day intervals.
Gray Leaf Spot (Blast) <i>Pyricularia grisea</i>	4-6	Apply preventative application before expected period of disease development. Continue applications at 14 day intervals.
Leaf Spot <i>Drechslera</i> Leaf, Crown, and Root	4-6	Apply when disease first appears and make applications at 14 day intervals as needed. Rotations and/or tank mix combinations with chlorothalonil,

Diseases <i>Bipolaris, Curvularia, Exserohilum</i>		iprodisone, or mancozeb (Protect) are recommended under severe conditions.
Pink Snow Mold <i>Microdochium nivale</i>	2-4	Apply in late Fall to early Winter before turf has stopped all growth activity. A second application may be used in combination with chlorothalonil, PCNB, or thiram (Spotrete™) at recommended rates before snow cover or during Spring thaw.
Rusts <i>Puccinia, Uromyces</i>	4-6	Apply at 14 day intervals when disease first appears. Rotations and/or tank mix combinations with chlorothalonil or mancozeb (Protect) are recommended.
Stripe Smut <i>Ustilago striiformis</i>	4-6	Apply at 14 day intervals when disease first appears. For prevention, apply in spring and fall.
Summer Patch <i>Magnaporthe poae</i>	4-6	For prevention, apply 3 applications starting late April or early May using 21 day intervals. Rotations and/or tank mix combinations may be used as part of the three application program. For suppression, apply at 14 day intervals when disease first appears. Water treatment into active root zone.
Bentgrass Dead Spot <i>Ophiosphaerella agrostis</i>	4-6	For prevention, apply in early June or based upon local Extension Service recommendations. Apply at 14 day intervals. Rotations and/or tank mix combinations may be used for season long disease prevention.

***Observe the maximum individual application rates and maximal seasonal applications limits in Table 2.**

COMMERCIAL HORTICULTURAL APPLICATIONS

Nursery, Greenhouse, Landscape & Interiorscape

Annual and Perennial Flowers, Bedding Plants, Foliage Plants, Ground Covers, plus Deciduous and Evergreen Trees and Shrubs

Do not use fruits, nuts or sap from treated trees as food or feed.

3336 F is a broad spectrum systemic fungicide which controls a variety of foliar, stem, and root diseases on a wide range of commercially important plants. 3336 F is also effective as a pre-plant dip on cuttings and bulbs. For soil drench applications, best crop protection is achieved with preventative treatments repeated every 21-28 days. For foliar applications, begin treatments when disease first appears, or during suspected periods of disease incidence. Apply additional applications every 7-14 days or as otherwise instructed for the prevention or control of the listed diseases. Use of a wetting agent is recommended

for plants that have leaves that are difficult to wet properly. Use of a spreader-sticker is recommended to enhance product performance in wet weather conditions or during periods of overhead irrigation. 3336 F may be applied as a ground application using hand held, mechanical or motorized spray equipment, or as a chemigation spray or through an applicable sprinkler irrigation systems; or as an overhead application where applicable. See specific instructions below.

Note: The "Directions For Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on Swedish Ivy (*Plectranthus australis*), Boston fern (*Nephrolepis exhalta*), and Easter Cactus (*Hatiora gaertneri*).

Application Instructions: Apply material with properly calibrated hand held, mechanical or motorized spray equipment or by chemigation thorough appropriate sprinkler irrigation, flood, or drip systems. Begin applications when disease first appears and repeat at 7-14 day intervals or as needed during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix 8-24 fl oz of 3336 F per 100 gal water (0.5-1.5 teaspoons per gal) and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Spray volume may range up to 400 gallons of finished spray per acre depending upon plant species and plant growth stage. For applications through irrigation systems, refer to use rates indicated in the foliar application chart. For small volume applications less than 100 gallons, divide recommended rate by 16 to get the number of teaspoons of 3336 F per gal.

Special Instructions For Proportional Injectors (e.g. Dosatron, Dosmatic, Anderson, and similar equipment)

Determine the treatment rate for crop and pathogen from the foliar application table below. Determine the injection ratio for the individual system to be used for application. For systems using a 1:100 ratio, measure and add the exact amount of recommended material per 100 gallons to each gallon of water in a stock bucket or tank. For systems using a 1:200 ratio, multiply the recommended amount per 100 gallons by 2. For systems using a 1:50 ratio, divide the recommended amount per 100 gallons added by 2. For systems using a 1:16 ratio, divide the recommended amount per 100 gallons by 6. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time. An injection ratio of 1:100 is recommended for most greenhouse and nursery systems. Do not apply more than 3.0 lbs AI per year.

FOLIAR APPLICATION*

Diseases Controlled	Rate of 3336 F	Remarks
---------------------	----------------	---------

Anthracnose <i>Colletotrichum</i>	12-16 fl oz/100 gallon	Apply as buds break or at first sign of disease. Repeat at 7-14 day intervals as needed during disease period.
Black Spot of Rose <i>Diplocarpon rosae</i>	12-16 fl oz/100 gallon	Apply early summer or at first sign of disease. Repeat every 7-14 days as needed during disease period.
Brown Rot and Blight <i>Monilinia, Sclerotinia, Whetzellinia</i>	12-16 fl oz/100 gallon	Apply late Spring or at first sign of disease. Repeat every 7-14 days as needed during disease period.
Fusicladium and Venturia Leaf Scabs on: Crabapple, Hawthorn, Pear, Mountain Ash, Pyracantha, etc.	12-16 fl oz/100 gallon	Apply as buds break. Repeat every 7-14 days during disease period. Effective control requires coverage during leaf expansion. Rotations and/or tank mix combinations with mancozeb (Protect), chlorothalonil or propiconazole can be utilized. Do not use fruit from treated crabapple or pear trees for food purposes.
Leaf Spots and Blights caused by: <i>Ascochyta, Blumeriella, Botrytis, Cercospora, Coccoomyces, Corynespora, Curvularia, Didymellina, Entomosporium, Fabraea, Fusarium, Ramularia, Rhizoctonia, Marssonina, Mycosphaerella, Myrothecium, Phoma, Physalospora, Schizothyrium, Septoria, Sphaceloma</i>	12-16 fl oz/100 gallon	Apply when disease symptoms first appear. Repeat every 7-14 days during disease period. Rotations and/or tank mix combinations with mancozeb (Protect) or chlorothalonil can be utilized.
Ovulinia Blight	8-16 fl oz/100 gallon	Apply as flowers open. Repeat every 7-14 days during disease period.
Powdery Mildews <i>Erysiphe, Microsphaera, Phyllactinia, Podosphaera, Oidium, Sphaerotheca</i>	12-24 fl oz/100 gallon	Apply when disease first appears and repeat every 7-14 days during disease period. Rotations and/or tank mix combinations with mancozeb (Protect) or triadimefon can be utilized.
Rust Diseases caused by: <i>Puccinia, Gymnosporangium, Uromyces</i>	12-16 fl oz/100 gallon	Apply late Spring or when symptoms first appear. Repeat every 7-14 days during disease period. Rotations and/or tank mix combinations with mancozeb (Protect) or chlorothalonil are recommended.
Tip Blight of Pine <i>Sphaeropsis sapinea, Diplodia pinea</i>	16-24 fl oz/100 gallon	Begin application in spring when new growth starts. Make a second application just before needles emerge from the sheath and a third application 14 days later. Thorough coverage is essential for optimal disease control.
Twig Blights, Cankers, and Diebacks <i>Diaporthe, Kabatina, Phoma, Phomopsis</i>	16-24 fl oz/100 gallon	Apply when symptoms first appear. Repeat every 7-14 days during disease period.

* For Cut Flowers do not exceed 0.5 lbs ai/acre/application

SOIL DRENCH APPLICATION

Diseases Controlled	Rate of 3336 F	Remarks
Stem, Crown, and Root Rots caused by: <i>Botrytis, Cythindrocladium, Fusarium,</i> <i>Gliocladium, Myrothecium, Penicillium,</i> <i>Ramularia, Rhizoctonia, Sclerotinia</i> Black Root Rot <i>Thielaviopsis</i>	8-16 fl oz/100 gal	Apply as a drench or directed spray using hand held, mechanical or motorized spray equipment or as a chemigation drench or directed spray using applicable sprinkler irrigation systems after seeding or sticking of cuttings (8 fl oz) or after transplanting (12-16 fl oz) to propagation beds, containers, pots, trays, or nursery or landscape beds at a rate to thoroughly soak the growing media through the root zone. A general guide is 0.25 –3.0 pints of finished mixture per sq ft depending on the media type and depth (about 4 fl oz per 4 inch pot or 8 fl oz per 6 inch pot). Repeat every 21-28 days for adequate crop protection. Note: 3336 F does not control <i>Pythium</i> or <i>Phytophthora</i> . Tank mix combinations with metalaxyl, mfenoxam, etridiazole, propamocarb, fosetyl-AI or mono and dipotassium salts of phosphorous acid are required for the control of <i>Pythium</i> and <i>Phytophthora</i> .

PLANT DIP APPLICATION

Diseases Controlled	Rate of 3336 F	Remarks
Plant or Cutting Diseases caused by: <i>Botrytis, Cythindrocladium, Fusarium,</i> <i>Gliocladium, Myrothecium, Penicillium,</i> <i>Ramularia, Rhizoctonia, Sclerotinia,</i> <i>Thielaviopsis</i>	16-24 fl oz/100 gal	Immerse plants or cuttings for 10-15 min. Remove and allow to drain. Note: Follow accepted hygiene practices to minimize the introduction and spread of water borne bacterial and water mold fungal diseases.
Bulb, Corm, and Rhizome Rots caused by: <i>Botrytis, Cythindrocladium, Fusarium,</i> <i>Gliocladium, Myrothecium, Penicillium,</i> <i>Ramularia, Rhizoctonia, Sclerotinia,</i> <i>Thielaviopsis</i>	16-24 fl oz/100 gal	Soak cleaned bulbs for 15-30 min in warm solution (80-85° F). For storage disease prevention, treat bulbs preferably within 48 hours after digging. After treatment, dry well before storing. If bulbs are for forcing, treat bulbs that have been heat-cured. Note: Follow accepted hygiene practices to minimize the introduction and spread of water borne bacterial and water mold fungal diseases.

HOMEOWNER ORNAMENTAL

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of equipment wash water.

Do not apply this product in a way that will contact any person, or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in the original container in a dry, temperature controlled area. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, contain/re-capture spillage and dispose of in accordance with the Pesticide Disposal instructions listed below.

Container Disposal:

If empty: Do not re-use empty container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency [or 1-800-CLEANUP or other appropriate number – see criteria below] for disposal instructions. Never place unused product down any indoor or outdoor drain.]

HOMEOWNER DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or residential grounds.

For use on ornamentals only.
Do not apply to home orchard/fruit trees.

HOMEOWNER GENERAL INFORMATION

3336 F is a broad-spectrum fungicide exhibiting preventative, curative and systemic properties. It is useful on a wide variety of ornamental disease problems. Apply 3336 F with ground or overhead equipment, using sufficient volume of spray to provide thorough coverage. Use the higher rates under conditions of severe disease pressure. Also, see local State Extension Service recommendations for application schedules. Do not apply this product in a way that will contact any person, or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.

Resistance Management: To avoid the development of tolerant strains of fungi, 3336 F should be used with fungicides of different modes of action. Cleary Chemical does not recommend the use of products containing thiabendazole in combination or rotation with 3336 F. These utilize similar chemistry and mode of action and can contribute to development of disease tolerance. If, after using 3336 F as recommended, the treatment is not effective, a tolerant strain of fungi may be present. Consult your local Cleary Chemical representative, your State Agricultural Experiment Station, or your State Cooperative Extension Service for proper disease identification and advice on the prompt use of some other suitable fungicide or disease control strategy. As long as recommended precautions are followed, 3336 F can remain useful for disease control.

Mixing Instructions: SHAKE WELL BEFORE USING. Some settling may occur during prolonged periods of non-use.

Add required amount of 3336 F to partially filled tank (1/2 total volume), agitate by mechanical means, and then add remaining required amount of water. Continuous agitation is recommended to keep the material in proper suspension. Once the product has completely dissolved, add tank mix product if used (see below). ***Do not tank mix 3336F with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur. High pH environments cause a shortened tank life for product. Buffer the tank water to pH 6-7 prior to the addition of 3336 F. Cleary's NutriGrow Magnum may be used for pH buffering by adding one teaspoon per gallon.*** Use spray mixture the same day it is prepared.

Tank Mixing Instructions: 3336 F is compatible with most commonly used pesticides. If tank mixing with other materials, add products in the following order: water soluble bags, wettable powders, dry flowables, liquid flowables, emulsifiable concentrates, and soluble materials such as fertilizers. No claim of compatibility with other products is implied. ***Do not tank mix with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur (see Mixing Instructions).*** Consult the intended tank mix partner product label for appropriate application rates and use instructions. Follow the label directions for the most restrictive of label precautions and limitations. This product cannot be mixed with any product containing a label prohibition against such mixing. Read and observe the most restrictive precautionary statements and other information appearing on product labels used in mixtures. 3336 F may be applied in conjunction with chemically neutral liquid fertilizers. Application in conjunction with highly alkaline fertilizers, such as aqueous ammonia, may cause a degradation of the pesticide, resulting in reduced performance and should be avoided.

HOMEOWNER HORTICULTURAL APPLICATIONS

Nursery, Greenhouse, Annual and Perennial Flowers, Bedding Plants, Foliage Plants, Ground Covers, plus Deciduous and Evergreen Trees and Shrubs

For use on ornamentals only. Do not apply to home orchard/fruit trees.

3336 F is a broad-spectrum systemic fungicide, which controls a variety of foliar, stem, and root diseases on a wide range of commercially important plants. 3336 F is also effective as a pre-plant dip on cuttings and bulbs. For soil drench applications, best crop protection is achieved with preventative treatments repeated every 21-28 days. For foliar applications, begin treatments when disease first appears, or during suspected periods of disease incidence. Apply additional applications every 7-14 days or as otherwise instructed for the prevention or control of the listed diseases. Use of a wetting agent is recommended for plants that have leaves that are difficult to wet properly. Use of a spreader-sticker is recommended to enhance product performance in wet weather conditions or during periods of overhead irrigation. **Note:** The "Directions For Use" of this product reflects the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on Swedish Ivy (*Plectranthus australis*), Boston Fern (*Nephrolepis exaltata*), and Easter Cactus (*Hatiora gaertneri*).

Application Instructions: Apply material with properly calibrated spray equipment. Begin applications when disease first appears and repeat at 7-14 day intervals or as needed during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. Apply 0.5-1.5 teaspoons of 3336 F per 1 gallon water as a full coverage spray to drip for the prevention and control of the diseases listed below. Typical residential spray volume is 150 gallons per acre. Continuous agitation is required to keep product in suspension. For Foliar applications, do not exceed 1.8 pounds active ingredient (1.8 quarts product) per treated acre in a single application.

FOLIAR APPLICATION

Diseases Controlled	Rate of 3336 F per gallon	Remarks
Anthracnose <i>Colletotrichum</i>	½ – 1 ½ teaspoons	Apply as buds break or at first signs of disease. Repeat at 7-14 day intervals as needed during disease period.
Black Spot of Rose <i>Diplocarpon rosae</i>	½ – 1 ½ teaspoons	Apply early summer or at first sign of disease. Repeat every 7-14 days as needed during disease period.

Brown Rot and Blight <i>Monilinia, Sclerotinia, Whetzellinia</i>	½ – 1 ½ teaspoons	Apply late Spring or at first sign of disease. Repeat every 7-14 days as needed during disease period.
Fusicladium and Venturia Leaf Scabs on: Crabapple, Hawthorn, Pear, Mountain Ash, Pyracantha, etc.	½ – 1 ½ teaspoons	Apply as buds break. Repeat every 7-14 days as needed during disease period. Effective control requires coverage during leaf expansion. Rotations and/or tank mix combinations with mancozeb (Protect), chlorothalonil or propiconazole can be utilized. Do not use fruit from treated crabapple or pear trees for food purposes.
Leaf Spots and Blights caused by: <i>Ascochyta, Blumeriella, Botrytis, Cercospora, Cocomyces, Corynespora, Curvularia, Didymellina, Entomosporium, Fabraea, Fusarium, Ramularia, Rhizoctonia, Marssonina, Mycosphaerella, Myrothecium, Phoma, Physalospora, Schizothyrium, Septoria, Sphaceloma</i>	½ – 1 ½ teaspoons	Apply when disease symptoms first appear. Repeat every 7-14 days as needed during disease period. Rotations and/or tank mix combinations with mancozeb (Protect) or chlorothalonil can be utilized.
Ovulinia Blight	½ – 1 ½ teaspoons	Apply as flowers open. Repeat every 7-14 days as needed during disease period.
Powdery Mildews <i>Erysiphe, Microsphaera, Phyllactinia, Podosphaera, Oidium, Sphaerotheca</i>	1-1 ½ teaspoons	Apply when disease first appears and repeat every 7-14 days or as needed. Rotations and/or tank mix combinations with mancozeb (Protect) or triadimefon can be utilized.
Rust Diseases caused by: <i>Puccinia, Gymnosporangium, Uromyces</i>	½ – 1 ½ teaspoons	Apply late Spring or when symptoms first appear. Repeat every 7-14 days as needed during disease period. Rotations and/or tank mix combinations with mancozeb (Protect) or chlorothalonil are recommended.
Tip Blight of Pine <i>Sphaeropsis sapinea, Diplodia pinea</i>	1 ½ - 1 ½ teaspoons	Begin application in spring when new growth starts. Make a second application just before needles emerge from the sheath and a third application 10-14 days later. Thorough coverage is essential for optimal disease control.
Twig Blights, Cankers, and Diebacks <i>Diaporthe, Kabatina, Phoma, Phomopsis</i>	1 ½ - 1 ½ teaspoons	Apply when symptoms first appear. Repeat every 7-14 days as needed during disease period.

SOIL DRENCH APPLICATION

Diseases	Rate of
----------	---------

Controlled	3336 F per gallon	Remarks
Stem, Crown, and Root Rots caused by: <i>Botrytis, Cythrodium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia</i> Black Root Rot <i>Thielaviopsis</i>	1-1 ½ teaspoons	Apply as a drench or directed spray at seeding (8 fl oz) or after transplanting (12-16 fl oz) to propagation beds, containers, pots, trays, or nursery or landscape beds at a rate to thoroughly soak the growing media through the root zone. A general guide is ¼ -3 pints per sq ft depending on the media type and depth (about 4 fl oz per 4 inch pot or 8 fl oz per 6 inch pot). Repeat every 21-28 days for adequate crop protection. Note: 3336 F does not control <i>Pythium</i> or <i>Phytophthora</i> . Tank mix combinations with metalaxyl, mefenoxam, etridiazole, propamocarb, Fosetyl-Al, or mono and dipotassium salts of phosphorous acid are required for the control of <i>Pythium</i> and <i>Phytophthora</i> .

PLANT DIP APPLICATION

Diseases Controlled	Rate of 3336 F per gallon	Remarks
Plant or Cutting Diseases caused by: <i>Botrytis, Cythrodium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia, Thielaviopsis</i>	1- 1 ½ teaspoons	Immerse plants or cuttings for 10-15 min. Remove and allow to drain. Note: Follow accepted hygiene practices to minimize the introduction and spread of water borne bacterial disease.
Bulb, Corm, and Rhizome Rots caused by: <i>Botrytis, Cythrodium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia, Thielaviopsis</i>	1-1 ½ teaspoons	Soak cleaned bulbs for 15-30 min in warm solution (80-85° F). For storage disease prevention, treat bulbs preferably within 48 hours after digging. After treatment, dry well before storing. If bulbs are for forcing, treat bulbs that have been heat-cured. Note: Follow accepted hygiene practices to minimize the introduction and spread of water borne bacterial disease.

LIMITED WARRANTY AND DISCLAIMER

CLEARY CHEMICALS LLC warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. To the extent consistent with applicable law, CLEARY CHEMICALS LLC makes no other expressed or implied warranty of fitness or merchantability or any other expressed or implied warranty. To the extent consistent with applicable law, neither CLEARY CHEMICAL nor seller shall be liable for consequential,

special or indirect damages resulting from the use or handling of this product including, but not limited to, loss of profits, business reputation, or customers, labor costs, or other expenses incurred in planting or harvesting. CLEARY CHEMICAL and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of CLEARY CHEMICALS LLC.

Amended: 9.18.08

Replaces: 5.17.04

Version: 9.18.08

NOTIFICATION

AUG 12 2009