

3336RG

ACCEPTED LABEL
 MAR 11 1997
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 1001-71

TURF GRANULAR SYSTEMIC FUNGICIDE

ACTIVE INGREDIENT

Thiophanate-methyl (dimethyl 4,4'-o-phenylenebis[3-thioallophanate]) 1.04%

INERT INGREDIENTS 98.96%

TOTAL 100.00%

Keep Out Of Reach Of Children

CAUTION

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Induce vomiting immediately by giving two glasses of water and sucking finger down throat. Call a physician. Never give anything by mouth to an unconscious person.
- IF IN EYES: Immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention if irritation develops or persists.
- IF ON SKIN: Immediately flush skin with plenty of water.
- IF INHALED: Remove victim to fresh air and check airway for blockage. If not breathing, give artificial respiration and seek additional medical help to restore breathing.

PRECAUTIONARY STATEMENTS

• HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. May cause irritation of eyes, nose, throat, and skin. Avoid breathing dust and contact with eyes and skin. Do not graze animals on treated turf. Do not feed clippings to livestock or poultry.

• ENVIRONMENTAL HAZARDS

For terrestrial (land use) only. Do not apply directly to water, where surface water is present, or to intertidal areas below the mean high water mark. Application equipment should be thoroughly cleaned after use. Do not contaminate water when disposing of equipment washwater. Do not apply, allow to drain, or flush equipment into non-target areas.

• STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal methods.

Storage: Store in a dry, climate controlled area in the original unopened container when not in use. Store in a secure area unavailable to unauthorized persons.

Pesticide Disposal: Product wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Shake empty product container to remove all possible material. When bags are empty, they may be disposed of in a sanitary landfill or by incineration or as allowed by State and local authorities. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any other requirements specific to your State or Tribe, consult the agency responsible for pesticides. **NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE SUCH AS SOD, OR FOR COMMERCIAL SEED PRODUCTION, OR FOR RESEARCH PURPOSES.** It is a VIOLATION OF FEDERAL LAW to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

Apply 3336^{RG} with ground based, hand operated or mechanical equipment sufficient to accurately deliver this product to the application site. Application equipment should be properly calibrated prior to use. Application should be water in by applicator or by homeowner with 0.1-0.25 inches of water as soon as possible to begin activation of product for disease control. If a rate range is given, use the higher rate under conditions of severe disease pressure. Contact your local State Extension service for recommendations for applications schedules appropriate to your area. No claim of compatibility with other pesticides is implied. **IMPORTANT**, if after using 3336^{RG} as recommended and the treatment is not effective, a tolerant strain of fungi may be present. Consult your local W. A. Cleary representative or your State Extension Service for advice on another suitable product for disease control.



W.A. Cleary Chemical
CORPORATION
 SOMERSET, NEW JERSEY 08875

TURF APPLICATIONS

3336^{RG} is a systemic fungicide for use on all fine turf applications on home lawns which consists of cool and warm season grasses such as, Bentgrass, Bluegrass, Bermudagrass, Centepede, Fescue, Turf Type Tall Fescue, Ryegrass, St. Augustine, Zoysia or their mixtures. 3336^{RG} is not phytotoxic to any of the above mentioned grasses when used in accordance with the label and may be used for the prevention and control of the diseases indicated below. 3336^{RG} may be applied to dry or moist leaf/soil conditions and should be activated with 1/4" to 1/2" of irrigation in the advent that natural rainfall is not sufficient. Irrigation should occur within 24 hours for best results.

RATE: 3-6 POUNDS PER THOUSAND SQUARE FEET

Dollar Spot

Apply when disease first appears. Continue at 7-14 day intervals as needed.

Large Brown Patch

Apply when disease first appears. Continue at 7-14 day intervals as needed.

Pink Snow Mold

Apply first application in November or just before grass plant have gone dormant or at first sign of disease. Best results are achieved when first application is made before turfgrass is completely dormant. Reapply as needed at first spring thaw.

Asarium Blight

Apply two applications at 10-14 day intervals beginning when disease first appears. Reapply as needed.

RATE: 9-12 POUNDS PER THOUSAND SQUARE FEET

Take All Patch

Apply first application in the fall when soil temperatures reach 60°F. Make summer applications when night temperatures are consistently above 70°F for a three day period. Repeat applications at 14 day intervals as needed for disease suppression. Water in application as described in the general information.

Bermuda Decline

For treatment of patch disease that generally occur in hot weather with high humidity. Wet conditions will encourage disease. Apply first application in the fall when soil temperatures reach 60°F. Make summer applications when night temperatures are consistently above 70°F for a three day period. Repeat applications at 14 day intervals as needed for disease suppression. Water in application as described in the general information.

Spring Dead Spot

Disease usually occurs when soil and air temperatures reach 60°F in the spring of the year. Apply first application in the fall when soil temperatures reach 60°F. Make summer applications when night temperatures are consistently above 70°F for a three day period. Repeat applications at 14 day intervals as needed for disease suppression. Water in application as described in the general information.

Foliar and Basal Rot Anthracnose

For foliar expression, apply as needed when disease first appears. Repeat as needed at 14 day intervals. For Basal Rot expression, apply as needed when night temperatures are consistently above 70°F for a three day period. Repeat applications at 7-14 day intervals as needed.

Summer Patch

Begin applications when soil temperatures reach 60°F. Apply 3 treatments at 21 day intervals. For suppression, apply when disease first appears at 10-14 day intervals as needed. Water application as described in general information.

NOTE: soil pH management has been shown to be an important cultural practice in the further management of this disease.

Necrotic Ring Spot

Apply at 14 day intervals as needed, beginning in spring when soil temperatures reach 60°F. Apply a fall application to target areas as needed.

Stripe Smut

Make 2 applications at 7-14 day intervals when disease first appears. For prevention, apply Fall of same season to reduce next years disease pressure.

SPREADER SETTINGS

Suggested Spreader Settings For 3 lb./1000 Application Rate

Spreader	Ground Speed	swath Width	Setting(s)
Central Drop			8
Cyclone Rotary	3 mph	8 feet	5
Cyclone Rotary 1B	3 mph	8 feet	3 1/2
Earthway Drop			11
Earthway Rotary	3 mph	8 feet	17
Ezee Flow			C4
Gandy			28
Greenview Drop-Spred II			3
Greenview Jet-Spred II	3 mph	8 feet	5
Greenview Jet-Spred III	3 mph	8 feet	5 1/2
Greenview Show 'N Tell			8
Lesco	3 mph	8 feet	E-F
Lely (WTR, WFR, HR, 1250)	5 mph	16 feet	4-4 1/2 III PTO at 450 RPM
Ortho Drop			7
Ortho Rotary	3 mph	8 feet	5
Red Devil Rotary	3 mph	8 feet	9
Republic Drop			7
Republic Rotary	3 mph	8 feet	5
Scott's Accu-Green			6 1/2
Scott's Drop PF1			6
Scott's Easy Green	3 mph	8 feet	28
Scott's RX7 Rotary	3 mph	8 feet	J
Scott's HBA Rotary	3 mph	8 feet	L
Scott's Rotary Retail	3 mph	8 feet	6
Scott's Speedy Green	3 mph	8 feet	5 3/4
Sears Craftsman Drop			7 1/2

LIMITED WARRANTY AND DISCLAIMER

W.A. CLEARY CHEMICAL CORPORATION warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. W.A. CLEARY CHEMICAL CORPORATION makes no other expressed or implied warranty of fitness or merchantability or any other expressed or implied warranty. In no case shall W.A. CLEARY or seller be liable for consequential, special or indirect damages resulting from the use or handling of this product including, but not limited to, loss of profits, business reputation, or customers, labor costs, or other expenses incurred in planting or harvesting. W.A. CLEARY and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of W.A. CLEARY CHEMICAL CORPORATION.



W.A. Cleary Chemical
CORPORATION

Corporate Headquarters
1049 Route 27, P.O. Box 10
Somerset, New Jersey 08875-0010

Chemical Sales and Distribution Center
178 Ridge Road, Suite A
Dayton, New Jersey 08810

Effective Date: 02/97

908-329-8399 800-524-1662

WAC215110M