

Best copy available
1007-24
02/10/1988

Agri-mycin 17

AGRICULTURAL STREPTOMYCIN

Active Ingredient: Streptomycin Sulfate
(Equivalent to 17% Streptomycin in base)

This material is a true chemical, with a chemical formula and a specific biological activity. Its use should be limited to the control of plant diseases. Seller makes no warranty, express or implied, concerning use of this product other than as indicated on the label. It is to be used in strict accordance with the label or handling not in strict accordance with the label.

KEEP...
CAUTION: FOR REPACKAGING USE ONLY

500 galls
20 oz
1 1/2 lb
2 1/2 lb
5 lb

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONS
EPA Reg No. 1007-24 EPA Est. 1007-IN-1
LOT NO. -40023-252

Net Contents: 200 lb (90.72 Kg)



Pfizer
CHEMICAL DIVISION
PFIZER INC., NEW YORK, N.Y. 10017

MADE IN U.S.A. 9.87-361 D-00

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS: Avoid contact with skin and eyes. Do not breathe dust or spray mist. Wear dust mask, chemical goggles and rubber gloves. Wash thoroughly after handling.
ENVIRONMENTAL HAZARD: Keep out of lakes, ponds and streams. Do not contaminate water by cleaning equipment or disposal of waste.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Storage: Keep tightly closed. Storage should be at a cool temperature when possible, and with minimum exposure to the atmosphere. Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Disposal: Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system.

REENTRY STATEMENT: Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

FARMWORKER NOTIFICATION: Written or oral warnings must be given to workers who are in or near treated areas or in an area about to be treated with this product. Oral warnings must be given to farm workers who are in or near treated areas or in an area about to be treated with this product. Written warnings must be given to farm workers who are in or near treated areas or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by the workers. Written warnings must be given in writing. Written warnings cannot be given if there is no reasonable way of reaching the workers. Warnings must include the following information: CAUTION. Area treated with AGRI-Mycin 17. (Name of application) Do not enter without appropriate protective clothing until sprays have dried. (Date of application) Do not enter without appropriate protective clothing until sprays have dried. (Name of application) Wash with plenty of water. For further information call _____ (Tel _____)

DISEASE AND CROP	BEST CONCENTRATION	FIRST	FOLLOW UP BEST SCHEDULE
Fire Blight (Pear, Apple, Crabapple)	200 ppm	10% bloom	Repeat at 5 day intervals until all late bloom is over (this could mean 17/15 apple bloom). (Continue to apply at 5/7 day intervals to control shoot and fruit infection. Do not apply within 30 days of harvest)
Fire Blight (Apple, Crabapple)	200 ppm	Full bloom	Apply at petal fall and late secondary bloom (continue to apply until 5/7 day interval. To maintain disease control but not later than 50 days before harvest)
Widening and Blue Mold (Apple)	200 ppm	Apply 10-15 days before setting fruit on plants in the field	Apply 10-15 days before setting fruit on plants in the field. Make three additional weekly applications at 10 lbs, 1 1/2 lbs, and 1 1/2 lbs per acre on apples. Do not apply 15 lbs per acre or later than 50 days before harvest.

Additional minimum reentry interval of 48 hours in 17 may be obtained from your local Agricultural Extension Agent or State Department of Agriculture.

DISEASE AND CROP	RECOMMENDED CONCENTRATION	FIRST SPRAY	FOLLOW UP SPRAY SCHEDULE
Fire Blight (Pear, Apple, Crabapple)	200 ppm	10% bloom	Repeat at 5 day intervals until all late bloom is over (this could mean 17/15 apple bloom). (Continue to apply at 5/7 day intervals to control shoot and fruit infection. Do not apply within 30 days of harvest)
Fire Blight (Apple, Crabapple)	200 ppm	Full bloom	Apply at petal fall and late secondary bloom (continue to apply until 5/7 day interval. To maintain disease control but not later than 50 days before harvest)
Widening and Blue Mold (Apple)	200 ppm	Apply first spray when varieties are in full stage when leafy twigs appear	Apply first spray 10 days before harvest (more spray sprays after petal fall until 5/7 day interval). Do not apply within 30 days of harvest.
Fire Blight (Apple)	200 ppm	Apply first spray when varieties are in full stage when leafy twigs appear	Apply first spray 10 days before harvest (more spray sprays after petal fall until 5/7 day interval). Do not apply within 30 days of harvest.
Fire Blight (Apple)	200 ppm	Apply first spray when varieties are in full stage when leafy twigs appear	Apply first spray 10 days before harvest (more spray sprays after petal fall until 5/7 day interval). Do not apply within 30 days of harvest.
Widening and Blue Mold (Apple)	200 ppm	Apply first spray when varieties are in full stage when leafy twigs appear	Apply first spray 10 days before harvest (more spray sprays after petal fall until 5/7 day interval). Do not apply within 30 days of harvest.
Widening and Blue Mold (Apple)	200 ppm	Apply first spray when varieties are in full stage when leafy twigs appear	Apply first spray 10 days before harvest (more spray sprays after petal fall until 5/7 day interval). Do not apply within 30 days of harvest.
Widening and Blue Mold (Apple)	200 ppm	Apply first spray when varieties are in full stage when leafy twigs appear	Apply first spray 10 days before harvest (more spray sprays after petal fall until 5/7 day interval). Do not apply within 30 days of harvest.
Widening and Blue Mold (Apple)	200 ppm	Apply first spray when varieties are in full stage when leafy twigs appear	Apply first spray 10 days before harvest (more spray sprays after petal fall until 5/7 day interval). Do not apply within 30 days of harvest.

