

# FOR DAIRY AND BEEF CATTLE

## DIRECTIONS FOR USE

- FOR DIRECT APPLICATION TO DAIRY CATTLE AND BEEF CATTLE, FOR CONTROL OF HORNFLIES: Apply 2 heaping tablespoonfuls (2 ounces) to the backs and flanks of infested animals. Rub lightly into hair. Repeat application no more than once in every 3 week period, if necessary.
- FOR CONTROL OF LICE AND TICKS ON BEEF CATTLE: Use as above. Treat animals thoroughly. Repeat treatments as necessary.
- AS AN AID FOR THE CONTROL OF FACE FLIES ON BEEF CATTLE: Dust the faces of cattle thoroughly. Frequent repeated applications may be necessary.
- FOR CONTROL OF HOG LICE: Apply to hogs, pens, and litter. One treatment may be sufficient. Treat animals thoroughly. Repeat in 10 days or thereafter as needed.
- LICE AND KEDS ON SHEEP: Treat animals thoroughly, preferably following shearing. Repeat as necessary.

Harmful if swallowed. Do not apply within 5 hours of milking or during milking. Do not contaminate feed or water. Avoid breathing of dust. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Do not use on poultry or in poultry houses. Do not reuse container. Destroy when empty. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.



E P A Reg. No. 1007-39

MADE IN U.S.A. 8



# CATTLE

# DRY ING

## FOR DAIRY AND Contains Malathion a

Controls Hornflies (and Hornflies, Lice and Ticks)

Active Ingredients:  
Methoxychlor, technical\*  
Malathion (O, O-dimethyl dithiophosphorothioic diethyl mercaptosuccinate)  
Inert Ingredients:

\*Equivalent to 4.4% 2, 2-bis (p-methoxyphenyl) ethane and 0.6% related reaction products

**CAUTION:** Keep out of reach of children.

SEE BACK PANEL FOR ADDITIONAL INFORMATION

NET WEIGHT: 1.5 LBS (680g)



# RY AND BEEF CATTLE

## RECTIONS FOR USE

ION TO DAIRY CATTLE AND BEEF CATTLE, FOR  
IES: Apply 2 heaping tablespoons (2 ounces) to  
of infested animals. Rub lightly into hair. Repeat  
than once in every 3 week period, if necessary.

AND TICKS ON BEEF CATTLE: Use as above. Treat  
Repeat treatments as necessary.

ONTROL OF FACE FLIES ON BEEF CATTLE: Dust  
oroughly. Frequent repeated applications may be

LICE: Apply to hogs, pens, and litter. One treat-  
ent. Treat animals thoroughly. Repeat in 10 days  
led.

EEP: Treat animals thoroughly, preferably follow-  
as necessary.

wallowed. Do not apply within 5 hours of milking  
Do not contaminate feed or water. Avoid  
oid contact with skin; wash thoroughly after using.  
of feed and foodstuffs. Do not use on poultry or in  
ot reuse container. Destroy when empty.  
to fish. Keep out of lakes, streams or ponds. Do  
ter by cleaning of equipment or disposal of wastes.



# CATTLE DUST DRY INSECTICIDE

## FOR DAIRY AND BEEF CATTLE

Contains Malathion and Methoxychlor

Controls Hornflies on Dairy Cattle  
and Hornflies, Lice and Ticks on Beef Cattle

Active Ingredients:	
Methoxychlor, technical* .....	5.0%
Malathion (O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate) .....	4.0%
Inert Ingredients: .....	91.0%

\*Equivalent to 4.4% 2, 2-bis (p-methoxyphenyl) - 1, 1, 1- trichloro-ethane and 0.6% related reaction products.

**CAUTION:** Keep out of the reach of children.

SEE BACK PANEL FOR ADDITIONAL CAUTIONS

**NET WEIGHT: 4 LBS.**



EPA Reg. No. 1007-39

MADE IN U.S.A. 8



Distributed by  
**AGRICULTURAL DIVISION**  
NEW YORK, N.Y. 10017