

places. Remove animals from buildings before spraying. Do not spray feed, forage, drinking water or feed containers. If drinking water or feed containers become contaminated, flush immediately with water.

**INSECTICIDAL PAINT:** Use 1.25% lindane prepared by adding 1 pint Lindane Compound to 1 gallon of water. Paint on and around side-walls, doors, windows, window screens, beams, baseboards, cracks, crevices and other hiding places. This serves as a spot treatment for these surfaces. Avoid contact with skin. Not for use in the home. This concentration may blister paint.

**FLIES ON MANURE PILES, REFUSE HEAPS:** Spray with 0.125% lindane prepared by adding 1 pint Lindane Compound to 10 gallons of water. Wet thoroughly every 3 or 4 weeks, or at 3 to 4 day intervals if manure or refuse is being added daily.

**CHIGGERS AND TICKS ON LAWNS:** Spray thoroughly with 0.06% lindane prepared by adding 1 pint Lindane Compound to 21 gallons of water.

**AGRICULTURAL INSECTS:** See leaflet.

**WARNING:** Harmful if swallowed. May be absorbed through skin. Avoid breathing spray mist and skin contact. In case of contact, wash immediately with soap and water. In case of contact with eyes, flush with water and get medical attention.

**CAUTION:** Do not use more than 0.025% lindane in finished spray on calves and other young livestock. Do not treat animals under three months of age. Do not treat sows within two weeks of farrowing nor for at least three weeks thereafter. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals. Do not use on dairy animals. Do not spray on any animal within 30 days of slaughter. Do not use on household pets or humans. Avoid contamination of feed and foodstuffs. Do not use or store near heat or open flame. Do not use this product in poultry houses, dairy barns or milk rooms.

See leaflet for additional cautions.

**CONTAINER DISPOSAL:** Do not reuse empty container. Destroy by crushing. Bury or discard in a safe place.

USDA Reg. No. 1007-45

Pfizer



For the control of certain insects on beef cattle and other livestock; and for the control of flies and certain other insects in livestock buildings.

To be diluted with water.

ACTIVE INGREDIENT: Lindane . . . . . 10.68% w/w  
INERT INGREDIENTS: . . . . . 89.32% w/w

**WARNING:** Keep out of reach of children.  
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

**NET CONTENTS: 16 FL. OZ. (1 PINT)**



Distributed by  
**AGRICULTURAL DIVISION**  
CHAS. PFIZER & CO., INC., NEW YORK, N.Y. 10017



**DIRECTIONS: MEASURE ACCURATELY AND  
STIR THOROUGHLY JUST BEFORE USING**

**HORNFLIES ON BEEF CATTLE:**

Spray animals thoroughly with 0.025% lindane prepared by adding 1 tablespoonful Lindane Compound to 1½ gallons of water; or 1 pint to 50 gallons. Repeat every 3 or 4 days or as often as necessary.

**LICE AND TICKS ON BEEF CATTLE, SHEEP, GOATS:** Spray animals with 0.03% lindane prepared by adding 2 tablespoonfuls Lindane Compound to 2½ gallons of water; or 1 pint to 42 gallons. Skin and hair of sheep and goats must be thoroughly wet. Repeat as necessary.

**LICE ON HOGS:** Spray hogs with 0.06% lindane prepared by adding 4 tablespoonfuls Lindane Compound to 2½ gallons of water; or 1 pint to 21 gallons. Wet skin thoroughly. Repeat as necessary. Do not apply to young animals.

**SARCOPTIC MANGE ON BEEF CATTLE AND HOGS, AND PSOROPTIC MANGE ON SHEEP AND GOATS:** Use 0.03% lindane prepared by adding 2 tablespoonfuls Lindane Compound to 2½ gallons of water; or 1 pint to 42 gallons. On beef cattle 2 gallons of the diluted spray per adult animal should be used at the rate of two sprays one week apart. Wet hogs thoroughly, including inside of ears. Pens and shelters should be sprayed also to prevent reinestation.

**RESIDUAL CONTROL OF FLIES, ROACHES, BOXELDER BUGS, SPIDERS, ANTS AND MOSQUITOES IN LIVESTOCK BUILDINGS:** For rough surfaces spray with 0.3% lindane prepared by adding 6 tablespoonfuls Lindane Compound to 1 gallon of water; or 1 pint to 4 gallons, and for smooth surfaces or for longer residual control use 0.5% lindane prepared by adding 7 tablespoonfuls Lindane Compound to ½ gallon of water, or 1 pint to 2½ gallons, applying 1 gallon for each 500 sq. ft. of surface, or until thoroughly wet, about to run off. For crawling insects apply thoroughly to cracks, crevices, baseboards and other hiding

ACCEPTED  
FEB 26 1970  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTRATION  
ED UNDER NO. 1007-45

01-1459-00-6



60-1460-00-3



## LINDANE COMPOUND

### Directions for Control of AGRICULTURAL INSECTS

**FOLLOW DIRECTIONS CLOSELY, MEASURE ACCURATELY, AND STIR EMULSION THOROUGHLY JUST BEFORE USING.**

Always mix with required amount of water. Measure carefully and mix thoroughly using 0.025% lindane in the final spray. Prepare by mixing 2 tablespoonfuls Lindane Compound with 3 gallons of water, or 1 pint with 50 gallons. See individual directions below and follow closely. In general, application should be made at first sign of infestation and repeated as necessary. Plants should not be sprayed when in full blossom. Consult your Agricultural Extension Service or State Experiment Station for recommendations as to timing and number of applications. Observe warnings on label on bottle.

**CAUTION:** Do not use Lindane in fields to be planted within 2 years with peanuts or root crops as it may adversely affect their flavor.

This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds.

#### VEGETABLES and BERRIES

**ASPARAGUS:** Asparagus Beetle. Do not treat cutting asparagus. Post-harvest application only. Maximum dosage: 280 gallons of 0.025% spray (0.6 pound actual Lindane) per acre.

**BEANS:** Aphids, Cucumber Beetle, Leafhoppers, Thrips, White Fly. Do not apply after edible parts start to form. Dosage: 100 gallons of 0.025% spray (0.2 pound actual Lindane) per acre. (Maximum dosage 0.6 pound actual Lindane per acre.)

**BR** **QL:** Aphids. Apply to soil around base of plants. Spray plants several times at 5 to 7 day intervals beginning when Aphids first appear. Do not apply after edible parts start to form. Dosage: 100 gallons of 0.025% spray (0.2 pound actual Lindane) per acre. (Maximum dosage 0.75 pound actual Lindane per acre.)

**CABBAGE:** Aphids, Cabbage Maggot, Flea Beetles, Tarnished Plant Bug, Thrips. Treat plant and apply to soil around base of plants. Do not apply after edible parts begin to form. Dosage: 100 gallons of 0.025% spray (0.2 pound actual Lindane) per acre. (Maximum dosage 0.75 pound actual Lindane per acre.)

**CANTALOUPE:** Aphids, Thrips. Dosage: 100 gallons of 0.025% spray (0.2 pound actual Lindane) per acre. (Maximum dosage 0.5 pound actual Lindane per acre.)

**CELERY:** Aphids, Climbing Cutworms, Tarnished Plant Bug, Thrips. Do not apply after plants are half grown or begin to bunch. Maximum dosage: 190 gallons of 0.025% spray (0.4 pound actual Lindane) per acre.

**CUCUMBERS, SQUASH:** Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Leaf Miners, Lygus Bug, Melon Worm, Pickle Worm, Squash Bug, Squash Vine Borer, Thrips. Spray under leaves also. Do not apply within one day prior to picking. Late application may result in some reduction in quality of squash. Dosage: 150 gallons of 0.025% Spray (0.3 pound actual Lindane) per acre. (Maximum dosage 0.75 pound actual Lindane per acre.) Do not apply this dosage after the blossoms open.

**EUCALYPTUS:** Aphids, Lace Wing Bug, Leaf Miners, Thrips. Do not apply after edible parts start to form. Maximum dosage: 240 gallons of 0.025% spray (0.5 pound actual Lindane) per acre.

**KALE, SPINACH:** Aphids, Armyworms, Cabbage Worms, Corn Earworm, Cutworm, Leaf Miners, Loopers, Potato Beetle, Flea Beetle, Webworm. Apply to soil around base of plants. Do

#### VEGETABLES and BERRIES (continued)

**STRAWBERRIES:** Aphids, Leafrollers, Pea Weevil, Spittle Bug, Strawberry Weevil. Do not apply after fruit starts to form. Maximum dosage: 72 gallons of 0.025% spray (0.15 pounds actual Lindane) per acre.

**SWEET CORN:** Aphids. Application should be made before silks appear. Do not feed treated forage to dairy animals or to animals being finished for slaughter. Maximum dosage: 2400 gallons of 0.025% spray (5.0 pounds actual Lindane) per acre.

**TOMATOES:** Aphids. Do not apply after fruit starts to form. Maximum dosage: 360 gallons of 0.025% spray (0.75 pounds actual Lindane) per acre.

#### TREES and SHRUBS

**APPLES, PEARS:** Aphids, Leafhoppers, Thrips. Do not apply within 60 days of harvest. Dosage: Use 0.025% spray (1/4 pound actual Lindane per 100 gallons). (Maximum dosage: 3.6 pounds actual Lindane per acre.)

**ARBORVITAE, CONIFER, CYPRESS, PINES, SPRUCES:** Aphids, Bagworms, Cypress Webbers.

**CHERRIES:** Aphids, Leafrollers, Thrips. Do not apply after fruit starts to form. Maximum dosage: 48 gallons of 0.025% spray (1.0 pound actual Lindane) per acre.

**GRAPEFRUIT, LEMONS, LIMES, ORANGES, TANGERINES:** Aphids, Thrips. Do not apply after fruit starts to form. Maximum dosage: 950 gallons of 0.025% spray (2 pounds actual Lindane) per acre.

**ELMS:** Woolly Elm Aphid, Woolly Elm Aphid.

**FIRS, LARCHES, NORWAY AND BLUE SPRUCES:** Larch Woolly Aphid, Pine Butterfly, Saw Fly Larvae, Twig Aphid.

**GENISTA:** Aphids.

**HOLLY:** Aphids, Thrips.

**NECTARINE:** Aphids, Leafrollers, Thrips. Do not apply within 60 days of harvest. Maximum dosage: 960 gallons of 0.025% spray (2.0 pounds actual Lindane) per acre.

**OLEANDERS:** Aphids.

**PEACHES, PLUMS, PRUNES:** Aphids, Leafrollers, Thrips. Do not apply within 60 days of harvest. Maximum dosage: 720 gallons of 0.025% spray (1.5 pounds actual Lindane) per acre.

#### FLOWERS and ORNAMENTALS

**BULBS (Amaryllis, Cyclamen, Daffodil, Gladiolus, Hyacinth, Iris, Lilies, Narcissus, Tulips):** Aphids, Bulb Fly, European Corn Borer, Iris Borer, Thrips. Apply when insects appear, and repeat as necessary.

**FLOWERS (Asters, Begonias, Carnations, Chrysanthemums, Coleus, Daisies, Delphinium, Forget-me-nots, Fuchsias, Gardenias, Geraniums, Hibiscus, Larkspur, Nasturtiums, Pansies, Pelargoniums, Peonies, Petunias, Pinks, Snapdragon, Sweet Peas, Zinnias):** Aphids, Blister Beetles, Cankerworms, Cucumber Beetles, Flea Beetle, Leafhoppers, Leaf Miners, Plant Bugs, Thrips.

**IVY:** Ivy Aphid.

**ORNAMENTAL FLOWERS (Azaleas, Camellias, Heather, Lilacs, Rhododendrons, Roses):** Aphids, Blackvine Weevil, Cucumber Beetles, Curculio, Fuller's Rose Weevil, Leaf Miner, Strawberry Weevil, Thrips, White Fly.

**SUCCULENTS (Agaves, Aloes, Cactus, Crassula, Mesembryanthemums, Sedums):** Aphids.

**TOYON, RED BERRIES:** Aphids, Thrips, White Fly.

**CAUTION:** Do not use Lindane in fields to be planted within 2 years with peanuts or root crops as it may adversely affect their flavor. This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds.

**VEGETABLES and BERRIES**

**ASPARAGUS:** Asparagus Beetle. Do not treat cutting asparagus. Post-harvest application only. Maximum dosage: 280 gallons of 0.025% spray (0.6 pound actual Lindane) per acre.

**BEANS:** Aphids, Cucumber Beetle, Leafhoppers, Thrips, White Fly. Do not apply after edible parts start to form. Dosage: 100 gallons of 0.025% spray (0.2 pound actual Lindane) per acre. (Maximum dosage 0.6 pound actual Lindane per acre.)

**BR** **OLI:** Aphids. Apply to soil around base of plants. Spray plants several times at 5 to 7 day intervals beginning when Aphids first appear. Do not apply after edible parts start to form. Dosage: 100 gallons of 0.025% spray (0.2 pound actual Lindane) per acre. (Maximum dosage 0.75 pound actual Lindane per acre.)

**CABBAGE:** Aphids, Cabbage Maggot, Flea Beetles, Tarnished Plant Bug, Thrips. Treat plant and apply to soil around base of plants. Do not apply after edible parts begin to form. Dosage: 100 gallons of 0.025% spray (0.2 pound actual Lindane) per acre. (Maximum dosage 0.75 pound actual Lindane per acre.)

**CANTALOUPE:** Aphids, Thrips. Dosage: 100 gallons of 0.025% spray (0.2 pound actual Lindane) per acre. (Maximum dosage 0.5 pound actual Lindane per acre.)

**CELERY:** Aphids, Climbing Cutworms, Tarnished Plant Bug, Thrips. Do not apply after plants are half grown or begin to bunch. Maximum dosage: 190 gallons of 0.025% spray (0.4 pound actual Lindane) per acre.

**CUCUMBERS, SQUASH:** Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Leaf Miners, Lygus Bug, Melon Worm, Pickle Worm, Squash Bug, Squash Vine Borer, Thrips. Spray under leaves also. Do not apply within one day prior to picking. Late application may result in some reduction in quality of squash. Dosage: 150 gallons of 0.025% Spray (0.3 pound actual Lindane) per acre. (Maximum dosage 0.75 pound actual Lindane per acre. Do not apply this dosage after the blossoms open.)

**EG** **NT:** Aphids, Lace Wing Bug, Leaf Miners, Thrips. Do not apply after edible parts start to form. Maximum dosage: 240 gallons of 0.025% spray (0.5 pound actual Lindane) per acre.

**KALE, SPINACH:** Aphids, Armyworms, Cabbage Worms, Corn Earworm, Cutworm, Leaf Miners, Loopers, Potato Beetle, special Flea Beetle, Webworm. Apply to soil around base of plants. Do not apply to kale after edible parts start to form. Do not apply to spinach after seedling stage. Maximum dosage: 360 gallons of 0.025% spray (0.75 pound actual Lindane) per acre.

**ONIONS:** Aphids, Onion Maggot, Thrips. Make first application when thrips average 5 to 6 per plant, or when plants are 2 to 3 inches tall. Do not spray green or spring onions. Maximum dosage: 480 gallons of 0.025% spray (1.0 pound actual Lindane) per acre.

**PEAS:** Aphids, Cucumber Beetles. Do not apply after pods start to form. Dosage: 100 gallons of 0.025% spray (0.2 pound actual Lindane) per acre. (Maximum dosage 0.3 pound actual Lindane per acre.)

**PEPPERS:** Aphids, Armyworms, Leaf Miners, Thrips. Do not apply after edible parts start to form. Maximum dosage: 280 gallons of 0.025% spray (0.6 pound actual Lindane) per acre.

imum dosage: 950 gallons of 0.025% spray (2 pounds actual Lindane) per acre.

**ELMS:** Woolly Apple Aphids, Woolly Elm Aphid.

**FIRS, LARCHES, NORWAY AND BLUE SPRUCES:** Larch Woolly Aphid, Pine Butterfly, Saw Fly Larvae, Twig Aphid.

**GENISTA:** Aphids.

**HOLLY:** Aphids, Thrips.

**NECTARINE:** Aphids, Leafrollers, Thrips. Do not apply within 60 days of harvest. Maximum dosage: 960 gallons of 0.025% spray (2.0 pounds actual Lindane) per acre.

**OLEANDERS:** Aphids.

**PEACHES, PLUMS, PRUNES:** Aphids, Leafrollers, Thrips. Do not apply within 60 days of harvest. Maximum dosage: 720 gallons of 0.025% spray (1.5 pounds actual Lindane) per acre.

**FLOWERS and ORNAMENTALS**

**BULBS (Amaryllis, Cyclamen, Daffodil, Gladiolus, Hyacinth, Iris, Lilies, Narcissus, Tulips):** Aphids, Bulb Fly, European Corn Borer, Iris Borer, Thrips. Apply when insects appear, and repeat as necessary.

**FLOWERS (Asters, Begonias, Carnations, Chrysanthemums, Coleus, Daisies, Delphinium, Forget-me-nots, Fuchsias, Gardenias, Geraniums, Hibiscus, Larkspur, Nasturtiums, Pansies, Pelargoniums, Peonies, Petunias, Pinks, Snapdragons, Sweet Peas, Zinnias):** Aphids, Blister Beetles, Cankerworms, Cucumber Beetles, Flea Beetle, Leafhoppers, Leaf Miners, Plant Bugs, Thrips.

**IVY:** Ivy Aphid.

**ORNAMENTAL FLOWERS (Azaleas, Camellias, Heather, Lilacs, Rhododendrons, Roses):** Aphids, Blackvine Weevil, Cucumber Beetles, Curculio, Fuller's Rose Weevil, Leaf Miner, Strawberry Weevil, Thrips, White Fly.

**SUCCULENTS (Agaves, Aloes, Cactus, Cruassula, Mesembryanthemums, Sedums):** Aphids.

**TOYON, RED BERRIES:** Aphids, Thrips, White Fly.

**AGRICULTURAL DIVISION**  
**CHAS. PFIZER & CO., INC.**  
NEW YORK, NEW YORK 10017

Printed in U.S.A.  
Revised Oct. 1969

60-1460-00-4

10/21/69

feed containers become contaminated, flush immediately with water.

**INSECTICIDAL PAINT:** Use 1.25% lindane prepared by adding 1 pint Lindane Compound to 1 gallon of water. Paint on and around side-walls, doors, windows, window screens, beams, baseboards, cracks, crevices and other hiding places. This serves as a spot treatment for these surfaces. Avoid contact with skin. Not for use in the home. This concentration may blister paint.

**FLIES ON MANURE PILES, REFUSE HEAPS:** Spray with 0.125% lindane prepared by adding 1 pint Lindane Compound to 10 gallons of water. Wet thoroughly every 3 or 4 weeks, or at 3 to 4 day intervals if manure or refuse is being added daily.

**CHIGGERS AND TICKS ON LAWNS:** Spray thoroughly with 0.06% lindane prepared by adding 1 pint Lindane Compound to 21 gallons of water.

**AGRICULTURAL INSECTS:** See leaflet.

**WARNING:** Harmful if swallowed. May be absorbed through skin. Avoid breathing spray mist and skin contact. In case of contact, wash immediately with soap and water. In case of contact with eyes, flush with water and get medical attention.

**CAUTION:** Do not use more than 0.025% lindane in finished spray on calves and other young livestock. Do not treat animals under three months of age. Do not treat sows within two weeks of farrowing nor for at least three weeks thereafter. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals. Do not use on dairy animals. Do not spray on any animal within 30 days of slaughter. Do not use on household pets or humans. Avoid contamination of feed and foodstuffs. Do not use or store near heat or open flame. Do not use this product in poultry houses, dairy barns or milk rooms.

See leaflet for additional cautions.

**CONTAINER DISPOSAL:** Do not reuse empty container. Destroy by crushing. Bury or discard in a safe place.

USDA Reg. No. 1007-45

**Pfizer**

# LINDANE Compound

For the control of certain insects on beef cattle and other livestock; and for the control of flies and certain other insects in livestock buildings.

To be diluted with water.

**ACTIVE INGREDIENT:** Lindane ..... 10.68% w/w  
**INERT INGREDIENTS:** ..... 89.32% w/w

**WARNING:** Keep out of reach of children.  
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

**NET CONTENTS: 32 FL. OZ. (1 QT.)**

**Pfizer**

Distributed by  
**AGRICULTURAL DIVISION**  
CHAS. PFIZER & CO., INC., NEW YORK, N.Y. 10017



**DIRECTIONS: MEASURE ACCURATELY AND STIR THOROUGHLY JUST BEFORE USING**

**HORNFLIES ON BEEF CATTLE:**

Spray animals thoroughly with 0.025% lindane prepared by adding 1 tablespoonful Lindane Compound to 1½ gallons of water; or 1 pint to 50 gallons. Repeat every 3 or 4 days or as often as necessary.

**LICE AND TICKS ON BEEF CATTLE, SHEEP, GOATS:** Spray animals with 0.03% lindane prepared by adding 2 tablespoonfuls Lindane Compound to 2½ gallons of water; or 1 pint to 42 gallons. Skin and hair of sheep and goats must be thoroughly wet. Repeat as necessary.

**LICE ON HOGS:** Spray hogs with 0.06% lindane prepared by adding 4 tablespoonfuls Lindane Compound to 2½ gallons of water; or 1 pint to 21 gallons. Wet skin thoroughly. Repeat as necessary. Do not apply to young animals.

**SARCOPTIC MANGE ON BEEF CATTLE AND HOGS, AND PSOROPTIC MANGE ON SHEEP AND GOATS:** Use 0.03% lindane prepared by adding 2 tablespoonfuls Lindane Compound to 2½ gallons of water; or 1 pint to 42 gallons. On beef cattle 2 gallons of the diluted spray per adult animal should be used at the rate of two sprays one week apart. Wet hogs thoroughly, including inside of ears. Pens and shelters should be sprayed also to prevent reinfestation.

**RESIDUAL CONTROL OF FLIES, ROACHES, BOXELDER BUGS, SPIDERS, ANTS AND MOSQUITOES IN LIVESTOCK BUILDINGS:** For rough surfaces spray with 0.3% lindane prepared by adding 8 tablespoonfuls Lindane Compound to 1 gallon of water; or 1 pint to 2½ gallons, and for smooth surfaces or for longer residual control use 0.5% lindane prepared by adding 7 tablespoonfuls Lindane Compound to ½ gallon of water; or 1 pint to 2½ gallons, applying 1 gallon for each 500 sq. ft. of surface, or until thoroughly wet, about to run off. For crawling insects apply thoroughly to cracks, crevices, baseboards and other hiding places. Remove animals from buildings before spraying. Do not spray feed, forage, drinking water or feed containers. If drinking water or

MADE IN U.S.A. 6



(Directions Continued)

other hiding places. Remove animals from buildings before spraying. Do not spray feed, forage, drinking water or feed containers. If drinking water or feed containers become contaminated, flush immediately with water.

**INSECTICIDAL PAINT:** Use 1.25% lindane prepared by adding 1 pint Lindane Compound to 1 gallon of water. Paint on and around sidewalls, doors, windows, window screens, beams, baseboards, cracks, crevices and other hiding places. This serves as a spot treatment for these surfaces. Avoid contact with skin. Not for use in the home. This concentration may blister paint.

**FLIES ON MANURE PILES, REFUSE HEAPS:** Spray with 0.125% lindane prepared by adding 1 pint Lindane Compound to 10 gallons of water. Wet thoroughly every 3 or 4 weeks, or at 3 to 4 day intervals if manure or refuse is being added daily.

**CHIGGERS AND TICKS ON LAWNS:** Spray thoroughly with 0.06% lindane prepared by adding 1 pint Lindane Compound to 21 gallons of water.

**AGRICULTURAL INSECTS:** See leaflet.

**WARNING:** Harmful if swallowed. May be absorbed through skin. Avoid breathing spray mist and skin contact. In case of contact, wash immediately with soap and water. In case of contact with eyes, flush with water and get medical attention.

**CAUTION:** Do not use more than 0.025% lindane in finished spray on calves and other young livestock. Do not treat animals under three months of age. Do not treat sows within two weeks of farrowing nor for at least three weeks thereafter. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals. Do not use on dairy animals. Do not spray on any animal within 30 days of slaughter. Do not use on household pets or humans. Avoid contamination of feed and foodstuffs. Do not use or store near heat or open flame. Do not use this product in poultry houses, dairy barns or milk rooms. See leaflet for additional cautions.

**CONTAINER DISPOSAL:** Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

USDA Reg. No. 1007-45  
MADE IN U.S.A. 6

Pfizer

LINDANE

Compound

For the control of certain insects on beef cattle and other livestock; and for the control of flies and certain other insects in livestock buildings.

To be diluted with water.

ACTIVE INGREDIENT:

Lindane..... 10.68% w/w

INERT INGREDIENTS:..... 89.32% w/w

**WARNING:** Keep out of reach of children.

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

NET CONTENTS: ONE FL. GAL.

Pfizer

Distributed by

AGRICULTURAL DIVISION  
CHAS. PFIZER & CO., INC., NEW YORK, N.Y. 10017



**DIRECTIONS:**  
MEASURE ACCURATELY  
AND STIR THOROUGHLY  
JUST BEFORE USING

**HORNFLIES ON BEEF CATTLE:**

Spray animals thoroughly with 0.025% lindane prepared by adding 1 tablespoonful Lindane Compound to 1½ gallons of water; or 1 pint to 50 gallons. Repeat every 3 or 4 days or as often as necessary.

**LICE AND TICKS ON BEEF CATTLE, SHEEP, GOATS:** Spray animals with 0.03% lindane prepared by adding 2 tablespoonfuls Lindane Compound to 2½ gallons of water; or 1 pint to 42 gallons. Skin and hair of sheep and goats must be thoroughly wet. Repeat as necessary.

**LICE ON HOGS:** Spray hogs with 0.06% lindane prepared by adding 4 tablespoonfuls Lindane Compound to 2½ gallons of water; or 1 pint to 21 gallons. Wet skin thoroughly. Repeat as necessary. Do not apply to young animals.

**SARCOPTIC MANGE ON BEEF CATTLE AND HOGS, AND PSOROPTIC MANGE ON SHEEP AND GOATS:** Use 0.03% lindane prepared by adding 2 tablespoonfuls Lindane Compound to 2½ gallons of water; or 1 pint to 42 gallons. On beef cattle 2 gallons of the diluted spray per adult animal should be used at the rate of two sprays one week apart. Wet hogs thoroughly, including inside of ears. Pens and shelters should be sprayed also to prevent reinfestation.

**RESIDUAL CONTROL OF FLIES, ROACHES, BOXELDER BUGS, SPIDERS, ANTS AND MOSQUITOES IN LIVESTOCK BUILDINGS:**

For rough surfaces spray with 0.3% lindane prepared by adding 8 tablespoonfuls Lindane Compound to 1 gallon of water; or 1 pint to 4 gallons, and for smooth surfaces or for longer residual control use 0.5% lindane prepared by adding 7 tablespoonfuls Lindane Compound to ½ gallon of water; or 1 pint to 2½ gallons, applying 1 gallon for each 500 sq. ft. of surface, or until thoroughly wet, about to run off. For crawling insects apply thoroughly to cracks, crevices, baseboards and  
(Continued on other panel)

01-1457-00-6